Zeitschrift:	Acta Tropica
Herausgeber:	Schweizerisches Tropeninstitut (Basel)
Band:	25 (1968)
Heft:	1
Artikel:	Miscellaneum : On the trypanosome-infection rate of "Glossina morsitans" in the Ulanga District (Tanzania)
Autor:	Geigy, R. / Kauffmann, M. / Diehl, P.
DOI:	https://doi.org/10.5169/seals-311529

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. <u>Siehe Rechtliche Hinweise.</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. <u>Voir Informations légales.</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. <u>See Legal notice.</u>

Download PDF: 14.05.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Miscellaneum

On the Trypanosome-Infection Rate of *Glossina morsitans* in the Ulanga District (Tanzania)

By R. GEIGY, M. KAUFFMANN and P. DIEHL

Swiss Tropical Institute, Basle, and Rural Aid Centre, Ifakara (Tanzania)

To further the knowledge of the epidemiological situation regarding trypanosomiasis in the Ulanga District (Tanzania), a survey was carried out in 1967 on the infection rate of freshly caught wild *Glossina morsitans*. For comparison, tsetse flies were examined a) in an area around Ifakara, where a lot of game and great numbers of tsetse flies are to be found, but no human trypanosomiasis has occurred so far and b) in the endemic region around Kilosa kwa Mpepo, where each year between 20 to 70 cases of sleeping sickness are reported and where we had found a rather high incidence of trypanosomes in wild mammals, namely 34.3% from the middle of October to the middle of December 1966 (GEIGY et al., 1967).

The Ifakara area was studied between the middle of August and the end of September, the Kilosa kwa Mpepo area in September and October 1967. A planned simultaneous survey of trypanosome infection in wild animals in the area around Ifakara had to be a abandoned because of the weather conditions: after prolonged rains the grass proved to be too high for hunting during the time at our disposal.

Material and methods

The tsetse survey of the Ifakara region was carried out from the Rural Aid Centre, whereas we installed a field laboratory for the work around Kilosa kwa Mpepo at Ngoheranga Mission, 18 miles north of Kilosa kwa Mpepo, as last year.

Males and females of G. morsitans only, the species predominant in both areas, were examined. To begin with, we tried to compare the age and sex ratio from catches off a bait cow and such caught on a Landrover, slowly driving over a track of about 2 miles through wooded savannah (Brachystegia woodland — miombo). As we were able to catch many more individuals in a much shorter time by the Landrover method and the ratio of males and females as well as of the age groups were similar with both methods, we gave up using bait animals very soon. Specially so, as the cow we used got soon infected with Trypanosoma congolense as well as Theileria parva and had to be slaughtered. As the number of male flies exceeded that of females considerably, we applied JACKSON'S (1947) wing fray method, using the wear of the hind margin of the wings, for age grading into 6 groups. For females, we started with SAUNDERS' (1962) ovarian method, but had to give it up as too time consuming. After consultation with SAUNDERS (personal communication) and after comparing our own results with both methods on a number of individuals, we decided, that the use of the wing fray method for females too would be sufficient for our purposes. Although, one still has to consider the fact, that as the male tsetse are more active than the females, one must suppose that females in the same wing fray categories are older than the corresponding males (JACKSON, 1947). Males as well as females were dissected for assessing the infection rate within 10 to 36 hours after being



caught. The flies were kept in a cool box (10 to 15° C) up to the time of dissection. Labrum, hypopharynx, salivary glands and gut were examined separately. In all cases, where the gut as well as the hypopharynx were showing trypanosomes, the entire proboscis was injected i.p. into a white mouse. The same was done with infected salivary glands. After JORDAN (1964) and HARLEY (1966) infections involving the salivary glands are of the brucei-type, those confined to the proboscis of the vivax- and those involving gut as well as proboscis supposed to be of the congolense-type. Infections found in the gut only, may be either brucei- or congolense-type. The determination of vivax- and mature brucei-type are clear cut. But as regards the species involved in gut-proboscis infections, these may be either congolense-type or a mixture of either brucei or congolense with vivax. Furthermore, in an area, where most of the trypanosome carrying wild animals show mixed infections, one must expect a high percentage of mixed infections in the tsetse fly as well. We therefore preferred to classify our results according to the localisation of the trypanosomes in the fly. Where metacyclic forms were present in such infections, the follow up of the infections in mice may clear the situation.

Results

Most of the flies were caught between 8.30 and 10.00 in the morning, single catches varying between 16 and 189 individuals. In the Ifakara region, a total of 1375 flies were caught over a period of 6 weeks from August 17th to September 27th. Of these, 1094 were dissected for gut, salivary glands and proboscis, i.e. wherever the gut was found infected, salivary glands and proboscis were

TABLE 1 Ifakara area

Infections classified according to localisation in male G. morsitans

The category "gut and salivary glands" has been left out, as no sure brucei-infections were found in the Ifakara area

Total of infections	* * *	1.6% (1)	4.1% (7)	10.2% (25)	15.9% (22)	19.5% (15)	21.5% (16)	11.2% (86)
-type) No. positive	proboscis	1.6% (1)	$3 \ \% \ (5)$	6.9% (16)	9.2% (11)	11.7% (7)	12.5% (7)	6.7% (47)
(Vivax) No. dissected	for proboscis	63	170	233	119	60	56	701
Gut and	proposeis		1.1% (2)	3.3% (9)	6.7% (11)	7.9% (8)	9 % (9)	4.5% (39)
Gut only **	1	2.4% (1)	2.1% (4)	6.3% (17)	2.4% (4)	$2 \ \% \ (2)$	4 % (4)	3.7% (32)
Remains of blood meal	*	2.4% (1)	2.6% (5)	1.1% (3)	1.2% (2)	1	4 % (4)	1.7% (15)
No. dissected	0+7+D	41	189	270	165	102	100	867
Mean age		9 d.	15 d.	25 d.	33 d.	34 d.	48 d. or older	
Age category		1	2	3	4	5	9	Total

be considered as proper "gut infections" but are mentioned to show, that at the time the survey was made, trypanosome carrying * In some cases, a few bloodforms were found in the partly digested remains of the last blood meal in the midgut. These cannot animals must have been quite frequent.

** Here at least some typical midgut forms were observed; their character remains undeterminate: they may develop later on into mature congolense or brucei-type infections.

*** Without gut infections.

TABLE 2 Kilosa kwa Mpepo area

Infections classified according to localisation in male G. morsitans

Total infected ***	1	2.5% (5)	7.5% (24)	13.2% (16)	13.1% (14)	38 % (22)		9.5% (81)
Proboscis only (vivax-type)	Ι	1.5% (3)	5 % (16)	7.4% (9)	8.4% (9)	17.3% (10)		5.5% (47)
Gut and salivary glands (brucei-type)]			0.95% (1)]		0.12% (1)
Gut and proboscis	1	1 % (2)	2.5% (8)	5.8% (7)	3.7% (4)	20.7% (12)		3.9%~(33)
Gut only **	1.8% (1)	$2 \ \% \ (4)$	4.1% (13)	4.1% (5)	3.8% (4)	5.2% (3)		3.5%~(30)
Remains of blood meal	l	3.1% (6)	1.9% (6)	2.4% (3)	1.9% (2)			2 % (17)
No. dissected G+P+S	55	196	317	122	106	58		854
Mean age	9 d.	15 d.	25 d.	33 d.	44 d.	48 d.	or older	
Age category	1	2	3	4	5	6		Total

* In some cases, a few bloodforms were found in the partly digested remains of the last blood meal in the midgut. These cannot be considered as proper "gut infections" but are mentioned to show, that at the time the survey was made, trypanosome carrying animals must have been quite frequent.

** Here at least some typical midgut forms were observed; their character remains undeterminate: they may develop later on into mature congolense or brucei-type infections.

*** Without gut infections.

examined too. 853 of the total number were examined for vivax-type infections (proboscis). The overall infection rate was 10.8% (without gut infections).

In the Kilosa kwa Mpepo region 994 flies were caught and dissected completely over a period of about 4 weeks from September until October 19th, showing an overall infection rate of 8.9%. As found by others authors, the greatest number were *vivax*-infections, followed by *congolense* and only one sure case of *brucei* (Kilosa kwa Mpepo area). As the percentage of trypanosome infections increases with the age of the flies (HARLEY, 1966), all results are classified according to age groups 1 to 6 (JACKSON, 1947). Graph 1 shows the relative distribution of the males and females caught in the two areas (on Landrover). The number of females proved to be very low in both areas, 16.5% around Ifakara and 14% only in the Kilosa kwa Mpepo region. Classified results of the different types of infection are therefore only given for male tsetse flies (table 1 Ifakara, table 2 Kilosa kwa Mpepo, combined results graph 2).



Discussion

There appears no significant difference in the incidence of trypanosomes in the transmitting tsetse flies examined in two different areas of the Ulanga District, one area, where only animal trypanosomiasis is known and another, which, in addition, is an endemic sleeping sickness area. In the latter, only one out of 994 tsetse flies harboured brucei-group trypanosomes, although along the track used for fly-catching Phacochoerus aethiopicus, Kobus ellipsiprymnus, Redunca arundinum, Hippotragus niger and Alcelaphus lichtensteini could be observed, all proved to be reservoirs for brucei-group trypanosomes. Some of the above mentioned animals show even a very high incidence (GEIGY et al., 1967). On the other hand, only mature *brucei*-infections can be definitely determined and these are to be found usually in age categories 4, 5 and 6, which represent a small number of the flies in our catches, namely 14%, 11.6% and 6.6% respectively. The evaluation of the tsetsefly infections is complicated even more by the fact, that most of the reservoir animals examined last year showed mixed infections. The tract we used in the Kilosa kwa Mpepo region is also frequently used by man, going hunting, collecting firewood or wild honey. Getting information about the localities, where people are bitten by infected tsetse flies proves to be very difficult. The population is thinly scattered over the area and usually only the dispensaries at which the diagnosis was made is known. A serological survey of the whole population of the entire area may help to clarify the situation.

Acknowledgements

We are indebted again to the Central Veterinary Laboratory (Dar es Salaam) for supplying us with the necessary white rats and mice as well as to the Swiss Capuchin Mission for their kind hospitality.

References

- GEIGY, R., KAUFFMANN, M. & BEGLINGER, R. (1967). A survey of wild animals as potential reservoirs of trypanosomiasis in the Ulanga District (Tanzania).
 — Acta trop. 24, 97-108
- HARLEY, J. M. B. (1966). Seasonal and diurnal variations in physiological age and trypanosome infection rate of females of *Glossina pallidipes* Aust., *G. palpalis fuscipes* Newst. and *G. brevipalpis* Newst. — Bull. ent. Res. 56, 595-614
- JACKSON, C. H. N. (1947). An artificially isolated generation of tsetse flies (Diptera). Bull. ent. Res. 37, 291-299
- JORDAN, A. M. (1964). Trypanosome infection rates in *Glossina morsitans sub*morsitans Newst. in Northern Nigeria. — Bull. ent. Res. 55, 219-231
- SAUNDERS, D. S. (1962). Age determination for female tsetse flies and the age compositions of samples of *Glossina pallidipes* Aust., *G. palpalis fuscipes* Newst. and *G. brevipalpis* Newst. — Bull. ent. Res. 53, 579-595