

**Zeitschrift:** Archives des sciences [1948-1980]  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 13 (1960)  
**Heft:** 9: Colloque Ampère

**Artikel:** Nuclear magnetic relaxation in vanadium  
**Autor:** Butterworth, J.  
**DOI:** <https://doi.org/10.5169/seals-738612>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Nuclear Magnetic Relaxation in Vanadium

by J. BUTTERWORTH

Metallurgy Division, A.E.R.E., Harwell, England

---

## *Summary.*

Measurements of the nuclear magnetic spin-spin and spin-lattice relaxation in the transition metal vanadium have been made at temperatures of 20.4° K, 77.3° K and 292° K. No evidence of the presence of localized unpaired electrons in vanadium was found. By comparison of the results with measured values of other physical properties of vanadium, it is deduced that there is such thorough admixture of *s*-states and *d*-states at the Fermi surface in vanadium that the spin susceptibility of the *s*-electrons is enhanced, by exchange interaction, to the same high degree as that of the *d*-electrons, namely, about 2.5 times. It is also deduced that the proportion of all the states at Fermi surface having *s*-character is roughly 0.17.

From a comparison of the measured values of the Knight shift,  $\Delta H/H$ , and the spin-lattice relaxation time,  $T_1$ , of the nuclear magnetic resonance in a metal, information can be derived about the state of the electrons in the metal. For, as Korringa showed [1], if the electrons responsible for the shift and the relaxation behave as uncorrelated electrons of largely *s*-character, the following simple relation exists between  $T_1$  and  $\Delta H/H$

$$T_1 \left( \frac{\Delta H}{H} \right)^2 = \frac{\hbar}{\pi k T} \left( \frac{\beta I}{\mu^2} \right)^2 \quad (1)$$

where  $T$  is the absolute temperature, and  $\beta$  and  $\mu$  are the magnetic moments of an electron and nucleus, respectively, while the other symbols have their conventional meanings. Conversely, departure from the Korringa relation indicates departure of the electrons from such simple behaviour.

Information about the electrons can also be obtained from measurements of the spin-spin relaxation, particularly if there is some question of the existence of localized states containing electrons of unpaired spin.

This note reports the results of measurements of  $T_1$ , and of the spin-spin relaxation of the  $^{51}\text{V}$  resonance in vanadium metal, and some inferences drawn from them. The results are of some interest because they are the first reported accurate measurements of nuclear magnetic

relaxation in a transition metal, where the unfilled shell of  $d$ -electrons might be expected to influence strongly the nuclear magnetic resonance. As a probable consequence of these electrons the results show much the largest departure from the Korringa relation observed so far.

The sample consisted of 2.4 gram of vanadium powder obtained by grinding the metal on an alumina wheel and subsequently sieving it through a 200 mesh. The metal was initially prepared electrolytically by the U.S. Bureau of Mines Electrometallurgical Section, and was found by analysis to have a purity of 99.7%, the principal impurity being 0.09% by weight of dissolved iron.




FIG. 1. INDUCTION DECAY, AFTER  $\frac{\pi}{2}$  PULSE, FROM TRANSVERSE NUCLEAR MAGNETIC MOMENT OF  $^{51}\text{V}$  NUCLEI IN VANADIUM METAL.

Fig. 1.

The measurements of relaxation were made at temperatures of 20.4° K, 77.3° K and 292° K, and at a frequency of 11.000 Mc/s, by a pulsed radio-frequency method incorporating coherent detection and post-detector integration. The pulse sequence used in the measurement of  $T_1$  consisted of an initial  $\pi/2$  pulse followed, after a variable period, by a further  $\pi/2$  pulse, while measurement of the induction decay following a single  $\pi/2$  pulse gave information about spin-spin relaxation.

The induction decay observed at all temperatures is shown in figure 1. We see from the overshoot that the decay is not that from a Gaussian line shape [2], for such a line would result in a decay having a time,  $t$ , dependence of the form  $\exp(-\gamma^2\sigma^2 t^2/2)$ , where  $\gamma$  is the magnetogyric ratio of the nucleus, and  $\sigma^2$  is the mean square deviation of the field at

the nuclei. The overshoot is a common feature of induction decays from rigid lattices [3]. The departure from Gaussian behaviour, however, does not begin until after some  $60\mu\text{s}$ , as is shown by figure 2, where the induction decay is shown as a function of  $t^2$ . The value of  $\sigma^2$  derived from the straight part of the curve is  $31 \text{ gauss}^2$ , which value may be compared with the calculated dipolar value [5] of  $19.3 \text{ gauss}^2$ , and that, derived from conventional measurements [5], of  $32 \text{ gauss}^2$ .




FIG. 2. AMPLITUDE OF INDUCTION DECAY, FROM  $^{51}\text{V}$  NUCLEI IN VANADIUM METAL, AGAINST  $(\text{TIME})^2$

Fig. 2.

In the reasonable agreement of the calculated dipolar line width with that derived from the measurements, and the temperature independence of the latter, we see no evidence for the existence in vanadium of localized unpaired electronic spins. The extra broadening is attributed to indirect spin-spin coupling of the nuclei via the conduction electrons.

The values of  $T_1$  obtained are given in Table 1, together with the product of  $T_1$  and temperature  $T$ .

We see that, within experimental error, the value of  $T_1$  varies inversely as the absolute temperature from  $292^\circ \text{K}$  down to  $20.4^\circ \text{K}$ , indicating that, over this temperature range, the dominant mechanism of spin-

lattice relaxation is the interaction between the nuclear spins and the spins of the conduction electrons. Thus, again, there is no sign of any effects due to spins of localised *d*-electrons.

TABLE 1.  
*Values of  $T_1$  and of  $T_1T$  against temperature.*

T (°K)	$T_1$ (m sec)	$T_1T$ (sec. °K)
20.4	$38.4 \pm 1\%$	$0.783 \pm 1\%$
77.3	$10.2 \pm 1\%$	$0.788 \pm 1\%$
292	$2.70 \pm 1\%$	$0.788 \pm 1\%$

If we assume that the electrons producing the spin-lattice relaxation are behaving as uncorrelated electrons of mainly *s*-character and use the measured value of  $T_1T$  in equation (1), we can calculate the value of Knight shift such electrons would produce. By this means we derive a value for  $\Delta H/H$  of 0.219%. Measured values, however, are some 2.5 times this, ranging from 0.55% [4] to 0.58% [5].

Such a discrepancy is in the wrong direction to be explained by invoking electron states of other than *s*-character, for such states would merely contribute to the relaxation without affecting the shift. Furthermore, because of their low spin density at the nucleus, their effect on relaxation would be small.

Consequently, we assume that  $T_1$  is largely determined by the effective density of *s*-states at the Fermi surface. As this density appears, from the temperature-independence of  $T_1T$ , to be constant, and because of the lack of a plausible mechanism for prolonging  $T_1$ , we conclude that  $\Delta H/H$  and, hence, the spin paramagnetism of the *s*-states is enhanced by a factor of about 2.5.

However, further information about the relation of the effective density of states of the conduction electrons to their spin paramagnetism,  $\chi_p$ , can be derived from measurements of the coefficient of electronic specific heat,  $\gamma_e$ , and the magnetic susceptibility of the metal. For, if the conduction electrons behave as uncorrelated the following simple relation exists between  $\chi_p$  and  $\gamma_e$  [6].

$$\frac{\chi_p}{\gamma_e} = \frac{3 \beta^2}{\pi^2 k^2} \quad (2)$$

This relation is analogous to the Korringa one, but refers to all the electronic states at the Fermi surface which are expected in vanadium to be of largely *d*-character, and not merely to those of *s*-character.

The measured value of  $\gamma_e$  in vanadium [7] is  $8.8 \times 10^{-3}$  Joule/gram atom. deg<sup>2</sup>. If we again assume that the electrons are uncorrelated and insert the value of  $\gamma_e$  in equation (2) we obtain a value for the spin susceptibility of these electrons of  $122 \times 10^{-6}$  emu/gram atom. Childs et al [8] have measured the total susceptibility of vanadium and, after making small corrections for diamagnetism, have arrived at a value for the spin susceptibility of  $308 \times 10^{-6}$  emu/gram atom. Thus, the spin susceptibility of the mainly *d*-type electrons at the Fermi surface is also some 2.5 times greater than that expected from their effective density of states.

In the present state of theory the explanation of this surprising coincidence must be to some extent speculation, but in the absence of any better theory, it is suggested that the enhancement of the susceptibility is a result of exchange interactions between the conduction electrons, while one interpretation of the fact that the susceptibility of the *s*-electrons is enhanced to the same high degree as that of the *d*-electrons is that there is such thorough admixture of *s*-states and *d*-states at the Fermi surface, that they are equally affected by exchange effects.

From the measured values of  $T_1$  and  $\gamma_e$ , we can deduce approximately the proportion,  $\xi$ , of the electrons at the Fermi surface having *s*-character, by means of Korringa's formula

$$T_1^{-1} = 2 \pi k T \hbar^{-1} \xi^2 n^2(E_0) a^2(s) \quad (3)$$

where  $n(E_0)$  is the total density of states at the Fermi surface, obtained from  $\gamma_e$ , and  $a(s)$  is the hyperfine coupling constant for an *s*-electron. Using the value of  $a(s)$  calculated by Knight [9] in equation (3) leads to a value for  $\xi$  of 0.17. The fact that such a reasonable value for  $\xi$  can be obtained from  $T_1$  alone tends to substantiate the view that the departure from the Korringa relation is mainly due to enhancement of the Knight shift.

Thanks are due to Dr. L. E. Drain, Dr. W. E. Gardner, Mr. A. D. Le Claire and Dr. W. Marshall for helpful discussions and to Mr. B. Lent for experimental assistance.

## REFERENCES

1. KORRINGA, J., *Physica*, **16**, 601 (1950).
2. ANDERSON, P.W., *J. Phys. Soc. Japan*, **9**, 316 (1954).
3. LOWE, I. J. and R. E. NORBERG, *Phys. Rev.*, **107**, 46 (1957).
4. KNIGHT, W. D., *Solid State Physics*, **2**, 122 (1956).
5. DRAIN, L. E., private communication.
6. MOTT and JONES, *Metals and Alloys*, 179-185. Oxford University Press, England (1936).
7. CORAK, W. S., B. B. GOODMAN, C. B. SATTERTHWAITHE and A. WEXLER, *Phys. Rev.*, **96**, 1442 (1954).
8. CHILDS, GARDNER and PENFOLD, *Phil. Mag.*, to be published.
9. KNIGHT, W. D., *ibid.*, 120.

## RELAXATION MAGNÉTIQUE NUCLÉAIRE DANS LE VANADIUM

*Résumé.*

Des mesures de relaxation magnétique nucléaire spin-spin et spin-réseau ont été effectuées dans le métal de transition Vanadium à des températures de 20,4° K — 77,3° K et 292° K. Aucune trace de présence d'électrons non appariés n'est apparue dans le Vanadium. La comparaison de ces résultats avec les valeurs mesurées à partir d'autres propriétés physiques du Vanadium permet de déduire qu'il y a un tel mélange d'états *s* et d'états *d* à la surface de Fermi du Vanadium, que la susceptibilité magnétique des électrons *s* est augmentée, par interaction d'échange, au même degré que celle des électrons *d*, à savoir environ 2,5 fois. On a également déduit que la proportion de tous les états à la surface de Fermi ayant un caractère *s* est environ 0,17.

---