

Zeitschrift: Archives des sciences [1948-1980]
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 13 (1960)
Heft: 9: Colloque Ampère

Artikel: High resolution paramagnetic proton resonance spectra of some metal-ethyl compounds
Autor: Klose, G.
DOI: <https://doi.org/10.5169/seals-738632>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 05.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

High resolution paramagnetic proton resonance spectra of some metal-ethyl compounds

by G. KLOSE

E. B. Baker [1] found, that the indirect spin-spin coupling between the nucleus of the lead isotope ^{207}Pb of leadtetraethyl and the protons of the CH_3 group is bigger than the coupling of the ^{207}Pb isotope with the protons of the CH_2 group attached direct to the metal atom. P. T. Narasimhan and M. T. Rogers obtained the same result for mercury diethyl [2].

We made a study of the spectra of tin tetraethyl and diethyl selenide in order to obtain the different coupling constants and the internal chemical shifts between the protons of the CH_3 and CH_2 groups.

The spectrum of tin tetraethyl of a frequency of 25 Mc/s consists of three intense central lines with several weak satellite lines on either side. These satellite lines are the result of the unequal spin coupling of the ^{117}Sn (spin 1/2, abundance 7,67%) and ^{119}Sn isotopes (spin 1/2, abundance 8,68%) with the CH_3 and CH_2 groups. The observed spectrum is in good agreement with the calculated spectrum (type: $B_3 A_2$, $B_3 A_2 X$, $B_3 A_2 Y$) with the following data:

$$\begin{aligned} J \text{ H}(\text{CH}_3) - \text{H}(\text{CH}_2) &= 7,9 \text{ c/s}; \delta = \nu_{\text{H}(\text{CH}_3)} - \nu_{\text{H}(\text{CH}_2)} = -9,3 \text{ c/s}; \\ J_3 \text{ }^{119}\text{Sn} - \text{H}(\text{CH}_3) &= 69,6 \text{ c/s}; J_3 \text{ }^{117}\text{Sn} - \text{H}(\text{CH}_3) = 66,5 \text{ c/s}; \\ J_2 \text{ }^{119}\text{Sn} - \text{H}(\text{CH}_2) &= 51,9 \text{ c/s}; J_2 \text{ }^{117}\text{Sn} - \text{H}(\text{CH}_2) = 49,6 \text{ c/s}. \end{aligned}$$

For diethyl selenide we obtained $J \text{ H}(\text{CH}_3) - \text{H}(\text{CH}_2) = 7,0 \text{ c/s}$ and $\delta = \nu_{\text{H}(\text{CH}_3)} - \nu_{\text{H}(\text{CH}_2)} = 36,2 \text{ c/s}$. Lines which are the result of a coupling between the ^{77}Se isotope (spin 1/2, abundance 7,5%) and the CH_3 and CH_2 groups were not observed.

The Pauling electronegativities for tin and selenium calculated from the internal chemical shifts confirm the suggestion of Shoolery [2], that the value of the constant in the original Dailey-Shoolery equation [3] must be changed from 1,71 to 2,10 for low electronegativities. A more detailed paper is in preparation.

1. BAKER, E. B., *J. Chem. Phys.*, 26, 960, 1957.
 2. NARASIMHAN, P. T., M. T. ROGERS, *J. Am. Chem. Soc.*, 82, 34, 1960.
 3. DAILEY, B. P., J. N. SHOOLERY, *J. Am. Chem. Soc.*, 77, 3977, 1955.
-