

Zeitschrift: Archives des sciences [1948-1980]
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 14 (1961)
Heft: 10: Colloque Ampère

Artikel: Entropy of spin systems in magnet resonance phenomena
Autor: Vojta, Günter
DOI: <https://doi.org/10.5169/seals-739628>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Entropy of Spin Systems in Magnetic Resonance Phenomena

by Günter VOJTA

Theoretisch-Physikalisches Institut, Universität, Leipzig,
und Arbeitsstelle für Statistische Physik, Leipzig,
der Deutschen Akademie der Wissenschaften zu Berlin

A well-known principal difficulty in statistical thermodynamics of irreversible processes is the correct introduction of the concepts of entropy and temperature [1]. It was shown by the author [2], that for relaxing spin systems under the influence of alternating magnetic fields the "entropy" defined by Wangsness [3, 4] is physically meaningless and that therefore a principle of minimum entropy production for simple or double resonance phenomena (e.g. Overhauser effect) cannot be established without further considerations.

Often the description of a spin state by means of occupation numbers N_i of energy niveaux is sufficient (one neglects then the nondiagonal elements of the density matrix ρ , these are proportional to the amplitude of the H_1 -field). In this case the entropy for a system of N (noninteracting) spins (in Boltzmann statistics) is simply

$$S = -Nk \sum (N_i/N) \ln (N_i/N), \quad (1)$$

and the minimum entropy production principle is to be formulated by use of a master equation for $\partial N_i/\partial t$ [5] or in terms of spin temperatures [6], the spin temperature being only an occupation number parameter and no longer a thermodynamic temperature.

For spin systems with irreversible processes there is no need to change the v. Neumann equilibrium definition of entropy

$$S = -Nk Tr(\rho \ln \rho) \quad (2)$$

(compare also [1, 7]). Callen [8] stated in the language of phenomenological thermodynamics of irreversible processes, that in absence of magnetic fields the microscopic density matrix ρ itself generally minimizes the rate

of entropy production. The author shows [9] for a spin system in a stationary state, that by use of a suitably orientated coordinate system, rotating with the frequency ω of the H_1 -field, the following statements are valid:

- 1) the transformed density matrix $\tilde{\rho}$ is diagonal;
- 2) the correct calculation of the entropy according to (2) amounts to the use of (1). The entropy is the same as in the original coordinate system;
- 3) a spin temperature (as an occupation number parameter) can be defined;
- 4) the principle of minimum entropy production is valid formally.

1. The diagonalization of ρ is performed with a unitary matrix U :

$$\tilde{\rho} = U^{-1} \rho U \quad (3)$$

with (in common notation)

$$U = e^{i\varphi I_y} e^{-i(\omega t - \psi) I_z} \quad (4)$$

(I_y, I_z : Pauli spin operators). With the stationary matrix [2, 10]

$\left(\text{for } I = \frac{1}{2} \right)$

$$\rho = \frac{1}{Z} \begin{pmatrix} 1 + \frac{1}{2}a_0 + \frac{1}{8}(a_0^2 + a_1 a_1^*) & a_1 e^{i\omega t} \\ a_1^* e^{-i\omega t} & 1 - \frac{1}{2}a_0 + \frac{1}{8}(a_0^2 + a_1 a_1^*) \end{pmatrix} \quad (5)$$

$$\text{with } Z = 2 \left\{ 1 + \frac{1}{8}(a_0^2 + a_1 a_1^*) \right\} \approx 2 \text{ (partition function)},$$

$$a_0 = \frac{\gamma \hbar}{kT} \left(H_0 - \frac{\omega \gamma H_1^2 \tau^2}{n} \right) \text{ (T : lattice temperature)},$$

$$a_1 = \frac{\gamma \hbar H_1}{kT} \left(1 - i\omega \tau \frac{1 + i(\gamma H_0 - \omega) \tau}{n} \right),$$

$$n = 1 + (\gamma H_0 - \omega)^2 \tau^2 + \gamma^2 H_1^2 \tau^2,$$

$$a_1 a_1^* = \left(\frac{\gamma \hbar H_1}{kT} \right)^2 \frac{\chi'^2 + \chi''^2}{\chi_0},$$

it follows

$$\tilde{\rho} = \frac{1}{2} \begin{pmatrix} 1 + \frac{1}{Z} \sqrt{a_0^2 + a_1 a_1^*} & \sigma \\ \sigma & 1 - \frac{1}{Z} \sqrt{a_0^2 + a_1 a_1^*} \end{pmatrix}. \quad (6)$$

From (3) ... (6) the angle φ between the new and the old z -axis can be calculated, and likewise the angle $\psi = \text{arc tg } (\chi''/\chi') = \text{arc tg } (v/u)$ (v : absorption, u : dispersion). The new coordinate system is inclined to the laboratory system and rotates with ω and a phase difference ψ behind the H_1 -field.

2. The entropy is, using (5) or (6),

$$\begin{aligned} S &= -NkTr(\rho ln\rho) = -NkTr(\tilde{\rho} ln\tilde{\rho}) \\ &= -Nk \left\{ \ln 2 - \frac{1}{2Z^2} (a_0^2 + a_1 a_1^*) \right\}. \end{aligned} \quad (7)$$

3. By writing (6) approximatively in exponential form and setting $Z \approx 2$, the spin temperature T_s is defined by

$$\frac{1}{Z} e^{-\tilde{H}/kT} = \frac{1}{Z} e^{-\gamma\hbar\sqrt{H_0^2 + H_1^2}/kT_s} \quad (8)$$

(\tilde{H} : formal Hamiltonian in the transformed coordinate system), or with $\beta_s = 1/kT_s$ by

$$\beta_s = \frac{1}{\sqrt{H_0^2 + H_1^2}} \sqrt{(H_0 - \omega\gamma H_1^2 \tau^2/n)^2 + H_1^2 (\chi'^2 + \chi''^2)/\chi_0^2}. \quad (9)$$

4. For the discussion of the entropy production see [9].

For spin systems obeying *FD*-statistics, e.g. electron spins in metals, the separation of the system entropy in lattice entropy and spin entropy in density matrix or occupation number formalism is not possible [11]. Our transformation to a rotating coordinate system is purely formal and has only the purpose of diagonalizing the density matrix, in contrary to the work of Redfield, Slichter, and others [12, 13], where an *effective* Hamiltonian is introduced and *physics in the rotating coordinate system* is studied.

1. See for example BLATT, J. M., *Progr. Theor. Phys.*, **24**, 741 (1960).
 2. VOJTA, G., *Ann. Physik*, **6**, 31 (1960). Compare LURCAT, F., *C. R. Acad. Sci. (Paris)*, **242**, 1686 (1956).
 3. WANGNESS, R. K., *Phys. Rev.*, **104**, 857 (1956).
 4. See also MEIJER, P. H. E., *Proc. Int. Symp. Transport Processes in Statistical Mechanics*, Brussels 1958, 319, and personal communication to the author.
 5. KLEIN, M. J., *Phys. Rev.*, **98**, 1736 (1955).
 6. VOJTA, G., Tagungsbericht Hochfrequenzspektroskopie, Leipzig 1960, p. 217-227.
 7. LANDSBERG, P. T., *Proc. Phys. Soc. (London)*, **74**, 486 (1959).
 8. CALLEN, H. B., *Phys. Rev.*, **105**, 360 (1957).
 9. VOJTA, G., to be published in *Ann. Physik*.
 10. —— *Z. Naturforschg.*, **12a**, 282 (1957).
 11. FICK, E., *Z. Physik*, **157**, 407 (1960); VOJTA, G., unpublished.
 12. REDFIELD, A. G., *Phys. Rev.*, **98**, 1787 (1955); **101**, 67 (1956). SLICHTER, C. P. and W. C. HOLTON, *Phys. Rev.*, **122**, 1701 (1961).
 13. SLICHTER, C. P., *Phys. Rev.*, **99**, 1822 (1955).
-

DISCUSSION

M. Kastler. — Lässt sich die Entropie-Definition auch durchführen bei einer Sachlage, die einer negativen absoluten Temperatur des Spinsystems (Besetzungs-umkehr) entspricht?

M. Vojta. — Ja in genau der gleichen Weise wie für eine positive absolute Temperatur des Spinsystems.