

Zeitschrift: Archives des sciences [1948-1980]
Herausgeber: Société de Physique et d'Histoire Naturelle de Genève
Band: 14 (1961)
Heft: 10: Colloque Ampère

Artikel: Proton spin-lattice relaxation in acetic acid
Autor: Hrynkiewicz, A.Z. / Krynicki, K. / Waluga, T.
DOI: <https://doi.org/10.5169/seals-739639>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Proton spin-lattice relaxation in acetic acid

by A. Z. HRYNKIEWICZ, K. KRYNICKI and T. WALUGA

Institute of Nuclear Physics, Polish Academy of Sciences,
Cracow 23, Poland

Résumé.

L'évidence expérimentale de l'accroissement non-exponentiel de la magnétisation des protons dans l'acide acétique liquide à diverses températures est présentée. Ce fait est expliqué par l'existence de deux différents temps de relaxation spin-réseau dans les groupes CH_3 et COOH.


The purpose of this communication is to provide some experimental evidence showing that in liquid acetic acid (CH_3COOH) the protons of the CH_3 group have a different spin-lattice relaxation time, T_1 , from the COOH proton.

Similar results have been obtained for methyl alcohol by Powles and Cutler [1] and for liquid toluene by different authors [2, 3, 4]. The simultaneous existence of two T_1 values leads to the non-exponential growth of nuclear magnetization in time.

T_1 measurements for acetic acid have also been reported by Bloembergen et al. [5] and Chiarotti et al. [6], who obtained the single relaxation times of 2.4 sec and 3.8 sec respectively. But the short observation time of the relaxation process involved in the method used made these authors unable to observe the non-exponential behaviour of magnetization. In addition, the sample used by Bloembergen et al. was not oxygen-free.

In the research here reported the sample of CH_3COOH was saturated and then the growth of nuclear magnetization M was observed, using the method of Hennel and Hrynkiewicz [7]. The growth time was varied within sufficiently wide limits to obtain the values of $-\ln(1 - M/M_0)$ between 0 and 3.8. The sample was prepared using C.P. glacial acetic acid. To purify, it was distilled four times. The atmospheric oxygen was removed by boiling at length as described by Hennel et al. [8].

The measurements were carried out at a resonant frequency of 28 Mc/sec at four temperatures, 20, 30, 50 and 80° C. The experimental results obtained for 20° C are shown in figure 1.



Growth of the magnetization M in time in liquid acetic acid at 20° C. The theoretical curve is drawn according to equation (1): $a = 3/4$, $b = 1/4$, $T'_1 = 3.12$ sec, $T''_1 = 9.55$ sec.

A theoretical curve of the following form has also been drawn:

$$1 - M/M_0 = a \exp\left(-\frac{t}{T'_1}\right) + b \exp\left(-\frac{t}{T''_1}\right), \quad (1)$$

where $a = 3/4$ and $b = 1/4$ according to the abundances of protons in the methyl and carboxyl groups. The relaxation times of these groups, T'_1 and T''_1 , (3.12 sec. and 9.55 sec.) have been chosen to fit the curve to the experimental points. The values of T'_1 and T''_1 shown in table 1 were obtained in the same way. The maximum random error was roughly estimated as 10% for T'_1 and 25% for T''_1 .

A non-exponential decay of the transversal magnetization for acetic acid is reported by Daszkiewicz [9].

The fact that two different relaxation times, T_1' and T_1'' , are present indicates that there is no rapid proton chemical exchange between the carboxyl and methyl groups. This result may be anticipated in view of the strong hydrogen bond (8.2 kcal/mole) in which the COOH proton is involved causing the well-known dimeric structure of acetic acid [10, 11].

TABLE 1.

Experimental values of proton T_1 for the methyl (T_1') and carboxyl (T_1'') groups in acetic acid.

°C	T_1' (sec)	T_1'' (sec)	T_1'/T_1''
20	3.12	9.55	3.06
30	3.58	12.3	3.44
50	4.76	15.1	3.17
80	6.92	19.2	2.77

The translational part of $1/T_1$ was estimated taking the formula of Blicharski et al. [12] as a basis. The viscosity of 1.211 cp at 20°C was taken. The obtained result was $(1/T_1)_{tr} = 0.106 \text{ sec}^{-1}$, i.e. $(T_1)_{tr} = 9.43 \text{ sec}$. The agreement with the experimental value $T_1'' = 9.55 \text{ sec}$ means that the spin-lattice relaxation time of the carboxyl group is mainly controlled by the intermolecular interactions.

As shown in table 1, the T_1'/T_1'' ratio is independent of the temperature within the limits of experimental error.

The authors wish to thank Dr. J. W. Hennel for helpful discussions.

REFERENCES

- POWLES, J. C. and D. CUTLER, *Arch. d. Sc.*, 12 (8th Colloque Ampère), 135 (1959).
- NEDERBRAGT, G. W. and C. A. REILLY, *J. Chem. Phys.*, 24, 1110 (1956).
- POWLES, J. G. and D. J. NEALE, *Proc. Phys. Soc. (London)*, 77, 737 (1961).
- BONERA, G., L. CHIODI and A. RIGAMONTI, *Nuovo Cimento*, 17, 198 (1960).
- BLOEMBERGEN, N., E. M. PURCELL and R. V. POUND, *Phys. Rev.*, 73, 679 (1948).
- CHIAROTTI, G., G. CRISTIANI and L. GIULOTTO, *Nuovo Cimento*, 1, 863 (1955).
- HENNEL, J. W. and A. Z. HRYNKIEWICZ, *Arch. d. Sc.*, 11 (7th Colloque Ampère), 238 (1958).

8. HENNEL, J. W., K. KRYNICKI, T. WALUGA and G. ZAPALSKI, *Acta Phys. Polon.*, 20, 77 (1961).
 9. DASZKIEWICZ, O., this Conference.
 10. HADZI, D. and N. SHEPPARD, *Proc. Roy. Soc. (London)*, 216 A, 247 (1953).
 11. BHAR, B. N., W. FORSLING and G. LINDSTRÖM, *Arkiv för Fysik*, 10, 59 (1956).
 12. BLICHARSKI, J., J. W. HENNEL, K. KRYNICKI, J. MIKULSKI, T. WALUGA and G. ZAPALSKI, *Bulletin Ampère*, 9 (9th Colloque Ampère), 452 (1960), formula (11).
-