Zeitschrift:	Archives des sciences [1948-1980]
Herausgeber:	Société de Physique et d'Histoire Naturelle de Genève
Band:	14 (1961)
Heft:	10: Colloque Ampère
Artikel:	Measurements of the temperature dependence of the proton spin-lattice relaxation time in liquid hydrogen sulphide
Autor:	Hennel, J.W. / Waluga, T.
DOI:	https://doi.org/10.5169/seals-739643

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 03.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Measurements of the Temperature Dependence of the Proton Spin-Lattice Relaxation Time in Liquid Hydrogen Sulphide

by J. W. HENNEL and T. WALUGA

Institute of Nuclear Physics, Polish Academy of Sciences, Cracow 23, Poland

Résumé.

Le temps de relaxation spin reseau T_1 , des protons dans le H_2S liquide a été mesuré dans le domaine de la température de — 18 à $+55^{\circ}$ C. Les résultats obtenus montrent que T_1 décroît avec l'accroissement de la température. La possibilité d'expliquer ce fait par l'action du champ magnétique provenant de la rotation a été discutée.

Recently Gutowsky et al. [3] presented some results of the spin-lattice relaxation time T_1 measurements showing that the fluorine T_1 in liquid CHF₂Cl and CHF₃ decreases with temperature in the range from about -100° C to at least room temperature. The authors explain this effect as due to spin-rotational interaction. The resulting contribution to the $1/T_1$ was described as proportional to the square of nuclear magnetic shielding constant and as dependent on "quenching" by intermolecular interactions which becomes less important at higher temperature.


The purpose of this report is to present an example of decrease of T_1 in the case of protons. As shown on figure 1 the proton T_1 in liquid hydrogen sulphide decreases by about 30 per cent in a temperature range from -10 to $+55^{\circ}$ C.

The sample used was prepared from very pure oxygen-free hydrogen and very pure, degassed sulphur in a similar way as that described by Klemenc and Bankowski [7]. The obtained hydrogen sulphide was twice fractionally distilled in vacuum at $--60^{\circ}$ C and carefully degassed. The measurements were made by the method of Hennel and Hrynkiewicz [4] at 28 Mc/sec. under the pressure of the saturated vapour.

It should also be noticed that there are numerous liquids in which the increase of T_1 with temperature is much smaller than that predicted by

COLLOQUE DE LEIPZIG 1961

theory. E.g. T_1 in methyl chloride remains constant within a range from $+10^{\circ}$ C to at least $+70^{\circ}$ C [1]. The decrease with temperature of the function $T_1\eta/T$ (η is the viscosity, T is the absolute temperature) in the following liquids, CHCl₃, CHBr₃, H₂O, H₂SO₄, CH₂Cl₂, NH₃, CH₃Cl, C₂H₅OH shown in our earlier papers [1, 5] and by Guillotto et al. [2] can



Results of measurements of the proton T_1 in liquid hydrogen sulphide.

also be regarded as an "insufficient" increase of T_1 . (The function $T_1\eta/T$ should be independent of the temperature if spin-rotational interaction is not taken into account.) Also in the case of H_2S the decrease of viscosity with temperature is known [6]. The objection may be raised that the relation between T_1 and viscosity is only a rough approximation; nevertheless the fact that all investigated liquids showed a decrease of $T_1\eta/T$ is remarkable. It is also interesting that the smallest decrease of this function was found in water in the temperature range from 0° C to 100° C where the strong hydrogen bonding is known.

It is possible that the decrease of T_1 in H_2S is also caused by spinrotational interaction. Although the magnetic shielding constant is small for protons [3] on the other hand the intermolecular interactions in H_2S , in the temperature region concerned, are weak and therefore the "quenching" may be small. It is interesting to investigate the temperature dependence of T_1 in other liquids not much below the critical temperature. Work along these lines is in progress.

REFERENCES

- 1. BLICHARSKI, J., J. W. HENNEL, K. KRYNICKI, J. MIKULSKI, T. WALUGA and G. ZAPALSKI, Bulletin Ampère, 9 (9th Colloque Ampere), 452 (1960).
- 2. GIULOTTO, L., G. LANZI and L. TOSCA, Arch. d. Sc., 10 (6th Colloque Ampère), 251 (1957); Nuovo Cimento Supp. to 9, 398 (1958).
- 3. GUTOWSKY, H. S., I. J. LAWRENSON and K. SHIMOMURA, Phys. Rev. Letters, 6, 349 (1961).
- HENNEL, J. W. and A. Z. HRYNKIEWICZ, Arch. d. Sc., 11 (7th Colloque Ampere), 238 (1958).
- 5. HENNEL, J. W., A. Z. HRYNKIEWICZ, K. KRYNICKI, T. WALUGA and G. ZAPALSKI, Arch. d. Sc., 11 (7th Colloque Ampère), 243 (1958).
- 6. HENNEL, J. W. and K. KRYNICKI, Acta Physica Polonica, 19, 523 (1959).
- 7. KLEMENC, A. and O. BANKOWSKI, Z. anorg. Ch., 208, 348 (1932).