

**Zeitschrift:** Archives des sciences [1948-1980]  
**Herausgeber:** Société de Physique et d'Histoire Naturelle de Genève  
**Band:** 29 (1976)  
**Heft:** 2

**Artikel:** Reduction of 2-chloro-6-methoxy-4-methylquinoline. Part II  
**Autor:** Pandey, R. K. / Surana, Asha / Joshi, B. C.  
**DOI:** <https://doi.org/10.5169/seals-739680>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 18.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## REDUCTION OF 2-CHLORO-6-METHOXY- 4-METHYLQUINOLINE

### PART-II

BY

R. K. PANDEY, ASHA SURANA and B. C. JOSHI<sup>1</sup>

In continuation of our earlier communication<sup>1</sup> the present study deals with the reduction of 2-chloro-6-methoxy-4-methylquinoline with tin and hydrochloric acid when alongwith 6-methoxyquinoline (2) another compound identified as 2,2'-bis-6-methoxy-4-methylquinoline was also isolated (3).

2-Chloro-6-methoxy-4-methyl quinoline [1] was reduced with tin and HCl when 6-methoxy-4-methylquinoline [2] as a major component was obtained. Besides this, another compound [3] in minor quantity was also obtained and on the basis of elemental analysis an empirical formula  $C_{11}H_{10}ON$  was suggested to 3. No chlorine was found to be present.

Molecular ion peak in mass spectrum was found at m/e 344, thus suggesting the molecular formula to 3 as  $C_{22}H_{20}O_2N_2$ .

Nmr spectra (in  $CDCl_3$  using TMS as internal reference standard) revealed the presence of a singlet at  $\delta$  2.55 for six protons and another singlet at  $\delta$  3.90 also for six protons. Eight aromatic protons were found to be present in the range  $\delta$  7.05-7.95. The singlet at  $\delta$  2.55 would account for the presence of two methyl groups and a singlet at  $\delta$  3.9 would account for the presence of two methoxy groups. The absence of the chlorine in the molecule revealed the involvement of position-2 of the quinoline in the reaction indicating the compound 3 as a bis molecule.

The mass spectrum of 3 gave a base peak at m/e 172 besides the molecular ion peak indicating the cleavage of the molecule at positions 2,2' giving an ion 4.

From all these observations the structure to this compound (3) was assigned as 2,2'-bis-6-methoxy-4-methylquinoline.

---

<sup>1</sup> Chemical Laboratories, University of Rajasthan, Jaipur-302004, India.

## EXPERIMENTAL

2-Chloro-6-methoxy-4-methylquinoline (41.5 g, 0.2M) was reduced with Conc. HCl (300 ml), H<sub>2</sub>O (300 ml) and tin (40.0g) on the lines followed earlier<sup>2</sup>. The reaction mixture treated in usual manner<sup>1</sup> gave a crude solid product. The tlc examination of this crude product revealed the presence of two components. These components were separated by repeated crystallisation with petroleum ether (40-60°), one which was more soluble was identified as 6-methoxy-4-methylquinoline (18.6 g, 48%, mp. 51°)<sup>3</sup> and another which was less soluble was identified as 2,2'-bis-6-methoxy-4-methylquinoline (3). (White needles, mp. 135°, 7.71 g, 10%). Its purity was ascertained through tlc examination over Silica Gel G plates, using ethyl alcohol, ethyl acetate, formic acid (6:2:2) as mobile phase (*R<sub>f</sub>* values: 2, .78; 3, .69).

## ANALYSIS

Calcd for	C <sub>22</sub> H <sub>20</sub> O <sub>2</sub> N <sub>2</sub> (3)
	C, 76.74%; H, 5.81%; N, 8.14%
Found	C, 76.38%; H, 5.64%; N, 8.38%.

## ACKNOWLEDGEMENT

The authors thank the Head, Chemistry Department for providing the necessary facilities and Defence Research and Development Orgn., Government of India for financial assistance.

## REFERENCES

- [1] D. KISHORE, Asha SURANA and Bhuwan C. JOSHI, *Archives des Sciences*, Vol. 26, (1973).
- [2] G. I. MIKHALOV and J. GEN, Chem. USSR. 6, 511-15, (1936). Chem. Abstr. 30, 63725, 3446 (1936), Russ. Pat. 39164.
- [3] K. N. CAMPBELL, R. STAUT TIPSON, R. C. ELDERFIELD, B. K. CAMPBELL, Mary A. CLAPP, W. J. GENSHER, Dwight MARRISON and W. J. MARAN. J. Org. Chem. 11, 803-11 (1946).