Zeitschrift:	Archives des sciences et compte rendu des séances de la Société	
Herausgeber:	Société de Physique et d'Histoire Naturelle de Genève	
Band:	45 (1992)	
Heft:	1: Archives des Sciences	
Artikel:	New Hamiltonians for high-order equations of the calculus of variations : a generalization of the Dorfman approach	
Autor:	Grandville, Olivier de la	
DOI:	https://doi.org/10.5169/seals-740331	

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 13.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# NEW HAMILTONIANS FOR HIGH-ORDER EQUATIONS OF THE CALCULUS OF VARIATIONS: A GENERALIZATION OF THE DORFMAN APPROACH

BY

# **Olivier de LA GRANDVILLE\***

## ABSTRACT

Through generalized Hamiltonians, which have a direct economic interpretation, we show that we can derive in a straightforward way high-order variational equations, as well as the Ostrogradski equation. The latter is of considerable importance, since it lies at the very heart of mathematical physics: in particular, it leads to the Laplace and to the Schrödinger equations.

#### RÉSUMÉ

L'objet de cet article est de montrer comment on peut définir de nouveaux Hamiltoniens afin de pouvoir dériver très simplement les équations d'Euler-Poisson et d'Ostrogradski appartenant au calcul des variations. L'équation d'Ostrogradski est centrale à la physique mathématique, car elle conduit aux équations de Laplace et de Schrödinger.

In a remarkable paper, Robert DORFMAN [1] had shown how the maximum principle could be derived from straightforward economic reasoning, specifically from capital theory. He had shown, in particular, that differentiating a modified Hamiltonian both with respect to the state variable and the control variable led directly to the maximum principle.

In this note we would like to point out that it is possible to extend this method to obtain important building blocks of the calculus of variations, namely the Euler-Poisson and the Ostrogradski equations; the latter leads to central results in physics. We will do so by introducing Hamiltonians of increasing generality, all of which have direct economic interpretations.

# I. From the Dorfman Hamiltonian to the Euler equation

Using Dorfman's notation, recall the problem of finding a control variable  $x \equiv x(t)$  which maximizes the functional  $\int_{0}^{T} u(k, x, t) dt$  where  $k \equiv k(t)$  is a state variable, under

<sup>\*</sup> University of Geneva, 3 place de l'Université, CH-1204 Geneva, Switzerland. I wish to extend to Robert Dorfman my deepest thanks for his extremely useful comments on this paper.

the constraint  $\dot{k} = f(k, x, t)$ . One procedure, suggested by Dorfman, is to form a modified Hamiltonian  $H^* = u(k, x, t) + \frac{d}{dt} (\lambda k)$ ; equating to zero the partial derivatives of H\* with respect to x and k will lead to the simplest version of Pontryagin's maximum principle.

Remember also the link existing between the maximum principle and the calculus of variations, which enables to get in one line the Euler equation. Set the constraint  $\vec{k} = f(k, x, t)$  equal to x, as a particular case. We then have to find  $\vec{k}(t)$  (or k(t)) such that  $\int_0^T u(k, \dot{k}, t)dt$  is maximized, which is the basic problem of the calculus of variations.

Form  $H^* = u(k, x, t) + \frac{d}{dt} (\lambda k) = u(k, \dot{k}, t) + \dot{\lambda}k + \lambda \dot{k}$ .  $H^*_k = 0$  implies  $u_k + \dot{\lambda} = 0$ ;  $H^*_{k} = 0$  implies  $u_{k} + \lambda = 0$  and  $\dot{u}_{k} + \dot{\lambda} = 0$ . Hence we must have  $u_k - \frac{d}{dt} u_{k} = 0$ , which is the Euler equation<sup>1</sup>.

### II. Generalizations

This reasoning can be generalized. Let us first introduce some definitions, which will then be applied to concrete calculus of variations problems.

• *price of a variable*: we define the *price* of the  $j^{th}$  variable of a functional as the partial derivative of the functional with respect to this variable: that is the rate of increase of the functional with respect to the variable at any time.

• *value of a variable*: the value of a variable is defined as the price of the variable times the variable itself.

• *value of a system*: we define the value of a system as the total effects of the set of variables. We shall separate short-term (or direct) effects from long-term (or indirect) ones. The short-term effect is the integrand of the functional. The long-term effect is the rate of increase of the value of all variables in the system.

Concept	Definition
• price of a variable	• partial derivative of the maximum value of the functional with respect to the variable
• value of a variable	• product between the price of the variable and the variable itself
• short-term effects of a set of variables	• integrand of the functional
• long-term effects of a set of variables	<ul> <li>sum of the time derivatives of the value of each variable = derivative of the sum of the values of the variables</li> </ul>
• total effects of a set of variables = value of a system of variables	• short-term plus long-term effects of a set of variables = integrand of the functional plus the derivative of the sum of the values of the variables

The following table summarizes these definitions.

<sup>&</sup>lt;sup>1</sup> A deep account of the relationship between the classical calculus of variations and optimal control theory is given in L. PONTRYAGIN et alii [4], chapter 5. A simpler presentation will be found in I.M. GELFAND and S.V. FOMIN [3], appendix II. The reader unfamiliar with the calculus of variations will find an excellent introduction in L. ELSGOLC [2].

In order to maximise (or minimise) a functional, we will then introduce a generalized Hamiltonian which will simply be the equal to the total effects of a set of variables, and we will maximise (or minimize) this Hamiltonian with respect to each variable.

As a first example, we will consider functionals depending on high-order derivatives; we will then turn towards multiple integrals, i.e. functionals involving functions of severable variables and their partial derivatives up to the n<sup>th</sup> order.

### 1. Functionals depending on high-order derivatives

Suppose that we want to find y(x) such that

$$\int_{a}^{b} F(x, y, y', ..., y^{(j)}, ..., y^{(n)}) dx$$

is maximized  $(y^{(j)} \text{ denotes the } j^{th} \text{ derivative of the function } y(x)$ ; by convention we set  $y^{(0)} \equiv y(x)$ ).

In this problem we define a system of n+1 prices, each corresponding to the n+1 variables  $y^{(0)}, ..., y^{(n)}$ :

$$\lambda j(x) = \frac{\partial}{\partial y^{(j)}} \int_a^b F(x, y, y', ..., y^{(n)}) dx, j = 0, ..., n$$

The value of each variable, i.e. the product of the unit value of the variable and the magnitude of the variable, is equal to  $\lambda j(x) \cdot y^{(j)}$ .

The short-term effects of the set of variables  $y, ..., y^{(n)}$  is just  $F(x, y, y', ..., y^{(n)}, and$ the long term effect is  $\frac{d}{dx} \left[\sum_{i=0}^{n-1} \lambda j(x) y^{(j)}\right]$ .

Hence the value of the system of variables for any x is equal to: 
$$H^{n} = F(x, y, y', y', y')$$
  

$$y^{(n)} + \frac{d}{dx} \left[\sum_{j=0}^{n-1} \lambda_{j}(x)y^{(j)}\right]$$

$$= (x, y, y', ..., y^{(n)}) + \sum_{j=0}^{n-1} [\lambda y_{j}^{(j+1)} + \lambda 'y_{j}^{(j)}]$$

$$= F(x, y, y', ..., y^{(n)}) + \lambda_{0}y' + \lambda_{0}'y + \lambda_{1}y'' + \lambda_{1}'y' + ...$$

$$+ \lambda_{j-1}y^{(j)} + \lambda '_{j-1}y^{(j-1)} + \lambda_{j}y^{(j+1)} + \lambda_{j}'y^{(j)} + ...$$

$$+ \lambda_{n-1}y^{(n)} + \lambda '_{n-1}y^{(n-1)}$$

Each of the components of  $\frac{d}{dx}(\lambda_j y^{(j)})$  is equal to  $\lambda_j y^{(j+1)} + \lambda'_j y^{(j)}$ : it is therefore made up of the rate of increase of  $y^{(j)}$  multiplied by its price, and of the price increase of  $y^{(j)}$ multiplied by  $y^{(j)}$  (this latter part corresponds to the "capital gain" in the Dorfman interpretation of his modified Hamiltonian).

Let us now derive the total value of the system H<sup>n</sup> partially with respect to each variable y, y', ...,  $y^{(n)}$  and set each derivative equal to 0. We then have the following system of n+1 differential equations:

...,

$$H_{y}^{n} = F_{y} + \lambda_{0}^{'} = 0$$

$$H_{y'}^{n} = F_{y'} + \lambda_{0} + \lambda_{1}^{'} = 0$$

$$\cdots$$

$$H_{y(j)}^{n} = F_{y(j)} + \lambda_{j-1} + \lambda_{1}^{'} = 0$$

$$\cdots$$

$$H_{y(n-1)}^{n} = F_{y(n-1)} + \lambda_{n-2} + \lambda_{n-1}^{'} = 0$$

$$H_{y(n)} = F_{y(n)} + \lambda_{n-1} = 0$$

Differentiate each equation  $H_{y(j)}^{n} = 0$  (j = 0, ..., n) j times with respect to x. We get:

(3)  

$$F_{y} + \lambda'_{0} = 0$$

$$\frac{d}{dx} F_{y'} + \lambda'_{0} + \lambda''_{1} = 0$$

$$\dots$$

$$\frac{d}{dx^{(j)}} F_{y^{(j)}} + \lambda^{(j)}_{j-1} + \lambda^{(j+1)}_{j} = 0$$

$$\dots$$

$$\frac{d^{(n-1)}}{dx^{(n-1)}}F_{y_{(n-1)}} + \lambda \frac{(n-1)}{n-2} + \lambda \frac{(n)}{n-1} = 0$$
$$\frac{d^{(n)}}{dx^{(n)}}F_{y_{(n)}} + \lambda \frac{(n)}{n-1} = 0$$

We can now write, from system (3):

. . .

$$\lambda \binom{(n)}{n-1} = -\frac{d(n)}{dx(n)} F_{y(n)}$$

$$\lambda \binom{(n-1)}{n-2} = -\frac{d(n-1)}{dx(n-1)} F_{y(n-1)} + \frac{d(n)}{dx(n)} F_{y(n)}$$

$$\lambda \binom{(n-2)}{n-3} = -\frac{d(n-2)}{dx(n-2)} F_{y(n-2)} + \frac{d(n-1)}{dx(n-1)} - \frac{d(n)}{dx(n)} F_{y(n)}$$

$$\dots$$

$$(4) \qquad \lambda \binom{(n-i+1)}{n-i} = (-1) \frac{d(n-i+1)}{dx(n-i+1)} F_{y(n-i+1)} + (-1)^2 \frac{d(n-i+2)}{dx(n-i+2)} F_{y(n-i+2)}$$

$$+ \dots + (-1)^{i} \frac{d(n)}{dx(n)} F_{y(n)} = \int_{2}^{2} \int_{2}^{1} (-1)^{j} \frac{d(n-i+j)}{dx(n-i+j)} F_{y(n-i+j)}$$

Hence, the variable  $\lambda'_0$  appearing in the first equation of system (3), and which corresponds to i = n in (4), is equal to

$$\lambda'_0 = \sum_{j=1}^n (-1)^j \frac{d^{(j)}}{dx^{(j)}} F_{y(j)}$$

We then get, from the first equation of (3), the Euler-Poisson equation which reads:

$$F_{y} + \sum_{j=1}^{n} (-1)^{j} \frac{d^{(j)}}{dx^{(j)}} F_{y(j)} = 0$$

### 2. Functionals defined by multiple integrals

This kind of economic reasoning can be further extended to functionals of the form

$$\int \dots \int_{R} F[x_{1}, \dots, x_{n}; z; z_{x_{1}}, \dots, z_{x_{n}}] dx_{1}, \dots, dx_{n}.$$

which have a considerable importance in physics, and which can be encountered in economics as well (for instance in dynamic location problems, or in stochastic optimization).

In order to get the Ostrogradski formula which sets a first order condition for  $z(x_1, ..., x_n)$  to be an extremal of this functional, let us form a general Dorfman-type Hamiltonian

$$H^{G} = F[x_{1}, ..., x_{n}; z; z_{x_{1}}, ..., z_{x_{n}}] + \sum_{i=1}^{n} \frac{\partial}{\partial x_{i}} [\lambda (x_{1}, ..., x_{n}) \cdot z(x_{1}, ..., x_{n})]$$

where  $\lambda$  ( $x_1, ..., x_n$ ) is defined by

$$\lambda(x_1, \dots, x_n) = \frac{\partial}{\partial z} \int \dots \int_{\mathbf{R}} F(x_1, \dots, x_n; z; z_{x_1}, \dots, z_{x_n}) dx_1, \dots, d_{x_n}.$$

As before,  $\lambda$  can be interpreted as the price of z. This Hamiltonian can conveniently be written as

$$H^{G} = F[x_{1}, ..., x_{n}; z; z_{x_{1}}, ..., z_{x_{n}}] + z \sum_{i=1}^{n} \lambda_{x_{i}} + \lambda \sum_{i=1}^{n} z_{x_{i}}$$

Differentiate  $H^G$  with respect to z and  $z_{x_1}$ , ...,  $z_{x_n}$ , and set the n+1 resulting partial derivatives to zero. We have:

(5)  $H_{z}^{G} = F_{z} + \lambda_{x_{I}} + \dots + \lambda_{x_{n}} = 0$  $H_{z_{x_{i}}}^{G} = F_{z_{x_{i}}} + \lambda = 0 \quad , i=1, \dots, n$ 

Differentiate each equation  $H_{z_{x_i}}^G = 0$  with respect to  $x_i$  (i=1, ..., n).

We get:

$$H_z^0 = F_z + \lambda_{x_1} + \dots + \lambda_{x_n} = 0$$

(6)

$$\frac{\partial}{\partial x_i} H^G_{z_{x_i}} = \frac{\partial}{\partial x_i} F_{z_{x_i}} + \lambda_{x_i} = 0 \quad , i = 1, ..., n$$

System (6) entails the Ostrogradski equation:

(7) 
$$F_z - \sum_{i=1}^n \frac{\partial}{\partial x_i} F_{z_{x_i}} = 0$$

from which we can obtain the Laplace and the Schrödinger equations.

It is possible to generalize this method to functionals involving functions of n variables and their partial derivatives up to the m<sup>th</sup> order. We are then led to a generalization of both the Euler-Poisson and the Ostrogradski equations. For simplicity, we will treat the case n = m = 2. We have to find z that maximizes the functional

$$\iint_{D} F(x_{1}, x_{2}; z; z_{x_{1}}, z_{x_{2}}; z_{x_{1}x_{1}}, z_{x_{1}x_{2}}, z_{x_{2}x_{2}}) dx_{1} dx_{2}$$

denoted  $\iint_{-} F(\dots) dx_1 dx_2$  for short.

We introduce a Dorfman-type Hamiltonian as follows: let

$$\lambda^0 = \frac{\partial}{\partial z} \iint_D F(\dots) dx_1 dx_2$$

and  $\lambda^{j} = \frac{\partial}{\partial z_{x_{j}}} \int \int F(\dots) dx_{1} dx_{2}$ , j = 1, 2

define the marginal contributions to the functional (or the prices) of z,  $z_{x_1}$  and  $z_{x_2}$ , respectively. A general Hamiltonian can then be written as follows:

(8) 
$$H^{2,2} = F(x_{1}, x_{2}; z; z_{x_{1}}, z_{x_{2}}; z_{x_{1}x_{1}}, z_{x_{1}x_{2}}, z_{x_{2}x_{2}}) + \frac{2}{i}\sum_{i=1}^{2} \frac{\partial}{\partial x_{i}} [\lambda^{0}z] + \sum_{j=1}^{2} \sum_{i=1}^{2} \frac{\partial}{\partial x_{i}} [\lambda^{j}z_{x_{j}}]$$
$$= F(...) + \lambda^{0}_{x_{1}}z + \lambda^{0}z_{x_{1}} + \lambda^{0}_{x_{2}}z + \lambda^{0}z_{x_{2}} + \frac{1}{\lambda^{1}z_{x_{1}x_{1}}} + \lambda^{1}z_{x_{1}x_{1}} + \lambda^{1}z_{x_{2}x_{1}} + \lambda^{1}z_{x_{1}x_{2}} + \frac{1}{\lambda^{2}z_{x_{2}}}z_{x_{2}} + \lambda^{2}z_{x_{2}x_{2}} + \lambda^{2}z_{x_{2}x_{2}} + \lambda^{2}z_{x_{2}x_{2}}z_{x_{2}} + \lambda^{2}z_{x_{2}x_{2}}z_{x_{2}}z_{x_{2}} + \lambda^{2}z_{x_{2}x_{2}}z_{x_{$$

(the first superscript in  $H^{2,2}$  refers to the number of variables of z; the second one indicates the highest order of the derivatives in the functional).

Differentiating  $H^{2,2}$  with respect to z,  $z_{x_1}$ ,  $z_{x_2}$ ,  $z_{x_1x_1}$ ,  $z_{x_1x_2}$ ,  $z_{x_2x_2}$ , we get:

(9a) 
$$H_{z}^{2,2} = F_{z} + \lambda \frac{0}{x_{1}} + \lambda \frac{0}{x_{2}} = 0$$

(9b) 
$$H_{z_{x_{l}}}^{2,2} = F_{z_{x_{l}}} + \lambda^{0} + \lambda_{x_{l}}^{1} + \lambda_{x_{2}}^{0} = 0$$

(9c) 
$$H_{z_{x_{2}}}^{2,2} = F_{z_{x_{2}}} + \lambda^{0} + \lambda_{x_{1}}^{2} + \lambda_{x_{2}}^{2} = 0$$

(9d) 
$$H_{z_{x_{l}x_{l}}}^{2,2} = F_{z_{x_{l}x_{l}}} + \lambda^{l} = 0$$

(9e) 
$$H_{z_{x_{l}x_{2}}}^{2,2} = F_{z_{x_{l}x_{2}}} + \lambda^{l} + \lambda^{2} = 0$$

(9f) 
$$H_{z_{x_2}x_2}^{2,2} = F_{z_{x_2}x_2} + \lambda^2 = 0$$

Differentiate (9b) and (9c) with respect to  $x_1$  and  $x_2$ , respectively; differentiate twice (9d), (9e) and (9f) with respect to  $x_1$ ,  $x_1$  and  $x_2$ , and  $x_2$ , respectively. We then have the following system:

(10) 
$$F_{z} + \lambda_{x_{1}}^{0} + \lambda_{x_{2}}^{0} = 0$$
$$\frac{\partial}{\partial x_{1}} F_{z_{x_{1}}} + \lambda_{x_{1}}^{0} + \lambda_{x_{1}x_{1}}^{1} + \lambda_{x_{1}x_{2}}^{1} = 0$$
$$\frac{\partial}{\partial x_{i}} F_{z_{x_{2}}} + \lambda_{x_{2}}^{0} + \lambda_{x_{1}x_{2}}^{2} + \lambda_{x_{2}x_{2}}^{2} = 0$$
$$\frac{\partial^{2}}{\partial x_{1}^{2}} F_{z_{x_{1}x_{1}}} + \lambda_{x_{1}x_{1}}^{1} = 0$$
$$\frac{\partial^{2}}{\partial x_{1}\partial x_{2}} F_{z_{x_{1}x_{2}}} + \lambda_{x_{1}x_{2}}^{1} + \lambda_{x_{1}x_{2}}^{2} = 0$$
$$\frac{\partial^{2}}{\partial x_{2}^{2}} F_{z_{x_{2}x_{2}}} + \lambda_{x_{2}x_{2}}^{2} = 0$$

System (10) yields the fourth-order partial differential equation

(11)  

$$F_{z} - \frac{\partial}{\partial x_{I}} F_{z_{x_{I}}} - \frac{\partial}{\partial x_{2}} F_{z_{x_{2}}} + \frac{\partial^{2}}{\partial x_{I}^{2}} F_{z_{x_{I}}x_{I}} + \frac{\partial^{2}}{\partial x_{I}\partial x_{2}} F_{z_{x_{I}}x_{2}} + \frac{\partial^{2}}{\partial x_{2}^{2}} F_{z_{x_{2}}x_{2}} = 0$$

which is a general Ostrogradski equation.

We have thus shown that the Dorfman's modified Hamiltonian approach is perfectly general: all the fundamental equations of the calculus of variations can be derived from extensions of that approach.

#### REFERENCES

- [1] R. DORFMAN, "An Economic Interpretation of Optimal Control Theory", American Economic Review, 1969.
- [2] L. ELSGOLC, Calculus of Variations, Reading, Mass.: Addison Wesley, 1962.
- [3] I. GELFAND and S. FOMIN, Calculus of Variations, Englewood Cliffs, N.J. Prentice Hall, 1963.
- [4] L. PONTRYAGIN, BOLTYANSKII, V., GAMKRELIDZE, R. and MISHCHENKO, E., The Mathematical Theory of Optimal Processes, Interscience Publisher, New York, 1962.

Manuscrit reçu le 10 mars 1992.