

# Coexistence of *Conochilus unicornis* with *C. hippocrepis* in Lake Geneva : and commensal organisms found in their gelatinous matrix

Autor(en): **Balvay, Gérard / Druart, Jean-Claude**

Objektyp: **Article**

Zeitschrift: **Archives des sciences et compte rendu des séances de la Société**

Band (Jahr): **48 (1995)**

Heft 3: **Archives des Sciences**

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-740260>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Archs Sci. Genève	Vol. 48	Fasc. 3	pp. 257-260	Décembre 1995
-------------------	---------	---------	-------------	---------------

## COEXISTENCE OF *CONOCHILUS UNICORNIS* WITH *C. HIPPOCREPIS* IN LAKE GENEVA, AND COMMENSAL ORGANISMS FOUND IN THEIR GELATINOUS MATRIX

BY

**Gérard BALVAY\* & Jean-Claude DRUART\***

*(Ms soumis le 27.09.1995, accepté le 3.11.1995)*

### ABSTRACT

**Coexistence of *Conochilus unicornis* with *C. hippocrepis* in Lake Geneva, and commensal organisms found in their gelatinous matrix.**- Coexistence of *Conochilus unicornis* with *C. hippocrepis* is confirmed in Lake Geneva. A number of commensal organisms in the gelatinous matrix of these colonial *Conochilus* was observed for the first time in Lake Geneva in 1992. Such colonies were intensively green in color from *Chlamydomonas gloeophila* var. *irregularis* and *Phormidium arcuatum* in May-June, becoming progressively colourless on and after mid-July.

**Key-words:** Lake Geneva, commensalism, *Conochilus unicornis*, *Conochilus hippocrepis*, *Phormidium arcuatum*, *Chlamydomonas gloeophila* var. *irregularis*.

### *CONOCHILUS* IN LAKE GENEVA

Lake Geneva is inhabited by two species of *Conochilus*, with *C. unicornis* Rousselet 1892, being more abundant than *C. hippocrepis* (Schrank, 1830). *Conochilus hippocrepis* was first recorded by IMHOF (1885). However, WEBER (1898) found only *C. unicornis* and concluded that Imhof confused *C. unicornis* with *C. hippocrepis*. In fact, these two species of *Conochilus* coexist in Lake Geneva as they do, for example, in Lake Glubokoe (MATVEEVA, 1986).

There are still some taxonomic problems within the genus *Conochilus*, due to the existence of intermediate forms between *C. unicornis* and *C. hippocrepis* (PEJLER, 1956). These two species are the two clearly distinguishable extreme members of the series. Between them there are transitional forms, both in the size and fusion of antennae (from completely fused, or nearly so, in *C. unicornis* to completely separate, or only slightly fused, in *C. hippocrepis*) as well as in the ratio of body size to length of foot (PEJLER, 1957; POURRIOT, 1965).

---

\* INRA, Institut de Limnologie, Station d'Hydrobiologie Lacustre, B.P. 511, F-74203 Thonon-les-Bains Cedex, France.

Most of our individual specimens could be rather securely assigned to one or the other of the two species, but sometimes the character of the antennae was not clearly definable in some of the preserved specimens. Moreover, considerable variation in size of colony and number of individuals occurs in colonies of *Conochilus*, with colonies intermediate in some respects between *C. unicornis* et *C. hippocrepis* (Table I).

TABLE I

Comparative data between *Conochilus unicornis* and *C. hippocrepis*

	<i>Conochilus unicornis</i>	<i>Conochilus hippocrepis</i>	Authors
Individual size ( $\mu\text{m}$ )	200-450	400-800	STEMBERGER, 1979
	250-380	500-800	RUDESCU, 1960
	320	500-600	BARTOS, 1959
	212-303	310-455	Present study
Size of colonies (mm)	0.5-3	2-4	KOSTE, 1978
	0.56-1.00	2-4	BARTOS, 1959
	0.80-1.00	4	RUDESCU, 1960
	0.40-0.76	0.52-0.98	Present study
Number of individuals in colonies		10-161	EDMONDSON & LITT, 1987
	2-25	60-100	BARTOS, 1959
	5-12	30-60	STEMBERGER, 1979
	5-25	30-200	KOSTE, 1978
	5->150	25->200	WALLACE, 1987
	4-38	12-98	Present work

### COMMENSAL SPECIES IN THE GELATINOUS MATRIX

Colonies of *Conochilus* are a normal component of zooplankton in Lake Geneva. In 1992, they became extraordinarily abundant in May and June (BALVAY, 1993). This period marks the first time that they were very visually obvious in zooplankton samples due to their green coloring induced by commensal species located in the gelatinous matrix of the colonies of *Conochilus*. On and after mid-July, fewer and fewer colonies were coloured and in mid-September all of them were colourless. At the same time, the same phenomenon was observed by our Swiss colleagues J. Naef and R. Revaclier in the "Petit Lac" (western part of Lake Geneva).

Microscopic examination of living samples revealed coexistence of colonies of *C. unicornis* and *C. hippocrepis* in which the gelatinous matrix was more or less intensively colonized with commensal algae.

Several kinds of symbiotic algae and protozoa have been seen growing in the matrix of *Conochilus* colonies (Table II). During the 1992 occurrence of *Conochilus* in Lake Geneva, two commensal species were observed: *Chlamydomonas gloeophila* var. *irregularis* Skuja (first record in Lake Geneva) and *Phormidium arcuatum* Skuja.

TABLE II

Organisms found in the gelatinous matrix of *Conochilus*

	<i>Conochilus unicornis</i>	<i>Conochilus hippocrepis</i>	<i>Conochilus</i> sp
CYANOPHYTA			
<i>Phormidium arcuatum</i> Skuja	12	12	5, 10
<i>Phormidium mucicola</i> Huber-P. et Naumann	1		3, 10
<i>Lyngbia endophytica</i> Elenkin et Hollerbach	7		
CHRYSOPHYTA			
	7		
EUCHLOROPHYTA			
<i>Carteria conochili</i> Skuja		9, 11	
<i>Chlamydomonas</i> sp	7		
<i>Chlamydomonas gloeophila</i> Skuja		9	
<i>C. gloeophila</i> var. <i>irregularis</i> Skuja	12	12	
PROTOZOA			
<i>Amoeba gracilis</i> Greeff	7		
<i>Amoeba mucicola</i> de Graff	7		
<i>Amoeba radiosa</i> Dujardin	7		
<i>Vorticella conochili</i> Stokes	6, 7, 8		
<i>Vorticella conosoma</i> Stokes	6, 7	2	
<i>Vorticella dimorpha</i> Stiller	4		
<i>Vorticella</i> sp	12	12	
1. HUBER-PESTALOZZI & NAUMANN, 1929	7. DE GRAFF, 1953		
2. KAHL, 1930	8. MORAVCOVA, 1956		
3. HUBER-PESTALOZZI, 1938	9. HUBER-PESTALOZZI, 1961		
4. STILLER, 1940	10. STARMACH, 1966		
5. SKUJA, 1948	11. ETTL, 1983		
6. FAURÉ-FREMIET, 1949	12. Present work		

*Chlamydomonas gloeophila* var. *irregularis* generally exhibits two different shapes: a rounded form and an elongated form. In Lake Geneva, colonies of *Conochilus* often were colonized with elongated forms.

Gelatinous matrix is seen only on freshly collected living colonies. The mucous is destroyed in preserved samples in which only solitary individuals may be observed. This fact could explain why *Phormidium arcuatum* and *P. mucicola* were previously found in some phytoplankton samples (BALVAY *et al.*, 1990; DRUART *et al.*, 1983).

#### ACKNOWLEDGMENTS

We wish to thank W. Koste who confirmed the identification of *C. hippocrepis*, J. Naef and R. Revaclier for their informations, and G. Chapuis for her technical assistance. The present work was supported by the "Commission Internationale pour la Protection des Eaux du Léman" (CIPEL) and the "Institut National de la Recherche Agronomique" (INRA).

## RÉSUMÉ

La coexistence de *Conochilus unicornis* avec *C. hippocrepis* a été observée dans le Léman en 1992. Plusieurs organismes commensaux ont été identifiés dans la matrice gélatineuse de ces rotifères dont les colonies, fortement colorées en vert par *Chlamydomonas gloeophila* var. *irregularis* et *Phormidium arcuatum* en Mai-Juin, sont devenues progressivement incolores après la mi-Juillet.

## REFERENCES

- BALVAY, G. 1993. Evolution du zooplancton du Léman. *Rapp. Comm. int. prot. eaux Léman contre pollut., Campagne 1992*: 97-112.
- BALVAY, G., J.C. DRUART & M. LAURENT. 1990. Deuxième complément à l'inventaire du plancton du Léman. *Archs Sci. Genève* 43: 159-166.
- BARTOS, E. 1959. Virnici - Rotatoria. *Fauna CSR* 15, 969 pp.
- DRUART, J.C., E. PONGRATZ & R. REVACLIER. 1983. Les algues planctoniques du Léman: historique et inventaire. *Schweiz. Z. Hydrol.* 45: 430-457.
- EDMONDSON, W.T. & A.H. LITT. 1987. *Conochilus* in Lake Washington. *Hydrobiologia* 147: 157-162.
- ETTL, H. 1983. Chlorophyta 1. Phytomonadina. Süßwasserflora von Mitteleuropa, Gustav Fischer Verlag (Ed.) Stuttgart, 9: 807 pp.
- FAURE-FREMIET, E. 1949. Deux espèces de *Vorticella* commensales des *Conochilus*. *Hydrobiologia* 1: 126-132.
- GRAAF, F. DE 1953. Commensal organisms in the colonies of *Conochilus unicornis* Rousselet. With a note on *Phormidium mucicola* Huber et Naumann. *Hydrobiologia* 5: 390-399.
- HUBER-PESTALOZZI, G. 1938. Das Phytoplankton des Süßwassers. *Die Binnengewässer* 16 (1): 342 pp.
- HUBER-PESTALOZZI, G. 1961. Das Phytoplankton des Süßwassers. *Die Binnengewässer* 16 (5): 744 pp.
- HUBER-PESTALOZZI, G. & NAUMANN, E. 1929. *Phormidium mucicola* Huber et Naumann, ein epibiont in der Gallerte pflanzlicher und tierischer Planktonorganismen. *Ber. deutsch. Botan. Ges.* 47: 67-76.
- IMHOF, O.E. 1885. Weitere Mittheilung über die pelagische und Tiefseefauna der Süßwasserbecken. *Zool. Anz.* 8 (190): 160-163.
- KAHL, A. 1930. Wimpertiere oder Ciliata (Infusoria). *Die Tierwelt Deutschlands* 18 (1): 886 pp.
- KOSTE, W. 1978. Rotatoria. Die Rädertiere Mitteleuropas. Gebrüder Borntraeger (Ed.) Berlin, Stuttgart, vol 1: 673 pp, vol. 2: 442 pp.
- MATVEEVA, L.K. 1986. Pelagic rotifers of Lake Glubokoe from 1897 to 1984. *Hydrobiologia* 141: 45-54.
- MORAVCOVA, V. 1956. Another rediscovery of commensal *Vorticella* upon *Conochilus* in Czechoslovakia. *Hydrobiologia* 8: 181-183.
- PEJLER, B. 1956. Introgression in planktonic Rotatoria with some points of view on its causes and conceivable results. *Evolution* 10 (3): 246-261.
- PEJLER, B. 1957. On variation and evolution in planktonic rotatoria. *Zool. Bidr. Upps.* 32: 1-66.
- POURRIOT, R. 1965. Notes taxonomiques sur quelques Rotifères planctoniques. *Hydrobiologia* 26: 579-604.
- RUDESCU, L. 1960. Trochelminthes, Rotatoria. *Fauna Republicii Populare Romine* 2 (2): 1192 pp.
- SKUJA, H. 1948. Taxonomie des Phytoplanktons einiger Seen in Uppland, Schweden. *Symb. Bot. Upsal.* 9 (3): 399 pp.
- STARMACH, K. 1966. Cyanophyta-Sinice, Glaucophyta-Glaukofity. *Flora Slodk. Polski* 2: 807 pp.
- STEMBERGER, R.S. 1979. A guide to Rotifers of the Laurentian Great Lakes. EPA-600/4-79-021, 186 pp.
- STILLER, J. 1940. Beitrag zur Peritrichenfauna des Grossen Plöner Sees in Holstein. *Arch. f. Hydrobiol.* 36: 263-285.
- WALLACE, R.L. 1987. Coloniality in the phylum Rotifera. *Hydrobiologia* 147: 141-155.
- WEBER, E.F. 1898. Faune rotatorienne du bassin du Léman. *Rev. suisse Zool.* 5: 263-785.