

The Anâtaxis phylogenetic method. 2. An exemple : reconstituting a whole dendrogram

Autor(en): **Bittar, Gabriel**

Objekttyp: **Article**

Zeitschrift: **Archives des sciences et compte rendu des séances de la Société**

Band (Jahr): **55 (2002)**

Heft 1

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-740288>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THE ANÂTAXIS PHYLOGENETIC METHOD. 2. AN EXAMPLE - RECONSTITUTING A WHOLE DENDROGRAM

BY

Gabriel BITTAR

(Ms. reçu le 3.1.2002, accepté le 23.5.2002)

ABSTRACT

The Anâtaxis phylogenetic method. 2. An example - reconstituting a whole dendrogram. - The WPGMA numerical taxonomy phenetics method is applied to the semi-matrix of dissimilarities (between terminal taxa) derived from a matrix of characters states to which a cladistic maximum parsimony method had also been applied. Because this semi-theoretical case displays homoplasy and lineage-dependent heterogeneity of transformation rates, the phenogram and the cladogram thus produced are phyletically quite different. The new phylogenetic method Anâtaxis is then applied to the same semi-matrix of dissimilarities, and it rapidly reconstitutes a dendrogram which is congruent with the cladogram.

Key-words: Cladistic Maximum Parsimony, Dissimilarity matrix, Evolutionary tree, Heterogeneity of transformation rates, Homoplasy, Numerical Taxonomy Phenetics, Outgroup-based method, Phylogenetic method, Splitting method.


We wish to phyletically study a number of taxa selected from within the **Craniata** phylum, more precisely we want to resolve the phyletic relationships within the **Vertebrata** sub-phylum. Accordingly, we use the **myxini taxon**, which is craniate but non-vertebrate, **as starting outgroup**.

taxons	caractères																																						
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36			
a_myxini	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
b_petromyzones	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
c_selacii	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
d_amia	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
e_teleostei	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
f_dipnoi	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
g_urodela	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
h_anura	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
i_sauria	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
j_ophidia	+	+	+	+	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
k_chelonia	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
l_crocodylia	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
m_aves	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
n_chiroptera	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
o_felidae	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
p_marsupialia	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

This is an *ad hoc* matrix, with characters and taxa chosen for demonstration purposes, and with no pretense at producing a taxonomically or systematically valid phyletic tree. Rather, this matrix was designed to include the main pitfalls associated in a phyletic reconstruction, and to demonstrate the validity and effectiveness of the Anâtaxis method. The characters are binary (presence/absence) morphological ones, and their definition might be obtained through the author.

¹ Universities of Geneva and Lausanne, po box 281, American River, Kangaroo Island, South Australia 5221, bittar@isb-sib.ch

With a Cladistic Maximum Parsimony (CMP) method (SWOFFORD, 1993), i.e. by looking for the tree that minimalises the sum of dissimilarities between all pairs of adjacent nodes, and with a post hoc rooting (R) done within the branch leading to the a_myxini node, we obtain the following phylogram, i.e. a cladogram in which the length of each branch is proportional to the dissimilarity between the two nodes which this branch connects (for the DelTran option, the transformations of characters 7 and 11-13 are done one branch later, in the Tetrapoda branch). The total length of this cladistically most parsimonious tree is 39 steps (all the steps are indicated above or below each branch), with a total of 5 homoplasies.



Now here is a phenogram, i.e. a tree produced by a Numerical Taxonomic Phenetics -NTP- method (SNEATH & SOKAL, 1973). It is the product of an application, on the Δ matrix of dissimilarities between pairs of taxa, of the clustering algorithm **WPGMA** (Weighted-Pair Group Method with Arithmetic averages) - and again, with a post hoc rooting (R) done within the branch leading to a_myxini.

The phenogram differs strongly from the cladogram, because, as a product of an NTP method, it is very sensitive to the lineage-dependent heterogeneity of evolutionary rates, and does not take into account homoplasy. Because of the high number of autapomorphies within the m_aves branch, the birds have been rejected to a basal position within the Amniota phenon; this adds two homoplasies to the tree. Because of the high number of autapomorphies within the Amniota branch, it has been rejected to a basal position relative to the f_dipnoi taxon and the Amphibia; this adds two homoplasies to the tree. And because of the high number of autapomorphies within the branch leading to the f_dipnoi and Tetrapoda, the remaining fish are found assembled together within a Pisces phenon; this adds one homoplasy to the tree.

Iteration 1.

With indices o for myxini and i for any vertebrate, median-normalise on the a_myxini taxon (median = 15.5) the Vertebrata Δ sub-matrix:

(in bold) $\text{diff}_{oi} = \text{med}(\Delta_{oi}) - \Delta_{oi}$

Δ		b	c	d	e	f	g	h	i	j	k	l	m	n
o=myxini	diff	14	11	10	10	5.5	2	0	-3	0	-3	-4	-8	-6
a_myxini	0	1.5	4.5	5.5	5.5	10	13.5	15.5	18.5	15.5	18.5	19.5	23.5	21.5
b_petromyzones	14		3.5	4.5	4.5	9	12.5	14.5	17.5	14.5	17.5	18.5	22.5	20.5
c_selacii	11			3.5	3.5	7	11.5	13.5	16.5	13.5	16.5	17.5	21.5	19.5
d_amia	10				2.5	7	8.5	12.5	15.5	12.5	15.5	16.5	20.5	18.5
e_teleostei	10					7	10.5	12.5	15.5	12.5	15.5	16.5	20.5	18.5
f_dipnoi	5.5						5	7	10	10	10	11	15	13
g_urodela	2							4	8	9	7	8	12	10
h_anura	0								8	11	7	8	12	10
i_sauria	-3									3	3	4	8	6
j_ophidia	0										6	7	11	9
k_chelonia	-3											3	7	5
l_crocodylia	-4												4	6
m_aves	-8													8

Vertebrata Δ^* sub-matrix median-normalised on taxon myxini,

with $\Delta_{ij}^* = \Delta_{ij} + \text{diff}_{oi} + \text{diff}_{oj}$:

Δ^*		b	c	d	e	f	g	h	i	j	k	l	m	n
o=myxini	diff	14	11	10	10	5.5	2	0	-3	0	-3	-4	-8	-6
a_myxini	0	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5	15.5
b_petromyzones	14		28.5	28.5	28.5	28.5	28.5	28.5	28.5	28.5	28.5	28.5	28.5	28.5
c_selacii	11			24.5	24.5	23.5	24.5	24.5	24.5	24.5	24.5	24.5	24.5	24.5
d_amia	10				22.5	22.5	20.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5
e_teleostei	10					22.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5
f_dipnoi	5.5						12.5	12.5	12.5	15.5	12.5	12.5	12.5	12.5
g_urodela	2							6	7	11	6	6	6	6
h_anura	0								5	11	4	4	4	4
i_sauria	-3									0	-3	-3	-3	-3
j_ophidia	0										3	3	3	3
k_chelonia	-3											-4	-4	-4
l_crocodylia	-4												-8	-4
m_aves	-8													-6

It is straightforward to define the sub-outgroup. Let us call **Gnathostomata** the non-lamprey vertebrates and gnathostomeX any member of this taxon. When analysing all the triads formed within the myxini-normalised Vertebrata ingroup, we notice that ALWAYS $\Delta^*(\text{petromyzones}; \text{gnathostome1}) = 28.5 = \Delta^*(\text{petromyzones}; \text{gnathostome2}) \gg \Delta^*(\text{gnathostome2}; \text{gnathostome1}) \{-8 \text{ to } 24.5\}$: all the triads involving petromyzones are ultrametric-like, and the lampreys taxon appears as the external one in all these triads.

Thus, within the Vertebrata subset, the b_petromyzones taxon is outgroup to the Gnathostomata ingroup.

Iteration 2.

Median-normalise on the b_petromyzones taxon (median = 14.5) the Gnathostomata sub-matrix of original Δ :

Δ		c	d	e	f	g	h	i	j	k	l	m	n
o=petromyzones	diff	11	10	10	5.5	2	0	-3	0	-3	-4	-8	-6
b_petromyzones	0	3.5	4.5	4.5	9	12.5	14.5	17.5	14.5	17.5	18.5	22.5	20.5
c_selacii	11		3.5	3.5	7	11.5	13.5	16.5	13.5	16.5	17.5	21.5	19.5
d_amia	10			2.5	7	8.5	12.5	15.5	12.5	15.5	16.5	20.5	18.5
e_teleostei	10				7	10.5	12.5	15.5	12.5	15.5	16.5	20.5	18.5
f_dipnoi	5.5					5	7	10	10	10	11	15	13
g_urodela	2						4	8	9	7	8	12	10
h_anura	0							8	11	7	8	12	10
i_sauria	-3								3	3	4	8	6
j_ophidia	0									6	7	11	9
k_chelonia	-3										3	7	5
l_crocodylia	-4											4	6
m_aves	-8												8

Gnathostomata Δ^* median-normalised on taxon b_petromyzones :

Δ^*		c	d	e	f	g	h	i	j	k	l	m	n
o=petromyzones	diff	11	10	10	5.5	2	0	-3	0	-3	-4	-8	-6
b_petromyzones	0	14.5	14.5	14.5	14.5	14.5	14.5	14.5	14.5	14.5	14.5	14.5	14.5
c_selacii	11		24.5	24.5	23.5	24.5	24.5	24.5	24.5	24.5	24.5	24.5	24.5
d_amia	10			22.5	22.5	20.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5
e_teleostei	10				22.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5	22.5
f_dipnoi	5.5					12.5	12.5	12.5	15.5	12.5	12.5	12.5	12.5
g_urodela	2						6	7	11	6	6	6	6
h_anura	0							5	11	4	4	4	4
i_sauria	-3								0	-3	-3	-3	-3
j_ophidia	0									3	3	3	3
k_chelonia	-3										-4	-4	-4
l_crocodylia	-4											-8	-4
m_aves	-8												-6

Let us call **Osteomorphi** the non-selacii gnathostomes. We notice the clear-cut candidacy of the selacii taxon as the next sub-outgroup: with the exception of when the f_dipnoi taxon is involved, there is an ultrametric-like triadic relationship, $\Delta^*(\text{selacii}; \text{non-dipnoi osteomorph1}) = 24.5 = \Delta^*(\text{selacii}; \text{non-dipnoi osteomorph2}) \gg \Delta^*(\text{non-dipnoi osteomorph2}; \text{non-dipnoi osteomorph1}) \{-8 \text{ to } 22.5\}$.

Otherwise, when the f_dipnoi taxon is involved, we have $\Delta^*(\text{selacii}; \text{non-dipnoi osteomorph}) = 24.5 > \Delta^*(\text{selacii}; \text{dipnoi}) = 23.5 \gg \Delta^*(\text{non-dipnoi osteomorph}; \text{dipnoi}) \{12.5 \text{ to } 22.5\}$. (...)

The c_selacii taxon appears as the external one in all triads formed from within the lampreys-normalised Gnathostomata subset. Thus, within this subset, the taxon c_selacii is outgroup to the Osteomorphi ingroup.

REFERENCES

- BALIC ŽUNIČ, T. & I. VICKOVIĆ. 1996. IVTON - program for the calculation of geometrical aspects of crystal structures and some crystal chemical applications. *J. Applied Cryst.*, 29, 305-306.
- BECKER, P.J. & P. COPPENS. 1974. Extinction within the limit of validity of the Darwin transfer equations. II. Refinement of extinction in spherical crystals of SrF_2 and LiF . *Acta Cryst.*, A30, 148-153.
- EBY, R.K. & F.C. HAWTHORNE. 1993. Structural relations in copper oxysalt minerals. I. Structural hierarchy. *Acta Cryst.*, B49, 28-56.
- GELATO, L.M. & E. PARTHÉ. 1987. STRUCTURE TIDY - a computer program to standardize crystal structure data. *J. Applied Cryst.*, 20, 139-143.
- HALL, S.R., H.D. FLACK & J.M. STEWART. 1992. Eds. Xtal3.2. Reference Manual. Universities of Western Australia, Geneva and Maryland.
- MANDARINO, J.A. 1981. The Gladstone-Dale relationship: part IV. The compatibility concept and its applications. *Can. Mineral.*, 19, 441-450.
- MARI, G. 1992. Les anciennes mines de cuivre du Dôme de Barrot (Alpes-Maritimes). Ed. Serre, Nice, 112 p.
- SARP, H., G. MARI, D. MARI & P. ROLLAND. 1994. Contribution à l'étude minéralogique des indices cuprifères de Roua (Daluis, Alpes-Maritimes, France). *Rivière Scientifique*, Nice, 47-56.
- SARP, H., G. MARI, D. MARI & P. ROLLAND. 1995. Sur la présence de vésigniéite dans les indices cuprifères de Roua (Daluis et Guillaumes, Alpes-Maritimes, France). *Rivière Scientifique*, Nice, 45-48.
- SARP, H., G. MARI, D. MARI & P. ROLLAND. 1996. Données nouvelles sur la minéralogie des indices cuprifères de Roua (Daluis et Guillaumes, Alpes-Maritimes, France). *Rivière Scientifique*, Nice, 27-36.
- SARP, H. & R. ČERNÝ. 1999. Gilmarite, $\text{Cu}_3(\text{AsO}_4)(\text{OH})_3$, a new mineral: its description and crystal structure. *Eur. J. Mineral.*, 11, 549-555.
- TOMAN, K. 1977. The symmetry and crystal structure of olivenite. *Acta Cryst.*, B33, 2628-2631.
- Vinchon, C. 1984. Sédimentogenèse et métallogénèse du Permien du Dôme de Barrot (Alpes-Maritimes). Thèse d'Etat. *Document B.R.G.M.*, no. 70, 445 p.