

Protogenetic coding bypass fixation : selectively robust to early earth alkalinity

Autor(en): **Turian, Gilbert**

Objektyp: **Article**

Zeitschrift: **Archives des sciences [2004-ff.]**

Band (Jahr): **59 (2006)**

Heft 1

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-738317>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Protogenetic coding bypass fixation selectively robust to early earth alkalinity

Gilbert TURIAN*

Manuscript received the 23rd march 2005, accepted the 28th october 2005

Abstract

Alkali-resistant phosphoramidic bonds could have provided an originally selective bypass to the alkalo-deficient acidophilic ribophosphoesterasic fixation of the prospective genetic code.

Keywords: coding, protogenetic, alkaline, fixation, bypass

Résumé

Fixation alcaline court-circuitante du codage protogénétique biogène. - L'alcalinité tellurique originelle pourrait avoir sélectionné les robustes liaisons phosphoramidiques pour la fixation du codage protogénétique des nucléobases, court-circuitant ainsi les liaisons ribophosphoestérasiques alcalo-déficientes du prospectif code génétique.

Mots-clés: codage, protogénétique, alcalin, fixation, court-circuit

Introduction

The precursors of the present-day genetic code, variously quoted as proto(s) (Kay 2000), first code (Wills & Bada 2000) or original code (De Duve 2002), to which we substitute "protocoding" in a more dynamic connotation, also consist of a list of correspondences between aminoacids specifically encoded by doublets or triplets of nucleobases either fixed stochastically by "frozen accident" (Crick 1968) or by preferential physical-stereochemical complementarity (Woese 1967; Mellersh 1993). We endorsed this last opinion in our anticodonic chart (Turian 1998), using the inversed base-paired letters of transfer-RNAs rather than those outstandingly deciphered by biochemists of the 60ties (Nirenberg & Mattei, 1961, in Kay 2000), namely C + C = G + G for glycine; C + G = G + C for alanine, etc.

What was then the environmental setting for the necessary validating fixation of such protogenetic coding? "The importance of being alkaline", as recently advertised by Russell (2003), confronted to both the original telluric alkalinity and to the necessary high pH robustness of a "pre-RNA world" coding fixation bond, sparked our evolutionary answer.

Primordial alkalinity of the early Earth rocked crust

Igneous rocks predominated in the primitive terrestrial lithosphere, among which silicate alkaline salts (Mg²⁺ - feldspars, Al³⁺ - kaolinite clays, etc.). The surface of these rocks might have been covered with microscopic pits with a probable high pH watery content, selectively favouring resistant intermolecular coding reactions.

In the vent systems of primitive oceans, mineral-catalyzed reduction of molecular nitrogen and oxides of nitrogen, suggested that hydrothermal environments and surrounding waters would have been the ammonia-rich and therefore alkaline. These conditions were most favourable to high pH serpentinization of olivine basaltic rocks of the early Earth crust (for references, see in Holm & Andersson 1998 and Zubay 2000) and, consecutively, to trigger the robustness of the protocoded processes.

* Prof. honoraire, 1212 Grand-Lancy, Geneva, Switzerland and Laboratoire de Bioenergetics, University of Geneva, 1254 Lullier, Switzerland.

Selectively bypassed ribodefective fixation of the protogenetic coding by alkaline-robust phosphoramidate bondings

Fixation of the proto-code on the ribophosphoesterasic of transfer RNA could not have occurred because of serious difficulties in the primordial synthesis of the sugar D-ribose (Shapiro 1988; Joyce & Orgel 1999), among which its lability in alkaline media (see Lahav 1999), even though possibly alleviated by boric acid present in interstellar space (Ricardo et al. 2004).

Consequently, among the predominant alkalinity of the original telluric environments, available protocoded nucleobases would have been deprived of ribose for their phosphorous fixation. This leads us to propose that the nucleotidic steps of pre-RNA synthesis would have been evolutively bypassed by phosphoramidic N → P bonds (Fig. 1), known to be selectively robust at high pHs (Lohrmann & Orgel 1973; Corbridge 1978), as confirmed for our nucleobase-phosphates (Turian et al. 1999; Turian & Rivara-Minten 2001). These bonds could thus have been the selective pressure factor for the chemical fixation of the protogenetic code.

Acidic transition to ribonucleotide fixation (RNAs)

The high pH dependance of protocoding would have been relevant in prebiotic to biotic evolution, following an environmental switch to acidic media such as the CO₂-enriched surface layers of the oceans (see

Russell 2003). Such acidification would have provided the necessary acidic opening of the N → P bondings by electric repulsion (Turian 2001). This would have permitted the median insertion of the hydroxylated precursors of ribose, glycolaldehyde and glyceraldehyde (see Eschenmoser's team work, 1997), produced by formose reaction in alkaline media, before an environmental acidic pH transition to bridge the opened bondings by the acidic pH-solid ribose.

Conclusion

Transition from N → P to N – C – P (C = ribose) bondings would have stabilized the proceedings from protogenetic to “modern” genetic code. Such biogenic filiation originating from geochemical alkalinity might have commanded protocoded “Protolife” before sparkling into the acidophilic RNA-DNA-genomics-proteomics Life systems.


Acknowledgements

We wish to thank Prof. Reto Strasser, Director of the Bioenergetics and Microbiology Laboratory, for his interest in our work, Dr Elisabeth Rivara-Minten for her expertised NMR analyses and Ms M. L. Manelli for typing the manuscript.

Fig. 1. Protogenetic coding alkaline-selected bypass fixation of pre-RNA.

(a) Phosphoramidate bondings (N à P) alkaline-selected (pH 8-10) bypasses of a phosphodiester linked ribopolynucleotide chain.

(b) Previous modellization (Turian, 2001) of protocoding nucleobase doublets, exemplified by: A (alanine) for C (cytosine) + G (guanine) riboselessly N – P fixed on the polyphosphate pre-RNA tape.



References

- **BRACK A.** (ed). 1998. *The Molecular Origins of Life*. Cambridge University Press, Cambridge, U.K.
- **CORBRIDGE DEC.** 1978. Phosphorus. An Outline of its Chemistry, Biochemistry and Technology. Elsevier Sci. Publ. Co.
- **CRICK FHC.** 1968. Origin of the genetic code. *J. Mol. Biol.* 38: 367-379.
- **DE DUVE C.** 2002. *Life Evolving. Molecules, Mind and Meaning*. Oxford University Press. 341 p.
- **ESCHENMOSE A.** 1997. Towards a chemical aetiology of nucleic acid structure. *Origins of Life*, 27: 535-553.
- **HOLM NG, ANDERSSON EM.** 1998. Hydrothermal systems, chapt. 4. In: Brack A (1998).
- **JOYCE GF, ORGEL LE.** 1999. Prospects for understanding the origin of the RNA world. *In: Gesteland RF, Cech TR, Atkins JF (eds.), The RNA World*, 2nd edition, Chapt 2: 49-77, Cold Spring Harbor, Labor. Press.
- **KAY LE.** 2000. *Who Wrote the Book of Life? A History of the Genetic Code*. Stanford University Press. Stanford California.
- **LAHAV N.** 1999. *Biogenesis: Theories of Life's Origins*. Oxford University Press.
- **LOHRMANN R, ORGEL LE.** 1973. Prebiotic activation processes. *Nature*, 244: 418-420.
- **MELLERSH AR.** 1993. Origins of Life Evol. *Biospheres*, 23: 261-270.
- **RICARDO A, CARRIGAN MA, OLCOTT AN, BENNER SA.** 2004. Borate minerals stabilize ribose. *Science*, 303: 196.
- **RUSSELL MJ.** 2003. The importance of being alkaline. *Nature*, 302: 580-581.
- **SHAPIRO R.** 1988. Prebiotic ribose synthesis: A critical analysis. *Origins of Life Evol. Biosphere*, 18: 71-85.
- **TURIAN G.** 1998. Polarity at onset of genetic coding. III. Prenucleic stereochemical interactions between amino acids and nucleobases doublets. *Archs Sci.* 51: 147-152.
- **TURIAN G.** 2001. Phosphoramidate-bonded polynucleobasephosphates as riboseless precursors of RNA? *In: Chela-Flores J et al. (eds.), First Steps in the Origin of Life in the Universe*, pp. 107-110. Kluwer Acad. Press, Dordrecht, The Netherlands.
- **TURIAN G, RIVARA-MINTEN E.** 2001. Prebiotic phosphoramidation of nucleobases by Mg²⁺-triggered decyclization of trimetaphosphate. *Archs. Sci.* 54: 233-238.
- **TURIAN G, RIVARA-MINTEN E, CATTANEO A.** 1999. Further ³¹P-NMR evidence of phosphoramidate bonding of nucleobases by Mg²⁺-enhanced nucleophilic attack on cyclic triphosphate. *Archs Sci.* 52: 209-216.
- **WILLS C, BADA J.** 2000. *The Spark of Life*. Perseus Publ., Cambridge Mass., 291 p.
- **WOESE CR.** 1967. *The Genetic Code: The Molecular Basis for Genetic Expression*, Harper & Row, New York.
- **ZUBAY G.** 2000. *Origins of Life on the Earth and in the Cosmos*, 2nd edition, Academic Press, 564 p.

