Zeitschrift:	Candollea : journal international de botanique systématique = international journal of systematic botany
Herausgeber:	Conservatoire et Jardin botaniques de la Ville de Genève
Band:	47 (1992)
Heft:	2
Artikel:	Taxonomic notes on the Hebe complex (Scrophulariaceae) in the New Zealand mountains
Autor:	Heads, Michael J.
DOI:	https://doi.org/10.5169/seals-879588

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 12.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Taxonomic notes on the Hebe complex (Scrophulariaceae) in the New Zealand mountains

MICHAEL J. HEADS

RÉSUMÉ

HEADS, M. J. (1992). Notes taxonomiques sur le complexe Hebe (Scrophulariacées) dans les montagnes de Nouvelle Zélande. *Candollea* 47: 583-595. En anglais, résumés français et anglais.

Descriptions et notes sont données pour tous les taxons de *Leonohebe* sect. *Buxifoliatae* et deux taxons d'*Hebe* sect. *Subdistichae*, Scrophulariacées alpines de Nouvelle Zélande.

ABSTRACT

HEADS, M. J. (1992). Taxonomic notes on the Hebe complex (Scrophulariaceae) in the New Zealand mountains. *Candollea* 47: 583-595. In English, French and English abstracts.

Descriptions and notes are given for all members of *Leonohebe* sect. *Buxifoliatae* and two members of *Hebe* sect. *Subdistichae*, alpine Scrophulariaceae of New Zealand.

Introduction

A synopsis of part of the *Hebe* complex of genera (Scrophulariaceae) has been published elsewhere (HEADS, 1987), with new names, brief diagnoses and typifications. The present paper provides fuller descriptions and notes on members of one section of *Leonohebe* and two species of *Hebe*. Distribution maps and discussion of generic delimitation are provided elsewhere (HEADS, 1992a, b).

Many important collections of the group have been made since it was last revised (MOORE, 1961). This study has benefited in particular from the extensive collections of Professor A. F. Mark and Mr. A. P. Druce.

Leonohebe Heads

Leonohebe sect. Buxifoliatae Moore ex Heads (1987: 10). Type species: Leonohebe odora (Hook. f.) Heads.

Prostrate to erect shrubs to 2 m high, final branches dark or bronzy green, sometimes flexed plagiotropically. Young leaf pairs coherent by the margins until nearly fully grown, or not coherent, bud sinus broad, shield-shaped or obtrullate-rhomboid, or elliptic with truncate or cordate base barely connate below, leaf-scar raised on pulvinate base. Irregularly lobed "reversion" foliage very rare. Inflorescence terminal or intercalary, of simple spikelets or branched spikes, each spike with

CODEN: CNDLAR ISSN: 0373-2967 47(2) 583 (1992) © CONSERVATOIRE ET JARDIN BOTANIQUES DE GENÈVE 1992 one to many pairs of flowers. Bracts opposite and connate throughout, at least lowermost bracts \pm leaf-like. Calyx with anterior lobes fused or not. Corolla white or lilac, 3-4-lobed with anterior lobe absent or narrowed or conduplicately folded enclosing style until anthesis, or corolla 5-lobed and \pm regular with two posterior lobes. Disc present. Capsule dorsally compressed or laterally compressed and \pm didymous. Chromosome number: $\times = 21$. Some populations of *L. odora* are tetraploid, *L. mooreae* is hexaploid and *L. masoniae* is an euploid: n = 59.

Distribution. - New Zealand: Auckland Is., Stewart Is., South Is., North Is.

Members of this section are important ecologically and dominate large areas of shrubland. The group is also phylogenetically important as certain species combine the bud structure of *Hebe sensu stricto* (Heads 1987) with the inflorescence of *Leonohebe*. Despite these features of interest the group has remained poorly understood. COCKAYNE & ALLAN (1927) despaired of the "many intergrading forms extremely difficult to classify", and possible additional taxa may remain lumped under taxa such as *L. odora* and *L. pauciramosa*.

CHEESEMAN's (1925) Veronica species 47-49 form the core of the section. Cheeseman, following HOOKER (1864), placed the group in Hebe s.s., next to what is now Hebe sect. Glaucae Heads. In contrast, MOORE (1967) placed the group in what can be regarded as another genus, Leonohebe, next to sect. Flagriformes. This latter treatment seems very perceptive and is followed here. Two species of L. sect. Buxifoliatae do share with Hebe s.s. the leaf margins coherent in the bud, but placing the section under Leonohebe is supported by the decussate inflorescence symmetry, and the bracts large and similar to vegetative leaves (cf. COCKAYNE & ALLAN, 1927). The flower and capsule of L. pauciflora and the strict, unbranched stems of L. pauciramosa also recall typical members of Leonohebe. Members of sect. Buxifoliatae are often confused with vegetatively similar members of Hebe sect. Subdistichae Heads, but the inflorescences are quite different.

Leonohebe pauciflora is very distinctive in this section. The four remaining species are closely related and neatly recombine the key inflorescence and stomata characters. Vegetative growth continues beyond the inflorescence in *L. pauciramosa, L. mooreae,* and *L. pauciflora* so that the inflorescence is intercalary (RICKETT, 1944), whereas in *L. odora, Hebe anomala* and *L. masoniae* all trace of terminal vegetative growth in the inflorescence is suppressed by the apex being totally "used up" in flowering. With respect to stomata, *Arctostaphylos* (Ericaceae) is another example of a difficult group which repaid a study of stomatal distribution (HOWELL, 1945). There is a notable "centre of taxonomic incongruence" at Arthur's Pass where an otherwise typical population of *L. odora* has adaxial stomata (see below).

Key to species of Leonohebe sect. Buxifoliatae

Note: Lead "B" below will incorrectly key Caswell Sound and Denniston Plateau populations of *L. mooreae*, and Arthur's Pass populations of *L. odora*.

Α	Fruit compressed laterally, corolla usually with 5 lobes 1. L. pauciflora	
Α	Fruit compressed dorsiventrally, corolla usually with (3)-4 lobes	В
В	Stomata on adaxial (upper) leaf surface and abaxial surface	С
В	Stomata on abaxial leaf surface only	D
С	Inflorescence intercalary ("lateral"), including terminal vegetative bud 4. L. pauciramosa	
С	Inflorescence terminal, not including vegetative bud 5. L. masoniae	
D	Inflorescence intercalary ("lateral"), including terminal vegetative bud 3. L. mooreae	
D	Inflorescence terminal, not including vegetative bud	E

- E Leaves usually < 3.5 mm wide, cuneate at base, anterior calyx lobes sometimes fused, anterior corolla lobe sometimes absent 2a. Veronica anomala
- E Leaves usually > 3.5 mm wide, anterior calyx lobes always free, anterior corolla lobe always present 2. L. odora

1. Leonohebe pauciflora (Simpson & Thomson) Heads (1987: 10). *Hebe pauciflora* Simpson (1943: 166). Type: G. Simpson March 1942 (CHR).

Closely branched subshrub to 20 cm high, branches prostrate, rooting, ascending distally. Leaves $5-7 \times 3.5-5$ mm, rhomboid, coriaceous, pairs coherent in bud by dense marginal pubescence, bud with open sinus, leaf margins thick, cartilaginous, entire, midrib forming rounded keel. Spikelets lateral, developing in pairs just behind vegetative apices, each spikelet with one or two pairs of sessile flowers, rachis apex aborting. Calyx 4-lobed, with dense, crinkled pubescence. Corolla white, usually pentamerous with "split" posterior corolla lobe. Ovary laterally compressed, seated on ringed disc. Capsule 4×3 mm, obcordate and didymous, dehiscence deeply loculicidal as well as septicidal.

Distribution. — Southwest South Island.

Habitat. — Rocky sites, in crevices and on ledges.

This species is quite distinct from all other members of the section by its foliage, perianth and fruit, although in the wild the plant may be mistaken for a depressed state of *L. petriei* var. *murrellii* as SIMPSON & THOMSON (1943) noticed, and MOORE (1961) recognised a "superficial resemblance" to *L. densifolia*. In fact, the inflorescences, flowers and fruits of *L. pauciflora* and *L. densifolia* all show similarities, and *L. pauciflora* may be an important link among *Leonohebe* sect. *Buxifoliatae*, sect. *Densifoliae* Heads, and the puzzling *L. petriei*.

Leonohebe odora (Hook. f.) Heads (1987: 10). Veronica odora Hook. f. (1844: 62). Hebe odora (Hook. f.) Ckne. (1929: 472). Type: J. D. Hooker 1460 (K) "Lord Auckland's Group" [not seen].

Multi-leadered woody plants, either with prostrate and rooting runners and orthotropic shoots to 15 cm high, or forming erect hemispherical shrubs to 1.5 m high. Branches often plagiotropically arched distally with phyllotaxis tending to distichous through twisting of the petioles. Internodes 1.5-5 mm long. Leaf pairs connate at the base by a line of hairs which extends along the lower petiole margins, meets the convergent abscission zones and continues down the stem as bifarious strips of pubescence. Leaves coherent in bud. Lamina 7-16 \times 3-7 mm, adaxial surface dark green and shining, without stomata, abaxial surface paler, with stomata. Margins, midrib and young stem yellowish green, margin entire, very rarely lobed ("reversion foliage"), bevelled, either smooth or thickly cartilaginous and minutely crenulate. Lamina glabrous, elliptic to slightly ovate or obovate, \pm acute, concave, midrib often not evident above but sometimes impressed, \pm keeled below. 2-3 veins sometimes visible parallel with the midrib, lamina truncate at the base, leaf pairs forming a broad shield-shaped sinus in the bud. Petiole 1-3 mm long, spreading to a broad, \pm pulvinate base. Inflorescence composed of terminal and usually some subjacent lateral spikes, each of at least (3)-4 pairs of sessile flowers, together forming an oblong, triangular or corymbose head 1-6 cm long, not including a vegetative bud, new growth coming from > 2 cm below inflorescence. Flowers and bracts strictly opposite throughout, bracts \pm connate at their base with a rim of hairs, rachis bifariously pubescent. Bracts broadly triangular, ± acute, 4 mm long, margin often minutely ciliolate below, lowest bracts leaf-like in texture. Calyx lobes ovate, minutely ciliolate, anterior pair fused to ca. 1/4, but overlapping to 3/4 of their length, posterior pair free to base, not overlapping. Corolla white or pink, tube = or > calyx, 6 mm long, lobes four, $2.5-6 \times 2-3$ mm, anterior lobe usually narrower, folded conduplicately and hooded, enclosing the style until anthesis. Disc lobed, to 1/4 length of ovary. Ovary glabrous, 1.0-1.5 mm long. Capsule dorsiventrally compressed, glabrous, ovate, $5(-6.5) \times 2-3$ mm, dehiscence septicidal with a little loculicidal dehiscence at apex. Chromosome number n = 21 or 42.

Distribution. — Auckland Is., Stewart Is., South Is., North Is.

Habitat. — Poorly drained sites — bogs, flushes, pakihi swamp, "swampy vegetation in stream flowing through mountain beech forest" (Thomas R.) or well drained sites — moraine, stable scree, rocky outcrops. Very rarely under forest canopy, but abundant on forest margins, in shrubland and in grassland. Found on soils from varied parent material, for example on soils derived from ultramafic rocks at Red Hill (Wairau). Sea-level (Auckland Is. and Stewart Is.) to 1440 m.

Often one main leader develops as a relay axis beyond the zone of inflorescence (Koriba's architectural model of HALLÉ & al., 1978), although these axes, and even short trunk axes may show some plagiotropy distally.

Many adaxial stomata have been seen on one piece from each of Moa Park, Craigieburn, Mt. Ida and Forbes Mts., and on all collections seen from Arthur's Pass. This last locality is well-known as a centre of endemism, as a northern boundary *(Hectorella-Hectorellaceae)* and now as a centre of taxonomic "incongruence".

Hebe odora var. prostrata was listed but not formally described by COCKAYNE (1909: 44). B. Patrick (pers. comm., 1987) has identified this with a distinct, prostrate form of Stewart Is. and Longwood Ra. (southern South Is.), which sometimes grows together with typical *L. odora*. HAIR (1967) records n = 42 for plants identified as *H. odora* var. prostrata, but also for several other South Is. populations of "Hebe" odora. Further field study is required. Fiordland specimens of *L. odora* have large, acute leaves to 16×7 mm, with leaf bases conspicuously swollen and with margins minutely crenulate (but less thickened than those of *L. mooreae*). Crenulate leaf margins also occur in some central Otago plants, and in all Stewart Is. and Auckland Is. plants. Extreme leaf dimensions are also recorded at, for example, Eyre Mountains (Given, Nov. 1971 (CHR): 15 \times 9 mm). Lobed "reversion" foliage is very rarely present (Johnson, 21.1.1974 (CHR) Mavora Hills).

2a. Veronica anomala J. F. Armstrong (1872: 271). Hebe anomala (J. F. Armstr.) Ckne. (1929: 468).
 Type: Armstrong (CM) "headwaters of the River Rakaia" [not seen].

MOORE (1961) described the cuneate-based leaves and fused anterior calyx lobes of this form and Armstrong described the corolla as 3-lobed. This character combination is very distinctive in a widely cultivated form which is probably near V. anomala. These plants are closely related to L. odora through their terminal inflorescence, for example. On the Rock and Pillar Ra. shrubs of L. odora are generally the typical form with leaves $8 \times 4 - 9 \times 5$ mm. However, Gray, 11.3.1972, (OTA) Rock and Pillar Ra., has a narrower leaf (9×3.5 mm) with a more well-defined mid-rib, and approaches the cultivated form referred to above. The leaf base is also less truncate in Gray's collection and the leaf pairs split very early in the bud. The status and distribution of V. anomala remain to be ascertained. Further to the type and the Gray collection, MOORE (1961) cites Armstrong material from Rangitata, and the localities: Rock and Pillar, Rangitata Valley, upper Rakaia suggest a standard northeast Otago-mid-Canterbury distribution pattern.

Leonohebe mooreae Heads (1987: 10) var. mooreae. Veronica menziesii Benth. sensu Simpson & Thomson (1942: 21). Hebe odora "menziesii form" Wardle (1975). Hebe sp. "i" Eagle (1982). Type: P. Wardle, 16.12.1978 (CHR) Douglas Range, south Westland.

Prostrate to erect shrub to 2 m high, branching often distinctly rhythmic, internodes to 8 mm long. Leaf base swollen, drying to a saddle-shaped inflation decurrent to the node below. Leaves sometimes dorsiventrally arranged, $16(-30) \times 4-8$ mm, narrowly ovate-elliptic, acute, all stomata abaxial but leaf \pm the same colour above as below, very coriaceous, margins bevelled, minutely crenulate. Midrib usually impressed above, conspicuous below, no other veins usually visible. Petiole 2 mm long, glabrous except where narrow bifarious grooves in the stem lined with pubescence intersect the petiole connature. Inflorescence intercalary, including a terminal vegetative bud, comprising one or two pairs of lateral spikes, each of these simple, very rarely tripartitely branched, to 4 cm long, and with at least 4 pairs of flowers. Rachis often pubescent, bracts strictly opposite throughout, connate, broad-triangular, ciliolate at least on lower margins at the connature, margins

membranous, often tinged purple. Flowers sessile, calyx 3-4 mm long, usually ciliolate at least apically, anterior lobes fused to 1/3 of their length, posterior two free. Corolla tube usually \pm equal to calyx lobes, lobes \pm clawed, 2.5-4 \times 5-6 mm. Disc lobed, to 1/3 length of ovary. Ovary glabrous, ovate, to 1.5 mm long. Fruit dorsally compressed, 6 \times 3 mm. Chromosome number n = 63 (Fig. 1).

Distribution. — Western South Island.

Habitat. — Poorly drained sites — bogs, *pakihi* swamp, lakeshores. Well-drained sites — rocky slopes and ledges. Occasionally in *Nothofagus* forest. most often in shrubland, where it may be structurally important, and also in grassland. On soils derived from a wide range of parent material, often found on soils on ultramafic rocks in tussock shrubland and dwarf *Nothofagus* forest, 5 m (Cascade R. mouth) – 1200 m.

Plants with adaxial stomata are known only from Caswell Sound, Fiordland, and Denniston Plateau, Nelson. The crenulate leaf margins recall *Coprosma serrulata* (Rubiaceae), also found in shrublands of southwest South Island. *Johnson*, 31.3.1977 (CHR), Cascade River mouth, Lake Jumbuck lakeshore scrub and forest, 5 m altitude, is a luxuriant form growing at Jackson Bay, a major biogeographic centre (HEADS, 1989). The collection has internodes up to 6 mm long, petioles up to 3 mm long, and leaves to 24×5.5 mm and had been identified, not surprisingly, as the unrelated *Hebe elliptica*. *Druce*, Jan. 1969 (CHR) Gouland Downs 1900', has corollas 15 mm long with the limb 15 mm broad, leaves acute, 22×4 mm. Sites of probable leader abortion are evident. This is another luxuriant form, again found at a locality of general biogeographic interest (HEADS, 1989).

3a. Leonohebe mooreae var. **telmata** Heads (1987: 10). *Hebe odora* "bog form" Wardle (1975). **Type:** *P. Wardle*, 16.12.1978 (CHR), Douglas Range, south Westland.

Shrub up to 60 cm tall. Lamina 10×4 mm. Spikelets smaller than those of var. *mooreae*, comprising 2 or 3 pairs of flowers only (Fig. 2).

Distribution. — South Is., Westland National Park. Douglas Range (type); between Jalf Glacier and head of Waikukupa River (*Wardle*, 26.2.1974, CHR); tops east of Mt. Fox (*Wardle & Fryer*, 27.2.1967, CHR).

Habitat. — Poorly drained low-alpine grassland and shrubland, 1160-1400 m.

Westland plants of var. *mooreae* have inflorescences of at least 5 pairs. The two forms of *L. mooreae* are probably subspecies or species and both occur together on the Douglas Range. Field study is required, as is a study of *L. mooreae* in Nelson, where some poorly known Karamea and Gouland Downs populations may belong with var. *telmata*.

4. Leonohebe pauciramosa (Ckne. & Allan) Heads (1987: 10). Hebe buxifolia var. pauciramosa Ckne. & Allan (1926: 27). Hebe pauciramosa (Ckne. & Allan) Moore (1961: 925). Type: L. Cockayne 8129 (WELT), Lake Harris.

Shrub of open habit to 50 cm high, stems seldom branched, the largest generally prostrate and rooting proximally, and orthotropic distally. Stem pubescence bifarious, in narrow grooves. Internodes to 5 mm long. Leaves coriaceous, dark bronze-green, splitting apart very early in bud, 6-8(-10) \times 4-6(-7) mm, widely obovate, obtuse, \pm mucronate, concave. Lamina abruptly truncate or cordate at base, bud with broad, obtrullate sinus. Lamina glabrous except for minute, prickle-like cilia on upper part of the margin. Midrib obvious on lower surface only, usually not evident above, keel of leaf often flattened abaxially just below apex. Petiole glabrous, 2 mm long, spreading at base. Inflorescences all intercalary, stem apex remaining vegetative, zone of inflorescence to 2 cm long. Rachis \pm pubescent, bracts all strictly opposite, connate, acute to obtuse, 2-3 mm long, with minute cilia. Calyx lobes 3-4 mm long, anterior two fused 50-90% of their length, ciliolate above. Corolla tube to 6 mm long, lobes ovate, sometimes clawed, to 5 \times 3 mm. Style held in the conduplicately folded anterior corolla lobe prior to anthesis. Disc lobed, 1/4-1/3 length of ovary. Capsule dorsally compressed, ovate-oblong, 5 \times 2 mm, often with a small amount of loculicidal dehiscence. Chromosome number n = 21.



Fig. 1. — Leonohebe mooreae var. mooreae. A, habit; B, flower; C, disc; D, distal portion of infructescence (A, B, C from type, D from I. M. Ritchie (CHR) Te Waewae).



Fig. 2. — Leonohebe mooreae var. telmata. A, habit; B, flower (posterior petal uppermost); C, oblique view of flower (posterior petal at bottom left) (A from type, B and C from Wardle & Fryer, 27.2.1967, CHR).

CANDOLLEA 47, 1992

Distribution. — Auckland Is., South Is., North Is. (Mt Taranaki only).

Habitat. — Poorly drained sites — swampy areas, seepages, flushes, streambanks. Welldrained sites — moraine, gravel in river and creek beds. Most often collected in grassland, and also known from forest margin, shrubland and herbfield, 450-1650 m.

Certain central Otago collections of this species, notably *Chaffey*, 30.12.1975 (CHR) Blue Lake, and *Mark*, 8.2.1985 (OTA) upper West Waikaia V., both from the Garvie Mts. (and cf. *Mark*, 22.11.1970 (OTA), Carrick Ra.) are more branched than the typical form. *Mark*, 8.2.1985 (previously identified as *Hebe buchananii*, which it resembles) is annotated "common on rock outcrops" which is also atypical for *L. pauciramosa* s.s.

5. Leonohebe masoniae Heads (1987: 10) var. masoniae. Hebe pauciramosa var. masonae Moore (1961: 926, 972). Type: R. Mason, 23.2.1946 (CHR), Cobb Valley.

Shrub to at least 20 cm high, internodes to 5 mm long, stem pubescence bifarious in grooves, petiole connature ciliate centrally, leaves separating early in bud. Leaves elliptic to slightly obovate or ovate, $8-16 \times 4-6$ mm, keel conspicuous, continued abaxially into short mucro. Leaf margin rounded, sometimes \pm bevelled near apex. Young leaves and leaves below inflorescence glabrous or with comb-like rim of caducous, erect, white cilia, stomata conspicuous on upper and lower leaf surfaces. Petiole to 3 mm long. Inflorescence usually terminal, simple, 1-4 cm long, sometimes branched with lateral spikes. Bracts opposite and connate throughout, conspicuous, leaf-like, to 7×4 mm, glabrous or with comb-like rim of marginal cilia. Calyx lobes to 5 mm long, anterior lobes fused to 1/2 of their length, densely ciliolate, corolla tube to 7 mm long, lobes white or mauve, rounded, clawed, to 6×5 mm. Disc to 1/4 length of ovary. Ovary with loculicidal ridge and slight septicidal groove, stigma bilobed. Capsule 6×3 mm. Chromosome number n = 59 (Fig. 3).

Distribution. — Western South Island.

Habitat. — Forest margin, shrubland, tussockland, 660-1350 m.

This variety is more lax than var. *rotundata*, and also differs through its longer and narrower leaves and the bifarious strips of pubescence which often line the stem grooves.

Habit and leaf size and shape in *L. masoniae* can be very similar to those of *L. mooreae*, but the adaxial stomata and rounded leaf margin of the former allow identification of even sterile material. *Wilson*, 9.12.1971 (CHR) is a puzzling collection from Worryline Stream, Ben Ohau Ra., Mt. Cook region. It has the compound inflorescence of *L. odora* but in most other respects resembles *L. masoniae*, which seems to be as variable as its more widespread relatives. Collections of this species have previously been identified as *Hebe odora*, *Hebe odora* "robust form" (i.e. *L. mooreae*), *H. pauciramosa* and *H. pauciramosa* var. *masoniae*. Sterile specimens from the Cobb Valley (*Druce*, Feb. 1977, CHR) which may belong here have several branches with lobed leaves, a feature also recorded in *L. odora*.

5a. Leonohebe masoniae Heads var. rotundata Heads (1987: 11). Type: A. P. Druce, Nov. 1980 (CHR), "Mineral Belt" north of Cobb Valley.

Erect shrub to 1 m high, leaves crowded at ends of branches and appressed, internodes short, comprised of swollen leaf-bases 1-2 mm long, stem pubescence sparse, bifarious, not in grooves but on flattened central portion of stem, older stems cicatricose. Leaves small and rounded, 5-7 \times 4-6 mm, concave. Margin smooth and rounded, neither crenulate nor bevelled. Midrib obscure above, keeled below, often extending to apex. Stomata conspicuous above and below. Petiole glabrous, 1-2 mm long. Young leaves (including those below the inflorescence) with a comb-like rim of erect, white, caducous cilia. Inflorescence a terminal, simple spike 1-2 cm long. Rachis with short, scattered, erect hairs. Bracts large and conspicuous, hiding the calyx, opposite and connate throughout. Lowest bracts \pm adjacent leaves but larger and with more membranous margins, to 9 \times 6 mm, margin usually ciliolate throughout. Calyx lobes to 6 \times 2 mm, obovate, ciliolate at apex,

two anterior lobes fused to 1/3 of their length. Corolla white or mauve, tube funnel-shaped, to 7 mm long, lobes (3)-4(-6), rounded, clawed, to 6×5 mm, anterior lobe folded conduplicately around style before anthesis. Disc 1/4 length of ovary. Capsule broadly obovate, dorsally compressed, 7×5 mm, slightly exceeding calyx (Fig. 3).

Distribution. — Northwest Nelson, South Is. Mt. Arthur, Mt. Peel, Lake Sylvester and near Cobb Dam, giving a distribution with a radius of some 8 kilometers. Collections seen: *Talbot*, Jan. 1962 (CHR) Cobb R.; *R. Melville*, 16.1.1962 (CHR) Cobb R.; *R. Melville*, 16.1.1962 (CHR) Mt. Cobb; *M. J. A. Simpson*, 16.1.1978 (CHR), Mt. Arthur; *M. A. & I. M. Ritchie*, 7.1.1969 (CHR) upper Takaka R.; *A. Campbell & C. McDonald*, 9.1.1978 (CHR) Mt. Peel; *J. Wells*, 5.11.1967 (OTA), Mt. Arthur.

Habitat. — Seepage areas in shrubland and tussockland, 1050-1500 m.

The variety name refers to the pulvinate "leaf-bases" and rounded leaves. As in related species, the leaves actually abscise above the abscission zone, so that a short portion of petiole remains attached to the "leaf-base".

Hebe Commerson ex Juss.

Hebe sect. Subdistichae Moore ex Heads (1987: 11).

H. dilatata Simpson & Thomson (1943: 164). Type: G. Simpson & J. S. Thomson (CHR), Blue Lake, Garvie Mts., 1370 m.

Prostrate, much-branched shrub, stems occasionally rooting adventitiously; stems with bifarious grooves including sparse pubescence, bud with narrow sinus, internodes 5(-10) mm long. Leaves \pm acute, midrib conspicuous and ridged below, and evident above, margin thickened especially apically and \pm bevelled. Inflorescence to 18 mm long, unbranched, bearing ca. 10 flowers, peduncle to 10 mm long, ca. 5 sterile bracts conspicuous at the base of each inflorescence. Lowermost bracts opposite. Calyx lobes rounded, ciliate, anterior lobes fused to 50-90% of their length, corolla tube < or = calyx, ovary glabrous. Capsule obovate, acute, 4 mm long.

Distribution. — Garvie Mts., South Island.

Habitat. — Alpine debris slopes, 1370 m.

The affinities of this species, placed by MOORE (1961) in "incertae sedis", lie with *H. cockay-neana*, rather than with *H. biggarrii* as Moore suggested. Apart from the holotype and two isotypes I have seen only two Simpson collections, both from the Garvie Mts. and matching the type.

Hebe crawii Heads (1987: 11). Type: A. F. Mark, 2.2.1971 (OTA), Excelsior Peak, Takitimu Mts., 4800'.

Decumbent, diffusely branched shrub to 40 cm tall, branchlets with bifarious pubescence, basal connature of leaves bearing rim of hairs, bud sinus narrow. Leaves $17-25 \times 9-12$ mm, broad obovate-spathulate, gradually narrowed to base, glaucous above and below, sometimes apiculate, margin cartilaginous and smooth. Inflorescence lateral, regularly compound (tripartite), peduncles and pedicels generally with some slight pubescence, pedicel to 2.5 mm, bracts to 3 mm long. Calyx lobes 4-5, 1.5 mm long, rounded-acute, corolla white, tube 1.0 mm long, lobes 2 mm long, broad, rounded. Disc short, \pm 4-lobed. Ovary glabrous, capsule 4 mm long, glabrous, acuminate (Fig. 4).

Distribution. — (Fig. 5): South Island.

Habitat. — Low-alpine shrubland, fellfield, boulderfield, partly stable talus slopes, 1200-1650 m.



Fig. 3. — Leonohebe masoniae var. rotundata.
A, habit; B, C, D, flowers (A from type, B and C from Campbell & McDonald, 9.1.1978, CHR).
L. masoniae var. masoniae: E, habit (Druce, Nov. 1976, CHR, Hope Range).



Fig. 4. — Hebe crawii.
 A, habit; B, leaf bud with narrow sinus; C, D, young fruits with persistent calyces; E, flowers at anthesis (anterior views on left, posterior views on right) (A-D from type, E from Dickinson, 13.1.1986, OTA, Gem Lake, Umbrella Mts.).



Fig. 5. — Hebe cockayneana: dotted stippled line; H. crawii: continuous line; H. dilatata: stippled circle; H. canterburiensis: hatched line.

Collections seen. — A. Wall, Feb. 1920 (CHR), Takitimu Mts.; P. Wardle, 31.1.1962 (CHR), Head of Aparima R., Takitimu Ra.; C. Meurk, 17.12.1972 (OTA), Corner Peaks, Takitimu Ra.; A. Mark, 7.2.1985 (OTA), Upper West Waikaia Valley, Garvie Mts.; K. Dickinson, 14.12.1985 (OTA), Gem Lake Cirque; K. Dickinson, 13.1.1986 (OTA), Gem Lake Cirque, Umbrella Mts.; K. Dickinson (OTA), Mt. Tennyson, southern Garvie Mts.; N. Simpson (OTA), Irthing Stream, SE Eyre Mts.; B. Rance (OTA), Robert Creek, NE Eyre Mts.; B. Rance, 7.1.1987 (OTA), Way Spur, Eyre Mts.

The large, spathulate and glaucous leaves are very distinctive and recall *Hebe cockayneana* (which has smaller, elliptic leaves), *H. pinguifolia* and *H. topiaria*. The tripartite inflorescence recalls northern members of *Hebe* sect. *Subdistichae*. The species has also been confused with *H. dilatata*.

ACKNOWLEDGMENTS

This study was supported by a fellowship from the E. L. Hellaby Indigenous Grasslands Research Trust and carried out in the Botany Department, University of Otago. Drs. P. J. Garnock-Jones, P. Weston and J. B. Wilson, and the late Dr. L. B. Moore kindly read versions of this paper and made helpful suggestions.

REFERENCES

ARMSTRONG, J. F. (1872). On some new species of New Zealand plants. Trans. New Zealand Inst. 4: 290-291.

CHEESEMAN, T. F. (1925). Manual of the New Zealand Flora. 2nd Ed. Government Printer, Wellington.

COCKAYNE, L. (1909). Report on a botanical survey of Stewart Island. Wellington.

COCKAYNE, L. (1929). New combinations in the genus Hebe. Trans. New Zealand Inst. 60: 465-472.

- COCKAYNE, L. & H. H. ALLAN (1926). Notes on New Zealand floristic botany, including descriptions of new species (No. 4). *Trans. New Zealand Inst.* 56: 21-33.
- COCKAYNE, L. & H. H. ALLAN (1927). The present taxonomic status of the New Zealand species of Hebe. Trans. New Zealand Inst. 57: 11-47.

EAGLE, A. (1982). Eagle's Trees and Shrubs of New Zealand. Second Series. Collins, Auckland.

- HAIR, J. B. (1967). Contributions to a chromosome atlas of the New Zealand flora 10. Hebe (Scrophulariaceae). New Zealand J. Bot. 5: 322-352.
- HALLÉ, F., R. A. A. OLDEMAN & P. B. TOMLINSON (1978). Tropical Trees and Forests: an architectural analysis. Springer-Verlag, Berlin.
- HEADS, M. J. (1987). New names in New Zealand Scrophulariaceae. Otago Bot. Soc. Newsletter 5: 4-11.
- HEADS, M. J. (1989). Integrating earth and life sciences in New Zealand natural history: the parallel arcs model. New Zealand J. Zoology 16: 549-485.
- HEADS, M. J. (1992a). Morphology, architecture and taxonomy in the Hebe complex (Scrophulariaceae). Submitted.
- HEADS, M. J. (1992b). Biogeography and evolution in the Hebe complex (Scrophulariaceae). Submitted.
- HOOKER, J. D. (1844). Flora Antarctica. Vol. 1, Part 1. Reeve, London.

HOOKER, J. D. (1864-1867). Handbook of the New Zealand Flora. Reeve, London.

HOWELL, J. T. (1945). Concerning stomata on leaves in Arctostaphylos. Wasmann Collector 6: 57-65.

- MOORE, L. B. (1961). Hebe (except whipcord species) and Ourisia. In: H. H. ALLAN (Ed.), Flora of New Zealand.: 1. Government Printer, Wellington.
- MOORE, L. B. (1967). How to look at Hebe. Tuatara 15: 10-15.
- RICKETT, H. W. (1944). The classification of inflorescences. Bot. Rev. 10: 187-231.
- SIMPSON, G. & J. S. THOMSON (1942). Notes on some New Zealand plants and descriptions of new species (No. 2). Trans. Roy. Soc. New Zealand 72: 21-40.
- SIMPSON, G. & J. S. THOMSON (1943). Notes on some New Zealand plants and descriptions of new species. *Trans. Roy* Soc. New Zealand 73: 155-171.
- WARDLE, P. (1975). Vascular plants of Westland National Park (New Zealand) and neighbouring lowland and coastal areas. New Zealand J. Bot. 13: 497-545.

Address of the author: Department of Biological Sciences, University of Zimbabwe, Harare, Zimbabwe.