

# The Puna vegetation in the valley of Rio Cazaderos, Catamarca Province, Argentina

Autor(en): **Martinez Carretero, Eduardo**

Objektyp: **Article**

Zeitschrift: **Candollea : journal international de botanique systématique = international journal of systematic botany**

Band (Jahr): **52 (1997)**

Heft 2

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-879451>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# The Puna vegetation in the valley of Rio Cazaderos, Catamarca Province, Argentina

EDUARDO MARTINEZ CARRETERO

## ABSTRACT

MARTINEZ CARRETERO, E. (1997). The Puna vegetation in the valley of Rio Cazaderos, Catamarca Province, Argentina. *Candollea* 52: 497-508. In English, English and Spanish abstracts.

The Puna vegetation in the valley of Río Cazaderos, Catamarca, Argentina, between 3600-4300 m s.m., was studied from the phytosociological point of view. Fifty-five floristic relevés following the Braun-Blanquet method were made and arranged in a table. The floristical relevés include xeric and wetland vegetation. Two altitudinal-vegetational zones are proposed: a lower elevation shrub zone, between 3600-3800 m s.m., dominated by the *Acantholippia punensis* community, with *A. punensis*, *Chiquiraga atacamensis*, *Lycium chañar*, etc., and a higher elevation zone, between 3800-4100 m s.m., characterized by the *Stipa chrysophylla* community, with *S. chrysophylla*, *Adesmia subterranea*, *Fabiana bryoides*, etc. The latter community shows a large spatial distribution from the southern parts of Bolivia to the provinces of Catamarca and, possibly, La Rioja, in Argentina. The maximum floristic diversity occurs in wetlands, where vegetational belts appear, composed of different plant communities, following a soil moisture gradient. The principal plant communities are: the *Carex gayana* community in deep, freshwater saturated soils; the *Oxychloe andina* community, above 4100 m s.m., near open and moving water; and the *Ranunculus mandoniana* community in wetlands with very shallow soils, saturated with fast-moving water. Other communities are present but limited in their distribution to specific edaphic conditions, such as the psammophilous *Lampaya hieronymi* community (3800-3900 m s.m.) in sandy places (sand accumulated by the wind); or the *Distichlis scoparia* community on saline soils where the water table is over 2 m below the surface. From our data we propose the hypothesis that plant dynamism in the wetlands is currently related to the effects of diminishing rainfalls on the hydric regime. We also suggest existence of new taxonomic units for Argentina, some of which were already suspected. From the *Calamagrostieta vicunarium* Rivas Martinez & Tovar, 1982, the following communities are present in the area: *Lobivio ferocis-Fabianion densae* Ruthsatz, 1993; *Urbanio pappigeriae-Stipion frigidae* G. Navarro, 1993; *Puccinellio oresigenae-Oxychloetum andinae* G. Navarro, 1993.

## RESUMEN

MARTINEZ CARRETERO, E. (1997). La vegetación puneña en el valle del Río Cazaderos, Catamarca, Argentina. *Candollea* 52: 497-508. En inglés, resúmenes en inglés y español.

Se estudia fitosociológicamente la vegetación puneña del valle del Río Cazaderos, Catamarca, comprendida entre los 3600-4300 m s.m. Cincuenta y cinco relevamientos florísticos fueron efectuados según el método de Braun-Blanquet y ordenados en una tabla de acuerdo al gradiente de xericidad. Dos zonas altitudinales de vegetación se proponen: una zona inferior, arbustiva, entre 3600-3800 m s.m., dominada por la comunidad de *Acantholippia punensis*, con *A. punensis*, *Chiquiraga atacamensis*, *Lycium chañar*, etc., y otra zona a mayor elevación, entre 3800-4100 m s.m., caracterizada por la comunidad de *Stipa chrysophylla*, con *S. chrysophylla*, *Adesmia subterranea*, *Fabiana bryoides*, etc. Esta última comunidad muestra una extensa distribución espacial desde el sur de Bolivia hasta la provincia de Catamarca y, posiblemente, de La Rioja en Argentina. La máxima diversidad florística ocurre en las vegas, donde los cinturones de vegetación, constituyendo diferentes comunidades vegetales, siguen un gradiente de humedad del suelo. Las principales comunidades vegetales son: la comunidad de *Carex gayana* en suelos profundos saturados de agua, la comunidad de *Oxychloe andina*, sobre los 4100 m s.m., en contacto con agua libre en movimiento, y la comunidad de *Ranunculus mandoniana* en vegas con suelos muy superficiales, saturados con agua en rápido movimiento. Otras comunidades están presentes pero limitadas en su distribución por condiciones edáficas específicas, tales como la comunidad de *Lampaya medi-*

*cinalis* (3800-3900 m s.m.) en lugares arenosos (arena acumulada por el viento), o la comunidad de *Dystichlis scoparia* en suelos salinos donde el agua freática supera los 2 m de profundidad. A partir de nuestros datos se propone una hipótesis del dinamismo vegetal en vegas, estrechamente relacionado con los efectos de la disminución de las lluvias sobre el régimen hídrico. Se sugiere además la existencia de nuevas unidades sintaxonómicas para la Argentina, algunas de las cuales se sospechaba su presencia. De los *Calamagrostieta vicunarum* Rivas Martínez y Tovar, 1982, las siguientes comunidades están presentes en el área: *Lobivio ferocis-Fabianion densae* Ruthsatz, 1993; *Urbanio pappigeriae-Stipion frigidae* G. Navarro, 1993; *Puccinellio oresigenae-Oxychloetum andinae* G. Navarro, 1993.

**KEY-WORDS:** Puna – Vegetation – Dynamismus.

### Introduction

There are only a few studies on the Puna vegetation in the Province of Catamarca. VERVOORST (1951) suggested the following zonation for the Puna vegetation in the area of Salinas de Laguna Verde, south of the Río Cazadero valley: a shrubby steppe between 3500-3800 m s.m. and a grassy steppe between 3800-4300 m s.m. For CABRERA (1957, 1968, 1976) and CABRERA & WILLING (1973), the Puna extends to the south through the mountains of the provinces of Catamarca and La Rioja between 3000-3700 m s.m. HUNZIKER's work (1952) in northern La Rioja describes the distribution of the Puna area on the border of both provinces. Following MARTINEZ CARRETERO (1995) this area would be included in the so called Distrito Jujeño (Jujeño district). From the geological point of view the Puna boundary lies between Antofagasta (Chile) and the Sierra de Quilmes, Las Cuevas and the Cordillera de San Buenaventura, in the Catamarca Province (TURNER & MENDEZ, 1979; ALLMENDIGER, 1986). It is stratigraphically characterized by Cenozoic volcanic units (TURNER & MENDEZ, 1979). The oldest units are mainly silicious volcanics and the youngest are andesitic strata overlying the Quaternary deposits constituted by tuffa and alluvial deposits. The Holocene deposits are mostly composed of sand and clay, generally reworked by floods and presently covered by sand dunes. Using an ESE-NW transect Fiambalá-C° Nacimiento, the following formations could be found: the Formación La Planchada to the west with Palaeozoic volcanic, andesitic rocks, followed by crystalline granitic Precambrian rocks, and finally Tertiary sediments from the upper Pliocene in the east (ALBA, 1979).

The entire area is markedly arid: Fiambalá, 1400 m s.m., receives 137 mm mean annual precipitation and Antofagasta de la Sierra (3400 m s.m.) 105 mm mean annual precipitation (Table 1). Aridity increases to the west, with only 62 mm of rain per year and a mean annual temperature of 6.8°C in Ollagüe (Chile).

Soils above 3900 m s.m. generally show freezing processes such as slope stripes (structural soil), gelifluction lobes, etc. CZAJKA (1951) found evidence of structural soil associated with stripes in Laguna Verde. In winter, frosts occur practically throughout all day, whereas in summer they only occur at dawn, when dew freezes affecting only the soil surface, mainly in wetlands. IGARZABAL (1984) found a daily freeze-thaw depth of 10-15 cm in bogs located above 4000 m s.m., in the Sierra de Sta. Victoria, Salta Province. In these bogs, permafrost can reach 3 m depth.

### Materials and Methods

In the area studied between Cazadero Grande, at 3400 m s.m., 27°24' SL-68°10' WL, and the spurs of the Cerros Nacimiento and Aguas Calientes, at 4250 m s.m., 27°16' SL-68°30' WL (Fig. 1), 55 phytosociological surveys were carried out. The results are arranged in Table 2. In bogs along the rivers Nacimiento and Aguas Calientes as well as on the slopes, 1 m<sup>2</sup> squares were cut 2 cm onto the soil, the thickness grazed by the camelid fauna, to determine the nutritional values of those samples (Table 3).



Fig. 1. General location of the study area.

	Tm c	Tm f	Ppm	Ppm c	Ppm f
Fiambalá	21,7	9,1	137	100	5
A. de la Sierra	12,0	4,4	105	75	6

Table 1. – Climatic data (according to DE FINA & al., 1959; 1941-50 Period). **Tm c**: mean temperature of the warmest month; **Tm f**: mean temperature of the coldest month; **Ppm**: mean annual precipitation; **Ppm c**: mean precipitation of the warmest three month period; **Ppm f**: mean precipitation of the coldest three month period.

## Results

### *Plant communities*

The analysis of Table 2 permits to establish the following plant communities in the study area:

#### *1. Acantholippia punensis* community (3600-3800 m s.m.).

This community reaches approximately 3800 m s.m., from the extensive Cazaderos Grande alluvial cone, to the W. It is a low shrub vegetation, 0.4-0.5 m high, and very open with a 25-30% mean cover. It is widely distributed on sandy, loess soils. Characteristic shrubs are *Lycium*





*chañar*, *Acantholippia punensis*, *Chuquiraga acanthophylla*, among others, and *Stipa chrysophylla* and *Maihueniopsis ovata* characterize the herbaceous layer.

VERVOOST (1951) distinguished this community as one of the altitudinal zones of the Puna shrub steppe. Several taxa such as *Ch. acanthophylla*, *L. chañar*, etc. are also characteristic of the *Lobivio ferocis-Fabianion densae* alliance RUTHSATZ, 1993 (NAVARRO, 1993) which comprises this community. It constitutes a plant community on rocky slopes between 3500-3900 m s.m. that extends from south-western Bolivia through north-western Argentina and appears to reach the La Rioja Province judging from HUNZIKER's floristic lists (1952). It shows a species impoverishment towards the south and the west.

## 2. *Stipa chrysophylla* community (3800-4100 m s.m.).

This gramineae-dominated community, has a mean cover of 20-30% with a scanty dwarf-shrub vegetation (pulvinate chamaephytes) or with Chamaephytes up to 0.4 m high. It constitutes a very extensive community, from the southern Bolivian Altiplano (NAVARRO, 1993) to Catamarca and, possibly, La Rioja provinces, between 4100-3800 m s.m. in the southern reaches of its distribution. Some of the elements of this community are included in the alliance *Urbanio pappigerae-Stipion frigidae* G. Navarro, 1993, characterized by *Stipa chrysophylla*, *Adesmia subterranea*, *Fabiana bryoides*, *Hoffmannseggia eremophylla*, etc. Among others, there are also *Nardophyllum armatum*, *Stipa speciosa* var. *breviglumis*, *Mulinum ulicinum* and *Stipa frigida*.

Facies of *Nicotiana petunioides*: this community occupies old alluvial cones at 4000 m s.m. and have a groundwater table at approximately 2-2.5 m in depth. *Adesmia subterranea* is the only woody species present. However, approximately 30% of the *Adesmia* plants are dead. This is perhaps due to the lowering of the groundwater table, possibly related to a dry cycle of 3-4 years with little contribution from snow. In these sectors, there is severe soil removal by burrowing rodents (*Ctenomys* sp.) which produces a plant facies of *Nicotiana petunioides*, accompanied by *Calandrinia picta*, *Ipomopsis gossypifera*, *Phacelia setigera* var. *humahuaguensis*, *Cristaria andicola*.

## 3. *Stipa frigida* community (>4000 m s.m.).

This is a microthermic, psammophyllous, gramineous plant community with a 40-50% cover, dominated by *Stipa frigida*, that extends across SSW exposed slopes, over basaltic soils with rocks of different sizes in a sandy matrix. *Fabiana bryoides*, *Adesmia erinacea* and *Ephedra rupestris* characterize the community above 4000 m s.m. This community extends through the Chilean Puna at similar elevation and on similar soil types (LAILHACAR, 1992).

### *Azonal communities*

## 4. *Lampaya hieronymi* community (3800-3900 m s.m.).

*Lampaya hieronymi* and *Adesmia horrida* characterize the base of north-facing slopes where the wind deposits sand, as well as sandy plains. This community is vicariant to the association *Acantholippio salsoloidis-Lampayetum castellanii* G. Navarro, 1993 in Bolivia and probably also in Jujuy (RUTHSATZ, 1977). Between 3800-3900 m s.m. above the Cazadero river, in drier sectors, *L. hieronymi* behaves as a colonizer, maintaining the population through shoots.

## 5. *Distichlis scoparia* saline soils community

This community generally occupies soils close to wetlands and small endorreic basins, in contact with xeric vegetation. The soils are more or less deep, disperse, covered by saline efflo-




Fig. 2. – Wetland profil and wetland occupied by eolic sands:  
 1. N exposed slope with *Ephedra americana* and *Acantholippia punensis*; 2. *Lampaya hieronymi*; 3. *L. hieronymi* and *Festuca orthophylla*; 4. wetland with cryogenic small domes (a) and depressions (b); 5. River.




Fig. 3. – Cryogenic dome profil, with different xericity levels.

Community	<i>C. gayana</i>	<i>R.mandonianus</i>	<i>S. chrysophylla</i>
N total	0,81	1,04	0,45
Prot. total	5,07	6,48	3,06
Ashs (%)	24,53	71,80	6,44
Ca (%)	1,55	8,38	0,16
Mg (%)	0,33	0,66	0,06
P (%)	0,07	0,14	1,76
Greases (%)	4,26	1,76	2,46
Fibers (%)	35,19	10,42	39,05

Table 3. – Nutritional value for three Puna communities (mean values of dried matter of 4 repetitions).



rescences where the water table is between 1.2-1.5 m deep. This community is characterized by *Distichlis scoparia* var. *erinacea*, the dominant species, accompanied by *Triglochin concinna*, *Scirpus nevadensis* var. *remireoides*, etc.

In Catamarca this community is vicariant to the association of *Distichlion humilis-Anthobryetea triandri* G. Navarro, 1993, that reaches its southeastern and southwestern limits in southwestern Bolivia, northeastern Chile and northwestern Argentina (NAVARRO, 1993).

### *Wetland Communities*

These communities have the highest floristic richness. They generally are dominated by grasses on water-saturated soils. The studied wetlands appear to belong to the minerotrophic type (SJÖRZ, 1959; IGARZABAL, 1984). Related to the dynamics of the river, vegetation zones are clearly differentiated according to their response to different degrees of water stress (Fig. 2).

#### **6. *Festuca orthophylla* community**

This plant community constitutes a vegetation zone on drier soils. Where wind accumulates sand between the *Festuca* tufts, xeric elements of the *Stipa frigida* community appear. This community has a mean cover of 60-70% and a height of 0.6 m.

*Deyeuxia curvula* and *D. eminens* var. *fulva* are also present in the community.

#### **7. *Carex gayana* community**

This plant community develops on deep soils (> 1 m), which are alkaline, water-saturated and very rich in organic matter (Table 3). *Carex gayana*, *Eleocharis geniculata*, *Deyeuxia velutina* and *Festuca desvauxii* prevail among others.

In this community, the presence of small cryogenic domes up to 0.4 m high is common, where fines textures prevail toward the surface. These formations characterize the drier sectors within the zone of water-saturated soils, where elements of xeric communities penetrate (Fig. 3).

Comparing this community with the *Ranunculus mandonianus* wetland, the lower values of Mg, Ca and P would be due to the differential partitioning of the assimilates between the aboveground and the belowground portion, and would also be due to the aboveground parts being submitted to grazing. The migration of those elements would respond to environmental conditions such as surface freezing, persistent cold winds, and continuous grazing. These environmental conditions would also explain the high fiber and lipid content as a plant response to mechanical pressure because of the freezing of the surface soils. According to RUTHSATZ (1978), the Cyperaceae dominance in these wetlands would partly be caused by the abundant adventitious roots, which allows the plants to reproduce asexually thus increasing the size of the population.

HALLOY (1983), during years of studying the high Andean wetlands in Tucumán, found no evidence of such asexual plant propagation by mechanisms such as cryoturbation. On the other hand, the abundance of roots would facilitate the absorption of the scarce nutrients such as P and N (poorly mineralized), providing a larger surface for gas exchange in a water-saturated environment.

This plant community is the most heavily grazed by guanacos (*Lama guanicoe*) and llamas (*Lama glama*). *Festuca desvauxii* also occupies gelifluction lobes on shady slopes, in places where snow accumulates, giving those sectors their uneven appearance.




Fig. 4. – Dynamism (xerosere) in *Carex gayana* wetland.

**8. *Oxychloe andina* community**

This plant community develops as a thick and spiny dwarf shrub vegetation near running water, above 4000-4100 m s.m. It is dominated by *Oxychloe andina* with a 100% cover, accompanied by *Scirpus atacamensis*, *Deyeuxia chrysophylla*, *Scirpus deserticola*, *Deyeuxia* aff. *simulans* and *Hypsela reniformis*, etc. It also forms a compact, thin belt at the base of the cryogenic domes, and in wetlands of *C. gayana*, near water-filled depressions. For RUTHSATZ (1978), the dwarf shrub growth forms of *O. andina* would respond to physiological drought (caused by low temperatures) and to low nitrogen content.

**9. *Ranunculus mandonianus* community**

This community, dominated by *Ranunculus mandonianus* and *Puccinellia oresigena* with a 85-90% cover, occupies the wetlands of the Aguas Calientes creek on consolidated, horizontally bedded sandstone substrates. The roots explore a saline soil no more than 10-15 cm deep, where there is horizontal run-off even under the herbaceous layers.

Because the soil development is very poor and saturated with running water, cryoturbation processes do not occur. This would be the reason for the lower fiber and lipid values of the dominant species. The higher values in calcium and ash content are related to the substrate (calcium consolidated sandstones).




Fig. 5. – Mean percentage cover of most conspicuous species from each vegetation unit of the wetland. **I:** *Distichlis scoparia* community; **II:** *Festuca argentina* community; **III:** *Festuca orthophylla* community; **IV:** *Carex gayana* community; **V:** *Oxychloe andina* community; **VI:** *Catabrosa werdermannii* community; **VII:** *Ranunculus mandonianus* community (see Table 2).

### *Other plant communities*

In rivulets, the aquatic plant community of *Potamogeton strictus* can be found, accompanied by *Catabrosa werdermannii*, *Myriophyllum brasiliense*, *Lemna* af. *gibba* and *Ranunculus cymbalaria* f. *exilis*.

The last two communities would indicate the presence in our region of *Puccinellio oresigenae-Oxychloetum andinae* G. Navarro, 1993, described for salty lagoons above 4100 m s.m. in Bolivia.

### *Rock outcrops vegetation*

This saxicolous plant community, characterized by *Senecio santelicensis*, *Cajophora coronata* and *Phacelia setigera* var. *setigera*, accompanied by *Stipa frigida*, can be found on rock outcrop fissures, or on soil deposited on rock edges.

### *Dynamic hypothesis of wetland communities*

Some wetland species tend to occupy habitats where eolic sand is deposited, acting as pioneers of subsequent more xeric stages. For example *Festuca orthophylla* installs itself on the drier, upper part of cryogenic domes produced by freezing of the community of *Carex gayana*. When the sand cover thickens, *Stipa frigida* arrives, indicating the encroachment of xeric communities over the wetland communities (Fig. 4). The same occurs with species from environments in contact with open water such as *Ranunculus mandonianus* and *Deyeuxia eminens*, that remain in small depressions where the water table is closer to the surface than in more xeric units. Fig. 5 shows the mean percentage cover of the most conspicuous taxa from each wetland vegetation unit.

## **Conclusion**

The plant communities phytosociologically studied in the area encompassed by Cazaderos Grande valley, Aguas Calientes and Nacimiento hills, show a close floristic relationship with those from northern Salta, Jujuy and part of southern Bolivia.

The floristic data obtained in this study confirm the inclusion of this zone in the Distrito Jujeño, as it has been proposed by the author, based on the distribution of 107 Puna taxa (MARTINEZ CARRETERO, 1995). New syntaxa in Argentina are indicated, some of them were already suspected: from *Calamagrostietea vicunarum* Rivas Martinez & Tovar, 1982:

- *Lovibio ferocis-Fabianion densae* Ruthsatz, 1993,
- *Urbanio pappigeriae-Stipion frigidae* G. Navarro, 1993,
- *Puccinellio oresigenae-Oxychloetum andinae* G. Navarro, 1993.

Besides, a dynamic xerosere scheme of the wetlands of the study area is proposed. A close relationship is found between plant communities and cryogenic processes. This is especially pronounced in wetlands with a moisture gradient that shows a remarkable fall from the water-saturated wetland to the cryogenic domes surface, where the species from xeric communities such as *Festuca orthophylla*, *Stipa frigida*, etc. colonize. It can also be pointed out that, starting from 3900-4000 m s.m., microthermal elements prevail associated with soils that indicate seasonal freezing, such as *Stipa frigida*, *Deyeuxia antoniana*, *Ephedra rupestris*, and others.

## ACKNOWLEDGEMENTS

To Fidel Roig and Arturo Corte for their comments and discussion of the text, and N. Horak for reviewing the English version.

## REFERENCES

- ALBA, E. (1979). Sistema de Famatina. *In: Geología Regional Argentina, 2º Simposio Academia Nacional de Ciencias, Córdoba* 1: 349-396.
- ALLMENDIGER, R. W. (1986). Tectonic development, Southeastern border of the Puna plateau, northwestern Argentine Andes. *Bull. Geol. Soc. Amer.* 97: 1070-1082.
- CABRERA, A. L. (1957). La Vegetación de la Puna Argentina. *Rev. Invest. Agric.* 11: 317-412.
- CABRERA, A. L. (1968). Ecología vegetal de la Puna Argentina. *In: TROLL, C., Geocology of the mountainous regions of the tropical americas. Colloq. Geogr.* 9: 117-121.
- CABRERA, A. L. (1976). *Las regiones fitogeográficas argentinas*. ACME Eds.
- CABRERA, A. L. & A. WILLINK (1973). *Biogeografía de América Latina*. OEA Serie de Biología, 120 pp.
- CZAJKA, G. (1951). Resultados de un viaje a la cuenca de Laguna Verde (Tinogasta-Catamarca) I. Observaciones geomorfológicas. *Actas de la XV Semana de Geografía. GAEA*: 39-49.
- DE FINA, A. L., A. GARBOSKY & L. SABELLA (1959). Difusión geográfica de cultivos índices en las provincias de Catamarca y La Rioja y sus causas. *Publ. Inst. de Suelos y Agrotécnia* n° 63.
- HALLOY, S. (1983). Movimientos criogénicos del suelo. *Actas 1a. Reunión Grupo Periglacial Argentino*: 109-132, IANIGLA.
- HUNZIKER, J. (1952). Las comunidades vegetales de la Cordillera de La Rioja. *Rev. Invest. Agric.* 11: 217-242.
- IGARZABAL, A. P. (1984). Comportamiento hidrológico de las turberas de montaña como estructuras criogénicas en las regiones de Puna y Cordillera Oriental. *Actas de la Segunda Reunión Grupo Periglacial Argentino*: 106-115.
- LAILHACAR, S. R. (1992). Evaluación nutritiva de los recursos forrajeros nativos y naturalizados de la II región. *Avances en Producción Animal* 17: 23-44.
- MARTINEZ CARRETERO, E. (1995). La Puna Argentina. Demarcación general y división en distritos florísticos. *Bol. Soc. Argent. Bot.* 31: 27-40.
- NAVARRO, G. (1993). Vegetación de Bolivia: El altiplano meridional. *Rivas Godaya* 7: 69-98.
- RUTHSATZ, B. (1977). Pflanzengesellschaften und ihre Lebensbedingungen in den Andinen Halbwüsten Nordwest Argentiniens. *Diss. Bot.* 39: 1-168.
- RUTHSATZ, B. (1978). Las plantas en cojín de los semidesiertos andinos del Noroeste argentino. *Darwiniana* 21: 491-520.
- SJÖRS, H. (1959). Bogs and fens in the Hudson Bay Lowlands. *Artic* 12: 3-19.
- TURNER, J. C. & V. MENDEZ (1979). Puna. *In: Geología Regional Argentina, 2º Simposio Academia Nacional de Ciencias, Córdoba* 1: 13-56.
- VERVOORST, F. (1951). Resultados de un viaje a la cuenca de Laguna Verde (Tinogasta-Catamarca) III. Observaciones sobre la vegetación entre Tinogasta y la cuenca. *Actas de la XV Semana de Geografía. GAEA*: 61-67.