

Some new and interesting lichens, allied fungi and lichenicolous fungi from southern and central Switzerland

Autor(en): **van den Boom, P.P.G. / Clerc, Philippe**

Objektyp: **Article**

Zeitschrift: **Candollea : journal international de botanique systématique = international journal of systematic botany**

Band (Jahr): **55 (2000)**

Heft 1

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-879509>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Some new and interesting lichens, allied fungi and lichenicolous fungi from southern and central Switzerland

P. P. G. VAN DEN BOOM
&
PHILIPPE CLERC

ABSTRACT

BOOM, P. P. G. van den & P. CLERC (2000). Some new and interesting lichens, allied fungi and lichenicolous fungi from southern and central Switzerland. *Candollea* 55: 87-103. In English, English and French abstracts.

Lichens, allied fungi and lichenicolous fungi, collected during a trip through southern and central Switzerland are listed. The following annotated list contains 258 identified species that are new and/or rare in Switzerland. *Agonimia vouauxii* (de Lesd.) ined., *Bacidia absistens* (Nyl.) Arnold, *Biatoridium delitescens* (Arnold) Hafellner, *Dermatocarpon leptophyllum* (Ach.) Lang, *Hymenelia similis* (A. Massal.) Poelt & Vezda, *Lepraria* aff. *flavescens* Clauzade & Cl. Roux, *Micarea botryoides* (Nyl.) Coppins, *M. leprosula* (Th. Fr.) Coppins & A. Fletcher, *Protoparmelia hypotremella* Herk, Spier & Wirth, *Rinodina rinodinoidea* (Anzi) H. Mayrhofer & Scheid., *Strigula* aff. *phaea* (Ach.) R. C. Harris, *Thelopsis flaveola* Arnold, *Tremella coppinsii* Diederich & Marson, *T. phaeophysciae* Diederich & M. S. Christ have been recorded for the first time in the country.

RÉSUMÉ

BOOM, P. P. G. van den & P. CLERC (2000). Nouveautés pour les lichens, champignons lichénicoles et autres champignons non lichénisés usuellement traités avec les lichens, de Suisse centrale et méridionale. *Candollea* 55: 87-103. En anglais, résumés anglais et français.

Une liste de 258 espèces de lichens, champignons lichénicoles et autres champignons non lichénisés usuellement traités avec les lichens est présentée, résultat d'une série d'excursions en Suisse, à travers l'Oberland bernois et le Valais central. *Agonimia vouauxii* (de Lesd.) ined., *Bacidia absistens* (Nyl.) Arnold, *Biatoridium delitescens* (Arnold) Hafellner, *Dermatocarpon leptophyllum* (Ach.) Lang, *Hymenelia similis* (A. Massal.) Poelt & Vezda, *Lepraria* aff. *flavescens* Clauzade & Cl. Roux, *Micarea botryoides* (Nyl.) Coppins, *M. leprosula* (Th. Fr.) Coppins & A. Fletcher, *Protoparmelia hypotremella* Herk, Spier & Wirth, *Rinodina rinodinoidea* (Anzi) H. Mayrhofer & Scheid., *Strigula* aff. *phaea* (Ach.) R. C. Harris, *Thelopsis flaveola* Arnold, *Tremella coppinsii* Diederich & Marson, *T. phaeophysciae* Diederich & M. S. Christ sont toutes des espèces nouvelles pour la Suisse.

KEY-WORDS: Lichens – Lichenicolous fungi – Switzerland.

Introduction

During a study trip through the Rhône valley (canton of Valais) and the area near Brienz (canton of Bern), in the summer of 1996, 20 sites were visited by the first author in order to survey the lichen flora. A part of this area close to Evolène (Val d'Hérens, canton of Valais) has

a relatively well-known lichen flora where the Dutch Bryological and Lichenological Study group recorded about 600 lichens and lichenicolous fungi during the recent field work it undertook (BOOM & al., 1993).

The present paper lists lichens and lichenicolous fungi that were found in the course of the survey of several localities ranging from 650 to 2200 m in the Rhône valley. The lichen flora of the Leuk area, near Sierre, is not particularly rich. It was however, gratifying to find some very interesting lichens on a fallen, rotting trunk, among them, an undescribed species of the genus *Gyalideopsis* which was growing close to *Absoconditella lignicola* Vezda & Pisut and *Catinaria atropurpurea* (Schaer.) Vezda & Poelt.

The survey of the area near Visperterminen did not show a particularly important lichen flora, with the exception of a small population of *Cetraria obtusata* (Schaer.) van den Boom & Sipman found among common terricolous lichens like *Catapyrenium cinereum* (Pers.) Körb., *Cetraria ericetorum* Opiz, *C. nivalis* (L.) Ach., *Dibaes baeomyces* (L. fil.) Rambold & Hertel, "*Lecidea*" *hypnorum* Lib. and *Rinodina mniarea* (Ach.) Körb. The most significant specimens from this area were "*Lecidea*" *rufofusca* (Anzi) Nyl. and *Rinodina conradii* Körb. which were growing in association with *Caloplaca sorocarpa* (Vain.) Zahlbr. and *Lecanora salicicola* H. Magn. on *Rhododendron* branches.

Rinodina rinodinoides (Anzi) Scheid., a rare species found in this area, was growing on a schistose stone on a west-exposed roof, of a barn in a meadow. It was growing among brown and yellow *Rhizocarpon* spp.

The sites covered by a survey at the area of Brienz (Bernese Oberland) vary from a very wet glacier valley to open fields with calcareous outcrops and low shrubs, from roadside trees to mixed forests, ranging from 960 to 2100 m. The glacier valley, with its varied exposed and damp rocks yielded an interesting lichen flora with extensive patches of *Polyblastia ardesiaca* (Bagl. & Car.) Zschacke growing with *Lepraria* sp. in most shaded areas. From the calcareous permanently wet rocks at the periphery of a gorge, that was within the radius of the spray of a waterfall, only crustose inconspicuous lichens like *Hymenelia similis* (A. Massal.) M. Choisy, *Ionaspis melanocarpa* (Kremp.) Arnold, *Eiglera homalomorpha* (Nyl.) Clauzade & Roux and *Thelidium incavatum* Mudd were collected. The first species seems to prefer habitats similar to this gorge and was present in a small amount.

The last important site, south of Brienz, along the stream Giessbach, was also the most interesting owing to its rather well developed *Lobarion* community found in a mixed forest with *Acer*, *Picea* and *Sorbus*. Interspersed in well-developed fertile patches of *Lobaria pulmonaria* (L.) Hoffm. were *Bacidia absistens* (Nyl.) Arnold, *Dimerella lutea* (Dicks.) Trevis, *Hypotrachyna sinuosa* (Sm.) Hale, *Megalaria pulvereae* (Borrer) Hafellner & Schreiner and *Parmeliella triptophylla* (Ach.) Müll. Arg. The last mentioned species was abundantly fertile.

Material and methods

Herbarium records have been included in table 1. Material collected during the survey was deposited in the private herbarium of the first author. Duplicates of some important collections were deposited in G. Standardized thin-layer chromatography (TLC) (CULBERSON & AMMANN, 1979) was used for identification of *Lepraria* sp. and some other collections.

The specimens which are listed below were studied mostly according to PURVIS & al. (1992), TÜRK & POELT (1993), SANTESSON (1993) or more recent publications were used mainly for the nomenclature. Coordinates (Coord.) are given according to the national grid system of Switzerland.

Collecting sites

Canton of Valais

- 1 = ENE of Sierre, N of Leuk, Albinen, slope along road, with *Abies* wood. Coord.: 615.000-132.300, alt. 1200 m, 2 August 1996.
- 2 = ENE of Sierre, N of Leuk, 0.2 km N of Albinen, along a (dry) stream on a slope. Coord.: 615.100-132.500, alt. 1260 m, 2 August 1996.
- 3 = ENE of Sierre, N of Leuk, 1.5 km N of Albinen, Dorbugrabu, *Picea* wood. Coord.: 615.000-133.700, alt. 1600 m, 2 August 1996.
- 4 = SW of Brig, SSE of Visp, Visperterminen, 1 km E of the village, path to Simplonpass and Gibidumpass, path W of a small lake. Coord.: 637.500-123.000, alt. 2050 m, 3 August 1996.
- 5 = SW of Brig, SSE of Visp, Visperterminen, 2 km E of the village, path to Simplonpass and Gibidumpass, near a small lake. Coord.: 638.600-123.000, alt. 2200 m, 3 August 1996.
- 6 = SW of Brig, SSE of Visp, Visperterminen, nearby the village, old barn and acid boulders in a meadow. Coord.: 636.000-123.000, alt. 1200 m, 3 August 1996.
- 7 = E of Sierre, 2 km south of Agarn, path to Unter-Meretschialp, north exposed *Picea* wood on a slope with big boulders and outcrops. Coord.: 616.900-125.400, alt. 1450 m, 4 August 1996.
- 8 = E of Sierre, 2.5 km E of Susten, N-S gorge, at the entrance, with calcareous rocks. Coord.: 617.900-128.400, alt. 650 m, 4 August 1996.
- 9 = SE of Sierre, Val d'Anniviers (N), Soussillon, path to Chandolin, Pramin, north side of stream, in a forest on a slope with calcareous outcrops. Coord.: 611.400-123.200, alt. 1600 m, 5 August 1996.
- 10 = SE of Sierre, Val d'Anniviers (N), Soussillon, path to Chandolin, Les Invers, south side of stream, wood on a slope with *Salix* and calcareous outcrops. Coord.: 611.500-122.900, alt. 1700 m, 5 August 1996.
- 11 = NW of Gletsch, Grimselpass, south exposed granite outcrops, S side of road. Coord.: 670.000-157.400, alt. 2160 m, 6 August 1996.

Canton of Bern

- 12 = NE of Grindelwald, Rosenlauri, road to Schwarzwaldalp, glacier valley, near outlet of gorge and damp area with calcareous outcrops near a waterfall. Coord.: 654.900-169.900, alt. 1380 m, 7 August 1996.
- 13 = 5 km S of Brienz, path from Chüemad to Tschingel, NW exposed meadow on a slope with calcareous outcrops. Coord.: 645.900-173.000, alt. 1850 m, 8 August 1996.
- 14 = 5 km S of Brienz, path from Chüemad to Tschingel, SW side of Tschingel, exposed calcareous outcrops. Coord.: 646.300-172.800, alt. 2000 m, 8 August 1996.
- 15 = 6 km SSE of Brienz, path E from Tschingel, S side of Axalphorn, exposed calcareous outcrops in a meadow on a slope. Coord.: 648.400-173.100, alt. 2100 m, 8 August 1996.
- 16 = 6 km SSE of Brienz, path NE from Axalphorn, Urserli, calcareous outcrops in karst area. Coord.: 648.200-173.600, alt. 1800 m, 8 August 1996.
- 17 = 3 km S of Brienz, road along Giessbach, 500 m after crossing, *Picea* wood with *Fagus*, big *Acer* and *Fraxinus* along 'road'. Coord.: 644.900-175.400, alt. 960 m, 9 August 1996.

- 18 = 3 km S of Brienz, road along Giessbach, 600 m after crossing, cave with steep calcareous outcrops. Coord.: 644.900-175.300, alt. 1100 m, 9 August 1996.
- 19 = 3 km S of Brienz, road along Giessbach, 700 m after crossing, near cottage of second bridge, *Alnus incana* along stream and big boulders. Coord.: 644.900-175.200, alt. 1080 m, 9 August 1996.
- 20 = 4.5 km S of Brienz, road along Giessbach, 100 m W of Giessbach, Schwand, north exposed wood on a slope with *Acer*, *Picea* and *Sorbus*. Coord.: 644.600-173.800, alt. 1200 m, 9 August 1996.

Annotated list of most interesting species

Abrothallus bertianus De Not. s.l.

Loc. 20, v.d. Boom 18113, on *Acer*: this lichenicolous fungus has been found on *Melanelia glabratula* (Lamy) Essl. According to the present literature, this species has been reported from several host species in *Cetraria* and *Parmelia* s.l. However the characteristics of this specimen do not exactly fit the description in CLAUZADE & al. (1989). A revision seems to be necessary. The ascospores are 1-septate and their dimensions are 12-15 × 5-6 µm, but the epithecium is brown and K⁺ greenish.

Absconditella lignicola Vezda & Pisút

Loc. 2, v.d. Boom 18132, on rotting wood of a big fallen trunk: it was growing close to an undescribed *Gyalideopsis*. *Absconditella lignicola* is already known from the canton of Grisons (DIETRICH & al., 1992) and central Switzerland (RUOSS, 1992). New to the canton of Valais.

Agonimia vouauxii (de Lesd.) comb. ined.

Syn. *Polyblastia vouauxii* de Lesd.

Loc. 13, v.d. Boom 18142, open place among low calcareous outcrops, on plant debris: *Agonimia vouauxii* has a granular to squamulose greenish to brownish papillate thallus, blackish perithecia (0.15-0.25 mm in diam.), 2-spored asci and large muriform ascospores (50-70 × 15-25 µm). Known to be present in dune areas in the Netherlands (BRAND & al., 1988) and NW France (de LESDAIN, 1910, 1914; ZSCHACKE, 1934), *A. vouauxii* has also been reported from Austria (TÜRK & POELT, 1993). In a paper on lichens and lichenicolous fungi from Belgium and Luxemburg, which is in preparation by M. Brand, P. Diederich, E. Sérusiaux & P. van den Boom, taxonomic notes are enclosed on some *Agonimia* species. This species is recorded for the first time in Switzerland.

Arthonia mediella Nyl.

Loc. 1, v.d. Boom 17747, 17750, on *Abies* in an *Abies* forest: *Arthonia mediella* is a species which is mainly found on conifers in boreal areas and the mountains of central Europe. New to the canton of Valais.

Arthonia muscigena Th. Fr.

Loc. 3, v.d. Boom 18122, on twigs of *Salix*, along a path in an *Abies* forest: this specimen was found among *Bacidia beckhausii* Körb., *B. subincompta* (Nyl.) Arnold and *Biatora helvola* Hellb.

The corticolous lichen flora on twigs and branches must have been much overlooked in Switzerland. Even the recent survey of DIETRICH & SCHEIDEGGER (1997) did not include it. This recording is interesting because it was found at an altitude of 1600 m. According to PURVIS & al. (1992), *A. muscigena* is a lowland species, known from western Europe. However according to WIRTH (1995b), this species is prevalent from central Europe to submediterranean areas at an altitude of c. 400 to 1100 m and has already been reported in Switzerland.

Bacidia absistens (Nyl.) Arnold

Loc. 19, v.d. Boom 18067, on *Alnus*; loc. 20, v.d. Boom 18095, on *Acer*: this species is mainly distributed in European countries along the Atlantic, but it is sometimes mentioned from continental areas, for example in the Alps (KALB, 1970). This species is recorded for the first time in Switzerland.

Bacidia vermifera (Nyl.) Th. Fr.

Loc. 12, v.d. Boom 17984, on *Acer* in a mixed forest. This *Bacidia* species has characteristic sigmoid spores, with dimensions of 25-30 × 2.5 µm. Conidia are short and bacilliform (c. 3.5 × 1.5 µm). The name *Bacidia hegetschweileri* (Hepp) Vain. has often been used for this species (EKMAN, 1996), but in his work, Ekman shows that the use of *B. vermifera* should be resumed. The absence of a green pigment in the hymenium and the proper exciple distinguish this species from *B. circumspecta* (Vain.) Malme (EKMAN, 1996). It is similar to *B. subcircumspecta* Coppin in pigmentation, but has different ascospores and conidia (PURVIS & al., 1992). In Switzerland this species is known only from Zurich. New to the canton of Valais.

Biatora ocelliformis (Nyl.) Arnold

Loc. 20, v.d. Boom 18110, on *Acer* in a mixed forest: this specimen is similar to material collected in southern Austria (Kärnten) where this species is locally rather common. According to PRINTZEN (1995) *B. ocelliformis* is widely distributed in the boreal zone, mainly in the more eastern part of central Europe. Recently this specimen was reported from Switzerland by PRINTZEN & PALICE (1999).

Biatora vernalis (L.) Fr.

Loc. 20, v.d. Boom 17919, on mosses, on vertical, wet and shaded calcareous outcrops: according to PRINTZEN (1995) *B. vernalis* has generally been mistaken for *B. subduplex* (Nyl.) Printzen, the most frequently found *Biatora* species in Europe. The latter species however has smaller spores (PRINTZEN, 1995, fig. 34) and a different ecology. PRINTZEN (1995) is intrigued by the absence of *B. vernalis* in southern Germany, Austria and Switzerland. This species had already been reported for Switzerland by STIZENBERGER (1882-1883) and several other lichenologists (LETTAU, 1919, 1954; MEYLAN, 1922; FREY, 1923, 1937; ZSCHACKE, 1926; BOOM & al., 1993; HINTEREGGER, 1994). Not all these specimens have been checked but those that have been checked turned out to be *B. subduplex* (PRINTZEN, 1995). Our specimen has been seen and identified by C. Printzen.

Biatoridium delitescens (Arnold) Hafellner

Loc. 20, v.d. Boom 18082, on *Acer* in a mixed forest: this species has been recorded in central, northern Europe and western Scotland. Being an extremely inconspicuous species, it is often overlooked. According to HAFELLNER (1994), in central Europe, it was reported not only from southern Germany but also from Spain. The Spanish record has been published by ETAYO & al. (1993). This species is recorded for the first time in Switzerland.

Caloplaca aurea (Schaer.) Zahlbr.

Loc. 15, v.d. Boom 17990, on vertical, east exposed calcareous outcrops in a meadow on a slope, growing over sand in crevices: this species has already been published by STIZENBERGER (1882-1883), MIGULA (1929), POELT (1965) and WESTBERG & KÄRNEFELT (1998) and it is found in the following cantons: Bern, Luzern, Nidwald, Obwald, Uri, Vaud and Valais.

Caloplaca chlorina (Flot.) H. Olivier

Loc. 12,13, v.d. Boom 17952, 17968, on a calcareous boulder in a meadow on a slope and on *Acer* near the outlet of a gorge: the specimens found were well developed. It as compared with the type collection of *C. isidiigera* Vezda (= *C. chlorina*). The thallus of the fertile epilithic collection is coarsely cracked and areolated, with an unevenly granular to isidiate surface. In the epiphytic specimen, the thallus is fertile, but without isidia-like granules.

Carbonea supersparsa (Nyl.) Hertel

Loc. 7, v.d. Boom 17846, on schist, growing on *Lecanora polytropa* in a shaded spot in a forest: previously known from Switzerland, but rarely collected, the species has already been reported and described by STIZENBERGER (1882-1883).

Catillaria picila (A. Massal.) Coppins

Loc. 16, v.d. Boom 18020, on a vertical and shaded calcareous rock: this rare species is characterized by the dark brown apothecia that are \pm constricted at base and 0.4-0.7 mm in diam.; the 0-(1) septate hyaline ellipsoid spores and the relatively wide hyphae of the excipulum. New to the canton of Bern.

Cercidospora sp.

Loc. 15, v.d. Boom 17993, on a calcareous rock, growing in the apothecia of *Aspicilia contorta* (Hoffm.) Kremp.: this lichenicolous fungus is closely related to *Cercidospora epipolytropa* (Mudd) Arnold, a species widely distributed throughout Europe and usually found on thallus and apothecia of *Lecanora polytropa* (Hoffm.) Rabenh. and related species.

Cetraria obtusata (Schaer.) van den Boom & Sipman

Loc. 5, v.d. Boom 17821, terricolous: this species was studied by BOOM & SIPMAN (1994) and already recorded in Switzerland (Bern, Valais and Uri). Its distribution is restricted to the alpine zone. It has often been mistaken for *C. aculeata* (Schreb.) Fr. or *C. muricata* (Ach.) Eckfeldt, but the yellow/orange C+ deep orange medulla and the typical pycnidia-bearing projections around the pseudocyphellae are typical identification marks.

Chaenothecopsis consociata (Nádv.) A. F. W. Schmidt

Loc. 3, v.d. Boom 17769, 17789, on *Picea* in a forest: it was growing on thallus of *Chaenotheca chrysocephala* (Ach.) Th. Fr., associated with *Calicium trabinellum* (Ach.) Ach. New to the canton of Valais.

Chaenothecopsis viridireagens (Nádv.) A. F. W. Schmidt

Loc. 12, *v.d. Boom 17940*, on a rotting overhanging stump in a damp area, nearby a waterfall: new to the canton of Bern.

Chromatochlamys muscorum (Nyl.) H. Mayrhofer & Poelt var. *muscorum*

Loc. 7, 10, 19, *v.d. Boom 17851, 17852, 17921, 18095*, growing on acid or calcareous boulders over mosses in rather extensive patches in open, but sheltered spots: new to the canton of Bern.

Dermatocarpon leptophyllum (Ach.) K. G. W. Lang

Loc. 8, *v.d. Boom 17863*, on a vertical, shaded and sheltered calcareous rock: this species is recognized by the single holdfast, the subglobose ascospores and the thinly pruinose, brownish thallus (ORANGE, 1998). This species is recorded for the first time in Switzerland.

Dimerella lutea (Dicks.) Trevis.

Loc. 20, *v.d. Boom 18076*, on *Acer* in a mixed forest, on a tree marked for logging: the *Lobarion* occurs here quite luxuriantly with species like *Hypotrachyna sinuosa*, *Nephroma bellum* (Spreng.) Tuck., *Parmeliella triptophylla* (fertile) and fertile *Lobaria pulmonaria*. New to the canton of Bern.

Dirina massiliensis Durieu & Mont. f. *sorediata* (Müll. Arg.) Tehler

Loc. 18, *v.d. Boom 18041*, on a steep shaded and sheltered calcareous rock in a forest on a slope: this species was associated with *Lepraria* aff. *flavescens* and *Leproplaca chrysodeta* (Räsänen) Laundon. New to the canton of Bern.

Eiglera homalomorpha (Nyl.) Clauzade & Roux

Loc. 12, *v.d. Boom 18127*, on a damp vertical sheltered calcareous rock: a small collections characterized by a I + blue apical dome of the asci, broad ellipsoid ascospores and a bluish N + red epithecium. This specimen has been compared with material from Austria, where this species is rarely collected (TÜRK & POELT, 1993). Previously recorded from Switzerland by BOISSIÈRE & al. (1989).

Endococcus propinquus (Körb.) D. Hawksw.

Loc. 10, *v.d. Boom 17977*, growing on an unidentified epilithic crust, on a vertical and wet rock in a wood on a slope: a species widely distributed in Europe, occurring on a wide range of hosts.

Farnoldia micropsis (A. Massal.) Hertel

Loc. 15, *v.d. Boom 17992*, on a calcareous outcrop in a meadow on a slope: new to the canton of Bern.

***Fuscidea lightfootii* (Sm.) Coppins & P. James**

Loc. 19, *v.d. Boom 18065*, on *Alnus*; loc. 20, *v.d. Boom 18107*, on *Acer*: the small specimen found was fertile, but only with a few apothecia. The typical ascospores, medianly constricted with obtuse ends, in combination with divaricatic acid in the thallus are typical for this species. At locality 19, it was found among sorediate species like *Buellia griseovirens* (Sm.) Almb. It is most probably rare in Switzerland. New to the canton of Bern.

***Hymenelia similis* (A. Massal.) Poelt & Vezda**

Loc. 12, *v.d. Boom 17939*, on a vertical and wet calcareous rock in the spread of a waterfall: this species was found several times in Carinthia (Austria) only in deep gorges where it was growing also under wet conditions (BOOM & al., 1996). The specimens in G have been checked and they appear to have been collected on the Salève in France and not in Switzerland as indicated by STIZENBERGER (1882-1883). This species is recorded so far for the first time in Switzerland.

***Hypotrachyna sinuosa* (Sm.) Hale**

Loc. 20, *v.d. Boom 18129*, on young *Acer* in mixed forest: this is an endangered species in Switzerland (CLERC & al., 1992). It was growing with *Megalaria pulverea*, *Lobaria pulmonaria* (fertile) and *Parmeliella triptophylla*.

***Lecania* cf. *cyrtellina* (Nyl.) Sandst.**

Loc. 10, *v.d. Boom 17883*, on *Salix*; loc. 16, *v.d. Boom 18008*, on a shrub (cf. *Rhododendron*): the apothecia are \pm biatorine, the ascospores are relative small (c. 2.5-3 μm) and mostly non-septate. These characteristics are typical for *L. cyrtellina*, but because of the lack of macroconidia, the identification is not certain.

***Lecania inundata* (Hepp ex Körb.) M. Mayrhofer**

Loc. 8, *v.d. Boom 17866*; loc. 18, *v.d. Boom 18048*, on a calcareous rock: in the latter it was growing on a steep rock face among *Lecania suavis* (Müll. Arg.) Mig. in a forest on a slope. New to the canton of Valais and to the canton of Bern.

***Lecania suavis* (Müll. Arg.) Mig.**

Loc. 18, *v.d. Boom 18043*, on a shaded and sheltered calcareous rock in a forest on a slope: mentioned in BOOM & al. (1993) for the canton of Valais. New to the canton of Bern.

***Lecanora handelii* J. Steiner**

Loc. 7, *v.d. Boom 17838*, on ferro-rich siliceous shaded outcrop, along a forest path: it was growing with *Lecanora epanora* (Ach.) Ach., *L. polytropa* and *Miriquidica leucophaea* (Rabenh.) Hertel & Rambold. New to the canton of Valais.

***Lecanora jamesii* J. R. Laundon**

Loc. 3, *v.d. Boom 17765*, on wood and on soil along a path in a *Picea* forest: *Lecanora jamesii* is mainly known in regions along the Atlantic ocean, from the British Isles to Portugal,

where it is locally abundant. Although this species was mentioned from the canton of Valais before (HANKO, 1983), it has been collected only rarely in Switzerland and Austria (TÜRK & POELT, 1993).

“*Lecidea*” *rufofusca* (Anzi) Nyl.

According to PRINTZEN (1995: 208) *Biatora porphyroplaca* Hinteregger & Poelt is probably identical with *L. rufofusca*.

Loc. 5, v.d. Boom 18123, on *Rhododendron*: already known from the canton of Valais (BOOM & al., 1993), this species has rarely been collected. It is known in boreal-alpine areas from a wide range of substrata, such as terricolous or epiphytic mosses, plant debris, shrubs etc. It has already been published for Switzerland by STIZENBERGER (1882-1883), MIGULA (1931) and HINTEREGGER (1994).

Lempholemma botryosum (A. Massal.) Zahlbr.

Loc. 18, v.d. Boom 18040, on a steep and shaded calcareous rock: this species was found in a species poor community, with only *Caloplaca cirrochroa* (Ach.) Th. Fr. as accompanying species.

Lepraria* cf. *cacuminum (A. Massal.) Lothander

Loc. 11, v.d. Boom 17873, on a south facing vertical surface on a granite outcrop: this specimen contains atranorin and most probably porphyritic acid (Rf 2-3 in solvent A). It was compared with specimens identified by Prof. C. Leuckert.

Lepraria* aff. *flavescens Clauzade & Roux

Loc. 18, v.d. Boom 18042, 18046, on a sheltered, shaded and steep calcareous rock: according to WIRTH (1995a) *L. flavescens* is the only species which occurs on calcareous rock and has a C+yellow thallus reaction. Our specimens contain atranorin and the unknown 5-6/2-3/5 (TLC); Spot tests give C+ deep yellow, K+yellow and P+yellow reactions. Morphologically it is similar to *L. flavescens* and it is most probably related to this species. It is recorded for the first time in Switzerland.

Lepraria nivalis J. R. Laundon

Loc. 12, v.d. Boom 17941, on a shaded vertical calcareous outcrop influenced by a waterfall: atranorin and protocetraric acid are detected by TLC. The thallus is white, ± lobed-like. Morphologically it is identical with a specimen collected in a similar habitat in Austria (BOOM & al., 1996). The specimen from loc. 16, v.d. Boom 18009, belongs to another chemotype, with gyrophoric acid as main substance and protocetraric acid. According to WIRTH (1995b), this species occurs in the Swiss part of the Jura.

Leptochidium albociliatum (Desm.) M. Choisy

Loc. 9, v.d. Boom 18125, on a well-lit calcareous rock among moss: associated species is *Caloplaca stillicidiorum* (Vahl.) Lynge. It is a small sterile collection, but with globose isidia and the characteristic convex, white-pubescent lobes. It is, however, similar to fertile collections (hb. v.d. Boom) from southern Europe.

Leptorhaphis epidermidis (Ach. ex Hepp) Th. Fr.

Loc. 12, v.d. *Boom 17955*, on *Betula*: The thallus seems to be absent in this specimen.

Lichenoconium erodens M. S. Christ. & D. Hawksw.

Loc. 9, v.d. *Boom 17909*, on *Lecanora* sp. on *Larix*: this common lichenicolous species is known from a wide range of hosts and is widely distributed in Europe.

Lichinella nigritella (Lettau) Moreno & Egea

Loc. 8, v.d. *Boom 17861*, on a vertical, moderately shaded and sheltered calcareous outcrop: this specimen was compared with the following exsiccatum: *Arnold Lich. Exsicc. No 1596* (*Collema furvum* Ach. f. *conchilobum* Flot. 'An Porphyrfelsen im Dorfe Jenesien bei Bozen; Tirol. Aug. 1893, collector: Kernstock') that is mentioned by LETTAU (1942) in the protologue as being *L. nigritella*. Both specimens obviously belong to the same species. *Lichinella nigritella* is a central European species with some known localities in the northern part of the mediterranean region (NIMIS, 1993).

Loxospora cismonica (Beltr.) Hafellner

Loc. 20, v.d. *Boom 18105*, on *Acer* in a mixed forest: according to DIETRICH & SCHEIDEGGER (1996), this species is rare in the pre-Alps and the central Plateau.

Megalalaria pulverea (Borrer) Hafellner & E. Schreiner

Loc. 20, v.d. *Boom 18098*, on *Acer* in a mixed forest: this species was growing abundantly in the *Lobarion* community. The specimen is sterile.

Micarea botryoides (Nyl.) Coppins

Loc. 3, v.d. *Boom 18136*, on a rotting wood along a path in a *Picea* forest: only a small collection with pynidia which contain the characteristic dark K- pigment, but apothecia are not found. It was growing together with *Dimerella pineti* (Ach.) Vezda. It was most probably overlooked in past surveys and is recorded here for the first time in Switzerland.

Micarea leprosula (Th. Fr.) Coppins & A. Fletcher

Loc. 19, v.d. *Boom 18073*, on calcareous outcrops, growing over mosses: this species was found in large amounts, containing typical pale greenish soorediate (sterile) patches. It is recorded for the first time in Switzerland.

Micarea sylvicola (Flot.) Vezda & V. Wirth

Loc. 7, 10, v.d. *Boom 17842, 17878*, on a vertical shaded and sheltered acid rock: this species has not been reported from Switzerland since the last century. In STIZENBERGER (1882-1883) this species was reported from the canton of Uri. New to the canton of Valais.

Miriquidica leucophaea (Flörke ex Rabenh.) Hertel & Rambold

Loc. 7, on a ferro-rich siliceous rock along a path in a forest: this species was growing close to *Lecanora polytropa*. New to the canton of Valais.

Placynthium filiforme (Garov.) M. Choisy

Loc. 14, v.d. Boom 17985, on a vertical and slightly overhanging and shaded calcareous rock: it was found growing close to *Squamarina cartilaginea* (With.) P. James, *S. gypsacea* (Sm.) Poelt and *Toninia alutacea* (Anzi) Jatta.

Placynthium tremniacum (A. Massal.) Jatta

Loc. 9, v.d. Boom 17905, on a calcareous outcrop in a forest on a slope: this specimen was collected together with *P. nigrum* (Huds.) Gray which was growing in close proximity. The 1-septate spores and pruinose thallus with somewhat stouter, \pm flat minute squamules, different marginal lobes, and the less developed prothallus distinguish this species from *P. nigrum* (GYELNIK, 1940).

Polyblastia ardesiaca Zschacke

Loc. 12, v.d. Boom 17937, 17945, on a calcareous rock: *Polyblastia ardesiaca* was previously known from two localities in Switzerland: Val Tremola and Guttannen (ZSCHACKE, 1934). This species was growing very abundantly on shaded and permanently wet rocks close to a waterfall in a deep gorge, associated with *Lepraria* sp.

Porina linearis (Leight.) Zahlbr.

Loc. 18, v.d. Boom 18054, on a steep shaded north facing calcareous outcrop: new to the canton of Bern.

Protoparmelia hypotremella Herk, Spier & V. Wirth

Loc. 19, v.d. Boom 18069, on *Alnus incana* along stream: this species is characterized by its granular thallus. The typical granules are isidia-like, convex, rounded to elongate or globose (APTROOT & al., 1997). This sterile species has never been found with ascomata or conidiomata. Our specimen was compared with specimens from north-western Europe (hb. v.d. Boom, hb. Diederich) where this species is locally common. It is recorded for the first time in Switzerland.

Rinodina griseosoralifera Coppins

Loc. 10, v.d. Boom 17923, on *Salix*; loc. 19, v.d. Boom 18058, on *Acer*: both collections were sterile but contain many characteristic discrete soralia. Atranorin and zeorin were detected with TLC. Only a few specimens have been recorded in the Alps (TÜRK & POELT, 1993). New to the canton of Valais and to the canton of Bern.

Rinodina orculata Poelt & M. Steiner

Loc. 3, v.d. Boom 17777, on *Sorbus*: this species has already been mentioned from Switzerland by HINTEREGGER & al. (1989) and ROPIN & MAYRHOFER (1993) as *R. trevisanii* auct. For a detailed description see also HINTEREGGER (1994). It is most probably an overlooked species.

Rinodina rinodinoides (Anzi) H. Mayrhofer & Scheid.

= *R. serpentinei* H. Mayrhofer & Poelt

Loc. 6, *v.d. Boom 17759*, on schistose stones on a roof of barn in a meadow on a slope: this species is recognized by the *Milvina*-type ascospores and the lecideine subimmersed apothecia. It is a rather rare species and has previously been recorded in several localities in mountainous areas from northern to southern Europe, including two localities from the alpine belt in Spain (MIGULA, 1929; MAYRHOFFER & al., 1992; GIRALT & LLIMONA, 1997). It is recorded for the first time in Switzerland.

Strigula* aff. *phaea (Ach.) R. C. Harris

Loc. 16, *v.d. Boom 18126*, on an unidentified shrub: rather a small specimen, it belongs to *Strigula* aff. *phaea*, as it seems morphologically similar to *S. phaea* (pers. comm. E. Sérusiaux). However only microconidia have been found. Macroconidia are most important for identification of species in the genus. This complex includes circa 5 species and is presently under investigation by Emmanuël Sérusiaux. It is recorded for the first time in Switzerland.

Thelopsis flaveola Arnold

Loc. 16, *v.d. Boom 18131*, on an unidentified shrub among calcareous outcrop: an inconspicuous species, found growing among *Lecania cyrtella* (Ach.) Th. Fr. and *Biatora subduplex*. Only a few localities are known in Austria and Germany, where this species was found growing on *Rhododendron* as mentioned by HINTEREGGER (1994). It is recorded for the first time in Switzerland.

Thermutis velutina (Ach.) Flot.

Loc. 19, *v.d. Boom 18060*, on a vertical east side of a large boulder: accompanying species were *Agonimia tristicula* (Nyl.) Zahlbr. and *Leptogium lichenoides* (L.) Zahlbr.

Tremella coppinsii Diederich & Marson

Loc. 20, *v.d. Boom 17746*, on *Acer*; growing on *Platismatia glauca* (L.) W. L. Culb. & C. F. Culb.: this lichenicolous fungus was abundant in extensive patches of the host and has been found so far only on *P. glauca* (DIEDERICH, 1996; DIEDERICH & MARSON, 1988). It is recorded for the first time in Switzerland.

Tremella phaeophysciae Diederich & M. S. Christ.

Loc. 12, *v.d. Boom 18133*, on *Acer*, growing on *Phaeophyscia orbicularis* (Neck.) Moberg: this species, recently described in DIEDERICH (1996) and so far only known from one host (*P. orbicularis*) is widely distributed, mainly in Western Europe. It is recorded for the first time in Switzerland.

Verrucaria foveolata (Flörke) A. Massal.

Loc. 12, *v.d. Boom 17954*, on stones along rotting trunk on a slope near a waterfall.

Verrucaria umbrinula Nyl.

Loc. 7, v.d. Boom 17844, on a vertical and shaded granite in a *Picea* wood on a slope: it is characterized by the dark brown rather thin rimose and matt thallus together with the circa $17 \times 7 \mu\text{m}$ ascospores and the circa $150 \mu\text{m}$ diam. \pm globose perithecia. A species, reported from northern, north-western and central Europe. It has been mentioned as present in Switzerland by STI-ZENBERGER (1882-1883, as *V. subnigrescens* Nyl.).

Vouauxiella lichenicola (Linds.) Petr. & Syd.

Loc. 17, v.d. Boom 18028, on *Fraxinus*, on *Lecanora chlarotera* Nyl.: this lichenicolous fungus is common in many parts of Europe. Its distribution pattern in Switzerland is unknown to us.

Substrate abbreviations

Ab = <i>Abies</i>	Sa = <i>Salix</i>
Ac = <i>Acer</i>	So = <i>Sorbus</i>
Al = <i>Alnus</i>	c = calcareous rock
Be = <i>Betula</i>	s = siliceous rock
Fr = <i>Fraxinus</i>	m = moss
Ju = <i>Juniperus</i>	w = wood
La = <i>Larix</i>	t = terricolous
Pic = <i>Picea</i>	sh = unidentified shrub
Rh = <i>Rhododendron</i>	

Table 1. – List of specimens

The numbers indicate the collecting sites (see in material and methods)

Abrothallus bertianus s.l.	20A	Biatoropsis usnearum	3Pic
Abconditella lignicola	2w	Bryoria capillaris	3Pic
Acarospora badiofusca	6s	Buellia griseovirens	19Al
Acarospora fuscata	6s	Buellia punctata	4w
Acrocordia conoidea	18c	Buellia schaeferi	3Pic 7Pic 9La
Agonimia tristicula	13c 15c 19c	Calicium trabinellum	3Pic
Agonimia vouauxii	13t	Caloplaca aurea	15c
Arthonia mediella	1Ab	Caloplaca cerina	12Sa
Arthonia muscigena	3Sa	Caloplaca chlorina	12Ac 13c
Arthonia radiata	1Ab 7So 12So 17Fr	Caloplaca herbidella	19Al
Arthopyrenia punctiformis	3So 12w	Caloplaca sinapisperma	19c
Arthrorhaphis citrinella	5t 10c	Caloplaca sorocarpa	4sh
Aspicilia cf. rolleana	11s	Caloplaca stillicidiorum	9c
Bacidia absistens	19Al 20Ac	Candelariella aurella	15c
Bacidia arnoldiana	8c 20Ac	Candelariella vitellina	4w
Bacidia bagliettoana	13c	Candelariella xanthostigma	10Sa
Bacidia beckhausii	3Sa 10Sa 16sh	Carbonea supersparsa	7s
Bacidia rubella	17Ac	Carbonea vitellinaria	11s
“Bacidia” sabuletorum	12c 13c 14c 16c	Catapyrenium cinereum	5t 13t 15t
Bacidia subincompta	3Sa	Catapyrenium lachneum	15c
Bacidia vermifera	12A	Catillaria picila	16c
Baeomyces rufus	4t	Catinaria atropurpurea	2w
Biatora chrysantha	1Ab 10c	Cercidospora sp.	15c
Biatora flavopunctata	15Rh	Cetraria ericetorum	5t
Biatora helvola	2w 3Sa 20Ac	Cetraria nivalis	5t
Biatora ocelliformis	20Ac	Cetraria obtusata	5t
Biatora subduplex	16sh	Cetrelia cetrarioides	19Al
Biatora vernalis	9c	Chaenotheca chrysocephala	20Pic
Biatoridium delitescens	20Ac	Chaenotheca furfuracea	7s 10s 12w

Chaenotheca trichialis	.3Pic 12Pic 20Pic
Chaenothecopsis consociata	.3Pic
Chaenothecopsis pusilla	.12w
Chaenothecopsis viridireagens	.12w
Chromatochlamys muscorum	.7s 10c 19c
Cladonia phyllophora	.4t
Clauzadea monticola	.10s 12c
Collema auriforme	.8c
Collema cristatum	.13c
Collema multipartitum	.8c
Collema undulatum	.8c 15c
Cyphelium inquinans	.9La
Cyphelium tigillare	.14w
Cystocoleus ebeneus	.10s
Dacampia hookeri	.16c
Dermatocarpon intestiniforme	.9c
Dermatocarpon leptophyllum	.8c
Dermatocarpon miniatum	.9c 13c
Dibaeis baemyces	.5t 14t
Dimerella lutea	.20Ac
Dimerella pineti	.3w 20Ac
Diplotomma epipolium	.18c
Dirina massiliensis f. soled.	.18c
Eiglera homalomorpha	.12c
Endocarpon pallidum	.15c
Endocarpon pusillum	.9c 15c
Endococcus propinquus	.10s
Evernia divaricata	.3Pic
Farnoldia jurana	.16c
Farnoldia micropsis	.15c
Fulgensia fulgens	.14c
Fuscidea lightfootii	.19A 120Ac
Gyalecta jenensis	.12c
Hymenelia similis	.12c
Hypocnomyce scalaris	.9La
Hypotrachyna sinuosa	.20Ac
Icmadophila ericetorum	.12w
Imshaugia aleurites	.9La
Ionaspis melanocarpa	.12c
Lecania cyrtella	.16sh 17Ac
Lecania cf. cyrtellina	.10Sa 16sh
Lecania inundata	.8c 18c
Lecania naegeli	.17Fr
Lecania suavis	.18c
Lecania turicensis	.18c
Lecanora albella	.19A1
Lecanora cadubriae	.4sh
Lecanora circumborealis	.4w
Lecanora crenulata	.18c
Lecanora dispersa	.10s
Lecanora epanora	.7s
Lecanora handelii	.7s
Lecanora intricata	.7s
Lecanora intumescens	.17Fr
Lecanora jamesii	.3w
Lecanora piniperda	.4sh 5Rh 9La
Lecanora rupicola	.4w
Lecanora salicicola	.5Rh
Lecanora varia	.4w
"Lecidea" berengeriana	.18c
"Lecidea" hypnorum	.5Rh 13c 16c
"Lecidea" lurida	.8c 9c 13c 15c
"Lecidea" rufofusca	.5Rh
Lecidella elaeochroma	.1Ab 10Sa
Lecidella elaeochroma var. soralifera	.17Fr
Lecidella euphora	.9Ju 12Sa 16sh
Lecidella stigmataea	.10s
Lecidoma demissum	.11t
Lempholemma botryosum	.18c
Lepraria cf. cacuminum	.11s
Lepraria eburnea	.12Pic
Lepraria aff. flavescens	.8c
Lepraria nivalis	.12c
Lepraria rigidula	.20Ac
Leprocaulon microscopicu	.9s
Leptoplaca chrysodeta	.18c
Leptochidium albociliatum	.9c
Leptogium gelatinosum	.13c
Leptogium lichenoides	.19c
Leptogium plicatile	.8c
Leptogium saturninum	.12Ac 19Ac
Leptogium tenuissimum	.16c
Leptorhaphis epidermidis	.12Be
Letharia vulpina	.3La
Lichenocodium erodens	.9La
Lichinella nigritella	.8c
Lobaria linita	.10c
Lobaria pulmonaria (fer.)	.19Ac 20Ac
Lobothallia radiosa	.15c
Loxospora cisonica	.20Ac
Loxospora elatina	.20So Ac Pic
Megalaria pulverea	.20Ac
Melanelia disjuncta	.9s
Melanelia glabratula	.20Ac
Micarea botryoides	.3w
Micarea denigrata	.3w 4w 7w 9La
Micarea leprosula	.19c
Micarea lignaria var. lignaria	.12w
Micarea misella	.12Pic
Micarea prasina	.20Ac
Micarea sylvicola	.7s 10s
Miriquidica instrata	.11s
Miriquidica leucophaea	.7s
Mycobilimbia obscurata	.3Sa 16c
Neofuscelia pulla	.6s
Nephroma bellum	.20Ac
Nephroma parile	.9t 10s
Nephroma resupinatum	.3Sa 7So 20So
Normandina pulchella	.19A1
Ochrolechia alboflavescens	.3Pic 9La
Ochrolechia androgyna	.19A1
Opegrapha rufescens	.17Fr
Opegrapha vulgata	.20Ac
Pannaria pezizoides	.10c 20So
Parmelina pastillifera	.17Fa
Parmeliella triptophylla	.20Ac,So
Peltigera aptosa	.4t
Peltigera collina	.7s 20So
Peltigera didactyla	.9c
Peltigera horizontalis	.10c
Peltigera leucophlebia	.10c
Peltigera neckeri	.2w
Peltigera polydactyla	.20So
Peltigera praetextata	.3t 7t 20Ac
Pertusaria coccodes	.9La
Phaeophyscia endophoenicea	.17Fr
Phaeophyscia orbicularis	.12Ac
Phlyctis argena	.1Ab
Physcia albinea	.9s
Physcia dubia	.13c
Physconia distorta	.17Fa
Physconia muscigena	.15c
Physconia perisidiosa	.17Fa
Placidium lachneum	.15c
Placynthiella oligotropa	.5t
Placynthium filiforme	.14c
Placynthium nigrum	.14c
Placynthium tremniacum	.9c

Polyblastia ardesiaca	.12c	Thamnomia vermicularis	.14c
Polyblastia cupularis	.12c	Thelidium decipiens	.16c
Polychidium muscicola	.9c	Thelidium incavatum	.12c
Porina aenea	20Ac	Thelopsis flaveola	.16sh
Porina linearis	.18c	Thermutis velutina	.19c
Porpidia cf. crustulata	.10s	Thyrea confusa	.8c
Protoparmelia hypotremella	.19Al	Toninia alutacea	.14c
Psora decipiens	.14c 16c	Toninia aromatica	.14c
Psoroma hypnorum	.10c	Toninia candida	.9c
Ramalina pollinaria	.7s	Toninia diffracta	.14c
Rhizocarpon badioatrum	.11s	Toninia lobulata	.14c 16c
Rhizocarpon geminatum	.6s	Toninia rosulata	.13c
Rhizocarpon obscuratum	.11s	Toninia sedifolia	.9c
Rhizoplaca chrysoleuca	.6s	Trapeliopsis flexuosa	.4w
Rhizoplaca melanophthalma	.6s	Trapeliopsis granulosa	.11t
Rinodina conradii	.5t 5Rh	Tremella coppinsii	.20Ac
Rinodina exigua	.2Sa 3Sa 9La 10Sa	Tremella phaeophysciae	.12Ac
Rinodina griseosoralifera	.10Sa 19Ac	Umbilicaria hirsuta	.6s
Rinodina milvina	.9s	Usnea cavernosa	.3Pic 7Pic
Rinodina mniaraea	.5t	Usnea scabrata s.l.	.3Pic 7Pic
Rinodina orculata	.3So	Usnea substerilis	.3Pic 7Pic
Rinodina parasitica	.9s	Verrucaria coerulea	.18c
Rinodina septentrionalis	.4w	Verrucaria foveolata	.12c
Rinodina rinodinoides	.6s	Verrucaria lecideoides	.15c
Sarea difformis	.1Ab	Verrucaria umbrinula	.7s
Solorina crocea	.5t	Vouauxiella lichenicola	.17Fr
Solorina spongiosa	.3t	Xanthoria candelaria	.4w
Squamarina cartilaginea	.14c	Xanthoria elegans	.15c
Squamarina gypsacea	.14c	Xanthoria fulva	.12Ac
Staurothele areolata	.13c 15c	Xanthoria sorediata	.13c
Steinia geophana	.14t	Xylographa vitiligo	.3w
Strigula aff. phaea	.16sh		

ACKNOWLEDGEMENTS

We are grateful to the following specialists for their help in the identification of some of the specimens: Mr. Maarten Brand (*Bacidia*, *Farnoldia*, *Lempholemma*), Dr. Othmar Breuss (*Dermatocarpon*, *Placidium*, *Verrucaria*), Dr. Brian Coppins (*Arthonia*, *Caloplaca*), Dr. Paul Diederich (lichenicolous fungi), Dr. Isabel Martinez and Dr. Orvo Vitikainen (*Peltigera*), Dr. Helmut Mayrhofer (*Chromatochlamys*, *Rinodina*), Dr. Christian Printzen (*Biatora*), Dr. Emmanuël Séruiaux (*Strigula*) and Dr. Leif Tibell (*Chaenothecopsis*).

REFERENCES

- APTROOT, A., P. DIEDERICH, C. M. VAN HERK, L. SPIER & V. WIRTH (1997). *Protoparmelia hypotremella*, a new sterile corticolous species from Europe and its lichenicolous fungi. *Lichenologist* 29: 415-424.
- BOISSIÈRE, J. C., S. DÉRUELLE & C. ROUX (1989). Liste provisoire des lichens récoltés dans la région de Chamonix (excursion A. F. L., 1988). *Bull. Assoc. Franç. Lichénol.* 14: 5-18.
- BOOM, P. P. G. VAN DEN & H. J. M. SIPMAN (1994). *Cetraria obtusata* comb. et. stat. nov., an overlooked lichen species from the central Alps. *Lichenologist* 26: 105-112.
- BOOM, P. P. G. VAN DEN, A. M. BRAND & H. J. M. SIPMAN (1993). Lichens from Evolène and surroundings (Switzerland). *Meylania* 3: 12-26.
- BOOM, P. P. G. VAN DEN, O. BREUSS, L. SPIER & A. M. BRAND (1996). Beitrag zur Flechtenflora Kärntens, Ergebnisse der Feldtagung der Bryologischen und Lichenologischen Arbeitsgruppe der KNNV in Weissbriach 1994. *Linzener Biol. Beitr.* 28: 619-654.
- BRAND, A. M., A. APTROOT, A. J. DE BAKKER & P. P. G. VAN DOBBEN (1988). *Standaardlijst van de nederlandse sekorstmossen. Checklist of the lichens of the Netherlands*. Stichting Uitgeverij Koninklijke Nederlandse Natuurhistorische Vereniging, Utrecht, 68 pp.
- CLAUZADE, G., P. DIEDERICH & C. ROUX (1989). Nelikenigintaj fungoj likenlogaj. Ilustrita determinlibro. *Bull. Soc. Linn. Provence*, Num. Spéc. 1: 1-142.
- CLERC, P., C. SCHEIDEGGER & K. AMMANN (1992). Liste rouge des macrolichens de la Suisse. *Bot. Helv.* 102: 71-83.

- CULBERSON, C. & K. AMMANN (1979). Standardmethode zur Dünnschichtchromatographie von Flechtensubstanzen. *Herzogia* 5: 1-24.
- DIEDERICH, P. (1996). The Lichenicolous Heterobasidiomycetes. *Biblioth. Lichenol.* 61: 1-198.
- DIEDERICH, P. & G. MARSON (1988). *Tremella coppinsii*, a new lichenicolous Basidiomycete from Sarawak. *Notes Roy. Bot. Gard. Edinburgh* 45: 145-176.
- DIETRICH, M. & C. SCHEIDEGGER (1996). The importance of sorediate crustose lichens in the epiphytic lichen flora of the Swiss Plateau and the Pre-Alps. *Lichenologist* 28: 245-256.
- DIETRICH, M. & C. SCHEIDEGGER (1997). Frequency, diversity and ecological strategies of epiphytic lichens in the swiss central plateau and the Pre-Alps. *Lichenologist* 29: 237-258.
- DIETRICH, M., C. KELLER & C. SCHEIDEGGER (1992). Botanischer Reichtum am Weg von Davos über die Berggüter Furgga zum Albula: Sommerexkursion 1991 im Anklang an die erste Exkursion der Schweizerischen Botanischen Gesellschaft 1890. 5. Flechten. *Bot. Helv.* 102: 41-47.
- EKMAN, S. (1996). The corticolous and lignicolous species of *Bacidia* and *Bacidina* in North America. *Opera Bot.* 127: 1-148.
- ETAYO, J., B. AGUIRRE & P. DIEDERICH (1993). Interesting or new lichens from the Atlantic Pyrenees and the north of the Iberian Peninsula. II. *Nova Hedwigia* 57: 179-194.
- FREY, E. (1923). Flechten der Schweiz. *Ber. Schweiz. Bot. Ges.* 32: 52-65.
- FREY, E. (1937). Die Flechtenvegetation des Aletschreservates und seiner näheren Umgebung. *Bull. Murith. Soc. Valais. Sci. Nat.* 54: 55-93.
- GIRALT, M. & X. LLIMONA (1997). The saxicolous species of the genera *Rinodina* and *Rinodinella* lacking spot test reactions in the Iberian Peninsula. *Mycotaxon* 62: 175-224.
- GYELNIK, V. (1940). Lichinaceae, Heppiaceae, Pannariaceae. In: RABENHORST, L., *Kryptogamenflora*, ed. 2, 9(1/2). 272 pp.
- HAFELLNER, J. (1994). On *Biatoridium*, a resurrected genus of lichenized fungi (Ascomycotina, Lecanorales). *Acta Bot. Fenn.* 150: 39-46.
- HANKO, B. (1983). Die Chemotypen der Flechtengattung *Pertusaria* in Europa. *Biblioth. Lichenol.* 19: 1-297.
- HINTEREGGER, E. (1994). Krustenflechten auf den Rhododendron-Arten (*Rh. ferrugineum* und *Rh. hirsutum*) der Ostalpen. *Biblioth. Lichenol.* 55: 1-346.
- HINTEREGGER, E., H. MAYRHOFER & J. POELT (1989). Die Flechten der Alpenrosen in den Ostalpen (*Rhododendron ferrugineum* und *Rh. hirsutum*). I. Einige Arten der Gattungen *Lecanora* und *Rinodina*. *Mitt. Naturwiss. Vereins Steiermark* 119: 83-102.
- KALB, K. (1970). Flechtengesellschaften der vorderen Ötztaler Alpen. *Diss. Bot.* 9: 1-118.
- LESDAIN, B. M. de (1910a). *Recherches sur les lichens des environs de Dunkerque*. Imprim. P. Michel, Dunkerque, 301 pp.
- LESDAIN, B. M. de (1914). *Recherches sur les lichens des environs de Dunkerque*, 1er supplément. Société dunkerquoise pour l'encouragement des sciences, des lettres et des arts, Dunkerque, 190 pp.
- LETTAU, G. (1919). Schweizer Flechten. II. *Nova Hedwigia* 60: 267-312.
- LETTAU, G. (1942). Flechten aus Mitteleuropa. VII. *Feddes Repert. Spec. Nov. Regni Veg. Beih.* 119: 265-348.
- LETTAU, G. (1954). Flechten aus Mitteleuropa. IX. *Feddes Repert. Spec. Nov. Regni Veg. Beih.* 56: 172-278.
- MAYRHOFER, H., C. SCHEIDEGGER & J. W. SHEARD (1992). On the taxonomy of five saxicolous species of the genus *Rinodina*. *Nord. J. Bot.* 12: 451-459.
- MEYLAN, C. (1922). Contribution à la connaissance des lichens du Jura. *Bull. Soc. Vaud. Sci. Nat.* 54: 287-294
- MIGULA, W. (1929). *Kryptogamen-Flora von Deutschland, Österreich und der Schweiz*. Band IV. Flechten 1. Teil: 1-527.
- MIGULA, W. (1931). *Kryptogamen-Flora von Deutschland, Österreich und der Schweiz*. Band IV. Flechten 2. Teil: 1-868.
- NIMIS, P. L. (1993). *The lichens of Italy*. Museo Regionale di Scienze Naturali, Torino, 897 pp.
- ORANGE, A. (1998). *Dermatocarpon leptophyllodes* and related species in the British Isles. *Lichenologist* 30: 1-20.
- POELT, J. (1965). Über einige Artengruppen der Flechtengattungen *Caloplaca* und *Fulgensia*. *Mitt. Bot. Staatssamml. München* 5: 571-607.
- PRINTZEN, C. (1995). Die Flechtengattung *Biatora* in Europa. *Biblioth. Lichenol.* 60: 1-275.
- PRINTZEN, C. & Z. PALICE (1999). The distribution, ecology and conservational status of the lichen genus *Biatora* in central Europe. *Lichenologist* 31: 319-335.

- PURVIS, O. W., B. J. COPPINS, D. L. HAWKSWORTH, P. W. JAMES & D. M. MOORE (1992). *The lichen flora of Great Britain and Ireland*. London: Natural History Museum Publications, 710 pp.
- ROPIN, K. & H. MAYRHOFER (1993). Zur Kenntnis corticoler Arten der Gattung *Rinodina* (lichenisierte Ascomyceten) in den Ostalpen und angrenzenden Gebieten. *Herzogia* 9: 779-835.
- RUOSS, E. (1992). *Flechten im Kanton Luzern. Untersuchungen zur Bioindikation und Floristik, sowie zur Immissionsökologie voralpiner Hochmoore*. Veröffentlichungen aus dem Natur-Museum Luzern, 3, 98 pp.
- SANTESSON, R. (1993) *The lichens and lichenicolous fungi of Sweden and Norway*. Lund, 240 pp.
- STIZENBERGER, E. (1882-1883). Lichenes helvetici eorumque stationes et distributio. *Ber. Tätigk. St. Gallischen Naturwiss. Ges.* 22: 255-522 (1882); 23: 201-327 (1883).
- TÜRK, R. & J. POELT (1993). *Bibliographie der Flechten und flechtenbewohnenden Pilze in Österreich*. Biosystematics and Ecology Series No. 3. Österr. Akademie der Wissenschaften, 168 pp.
- WESTBERG, M. & I. KÄRNEFELT (1998). The genus *Fulgensia* A. Massal. & De Not., a diverse group in the Teloschistaceae. *Lichenologist* 30: 515-532.
- WIRTH, V. (1995a). *Flechtenflora* (2). Stuttgart: Ulmer, 661 pp.
- WIRTH, V. (1995b). *Die Flechten Baden-Württembergs*, Teil I und II. Stuttgart, Ulmer. 1006 pp.
- ZSCHACKE, H. (1926). Die Flechten des Davoser Tales. *Mitth. Naturf. Ges. Davos* 1925/26: 1-59.
- ZSCHACKE, H. (1934). Epigloaceae, Verrucariaceae und Dermatocarpaceae. *In*: RABENHORST, L., *Kryptogamenflora* ed. 2, 9(1/1): 44-695.

Addresses of the authors: PPGvdB: Arafura 16, 5691 JA, Son, The Netherlands.

PC: Conservatoire et Jardin botaniques de la Ville de Genève, Case Postale 60, CH-1292 Chambésy/GE, Suisse.

