

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 61 (1986)  
  
**Artikel:** Cyclic homology of groups and the Bass conjecture.  
**Autor:** Eckmann, Bruno  
**DOI:** <https://doi.org/10.5169/seals-46927>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Cyclic homology of groups and the Bass conjecture

BENO ECKMANN

### 0. Introduction

**0.1.** The cyclic homology  $HC_i(\mathbb{Q}G)$  of a group algebra  $\mathbb{Q}G$  decomposes into a direct sum indexed by the conjugacy classes  $[x]$  in  $G$ , as shown by Burghlea [7] (see Section 1.3 below). We will consider certain classes of groups of finite homology dimension over  $\mathbb{Q}$ ,  $hd_{\mathbb{Q}}G = n$ , and show that for  $i \geq n$  the terms in  $HC_i(\mathbb{Q}G)$  corresponding to conjugacy classes  $[x]$  of elements of infinite order vanish. The groups  $G$  with  $hd_{\mathbb{Q}}G = n < \infty$  for which this will be done are

- (a) Nilpotent groups  $G$ ,
- (b) Torsion-free solvable groups,
- (c) Linear groups  $G \subset GL_r(F)$  where  $F$  is a field of characteristic 0,
- (d) Groups of cohomology dimension  $cd_{\mathbb{Q}}G \leq 2$  (here  $n \leq 2$ ).

We recall (Serre [10]) that if in (c)  $F$  is a number field and  $G$  finitely generated, then  $G$  is always of finite virtual cohomology dimension, whence  $hd_{\mathbb{Q}}G = n < \infty$ . The case (b) actually falls under (c), but we prefer to give a simple direct argument, cf. Remark 2.3' below.

As an immediate consequence of that vanishing result it follows that the character maps from  $K$ -theory of  $\mathbb{Q}G$  to  $HC_*(\mathbb{Q}G)$ , see Karoubi [8], have vanishing components in the summands indexed by  $[x]$  with  $x$  of infinite order—for all the groups listed above.

In particular, the character map  $Ch_0^0: K_0(\mathbb{Q}G) \rightarrow HC_0(\mathbb{Q}G)$  can easily be seen to be the “Hattori–Stallings rank”  $r_P$  of finitely generated projective  $\mathbb{Q}G$ -modules  $P$  (representing elements of  $K_0(\mathbb{Q}G)$ ), see Section 3.2. For the groups above it thus follows that  $r_P$  is concentrated on the conjugacy classes  $[x]$  of elements  $x$  of finite order; hence on [1] if  $G$  is torsion-free. This is a contribution towards the strong Bass conjecture [3, p. 156]. Note that the case (c) yields a weaker statement than Bass’ result on linear groups [3, p. 156/57]; but our method is entirely different and stems from a result more general in another direction. The result establishing the Bass conjecture over  $\mathbb{Q}G$  in the other cases seems to be new.

## 1. Cyclic homology of groups

**1.1.** Let  $G$  be a group,  $\mathbb{Q}G$  its rational group algebra, and  $HC_i(\mathbb{Q}G)$ ,  $i \in \mathbb{Z}$ , the cyclic homology of  $\mathbb{Q}G$  in the sense of Connes; we will call it here in short the cyclic homology of  $G$ . It is related to the Hochschild homology  $HH_i(\mathbb{Q}G)$  of  $\mathbb{Q}G$ , with bimodule-coefficients in  $\mathbb{Q}G$  by left and right multiplication, through the “Connes–Gysin exact sequence”

$$\cdots \rightarrow HH_i(\mathbb{Q}G) \rightarrow HC_i(\mathbb{Q}G) \xrightarrow{S} HC_{i-2}(\mathbb{Q}G) \rightarrow HH_{i-1}(\mathbb{Q}G) \rightarrow \cdots \quad (1.1)$$

It is a standard fact (see [9]) that Hochschild homology of  $\mathbb{Q}G$  with bimodule coefficients can be expressed as homology of  $G$  with the same coefficient module turned into a right  $G$ -module; in the present case this is  $\mathbb{Q}G$  with  $G$ -action by conjugation in  $G$ . It thus follows that for groups  $G$  of finite homology dimension  $hd_{\mathbb{Q}}G = n < \infty$  over  $\mathbb{Q}$  (i.e., for all  $\mathbb{Q}G$ -module coefficients) the cyclic homology of  $G$  stabilized above  $n$ :

$$\begin{aligned} HC_{n+2k}(\mathbb{Q}G) &= HC_n(\mathbb{Q}G), \\ HC_{n+2k+1}(\mathbb{Q}G) &= HC_{n+1}(\mathbb{Q}G) \end{aligned}$$

for  $k = 0, 1, 2, \dots$

**1.2.** The conjugation module  $\mathbb{Q}G$  obviously decomposes into a direct sum of right  $\mathbb{Q}G$ -module indexed by the conjugacy classes  $[x]$  of  $G$ ;  $x$  is an arbitrary but fixed representative of  $[x]$ :

$$\mathbb{Q}G = \bigoplus_{[x]} D_x$$

where  $D_x$  is the  $\mathbb{Q}$ -module over the elements  $x' \in [x]$  as basis, and with  $\mathbb{Q}G$ -action by  $x' \mapsto y^{-1}x'y$ ,  $y \in G$ . If  $C_x$  is the centralizer of  $x$  in  $G$ ,  $D_x$  is isomorphic to  $\mathbb{Q}(G/C_x)$ , the right  $\mathbb{Q}G$ -module generated by the right cosets modulo  $C_x$ ; the isomorphism is given by  $x' \mapsto C_x z$  where  $z \in G$  is such that  $z^{-1}xz = x'$ . Thus

$$\mathbb{Q}G = \bigoplus_{[x]} \mathbb{Q}(G/C_x) = \bigoplus_{[x]} (\mathbb{Q} \otimes_{C_x} \mathbb{Q}G),$$

and finally

$$HH_i(\mathbb{Q}G) = \bigoplus_{[x]} H_i(G; \mathbb{Q} \otimes_{C_x} \mathbb{Q}G) = \bigoplus_{[x]} H_i(C_x; \mathbb{Q}) \quad (1.2)$$

with trivial  $G$ -module coefficients  $\mathbb{Q}$ .

*Remark 1.1.*  $HH_0(\mathbb{Q}G)$  is the  $\mathbb{Q}$ -module having the conjugacy classes  $[x]$  in  $G$  as basis. This can also be seen directly from the well-known fact that  $HH_0(\mathbb{Q}G)$  is  $\mathbb{Q}G/\{\lambda\mu - \mu\lambda\}$ , where  $\{\lambda\mu - \mu\lambda\}$  denotes the  $\mathbb{Q}$ -submodule generated by all  $\lambda\mu - \mu\lambda$ ,  $\lambda, \mu \in \mathbb{Q}G$ ; i.e., the  $\mathbb{Q}$ -submodule generated by all  $xy - yx$ ,  $x, y \in G$ . We write  $\overline{\mathbb{Q}G}$  for this factor- $\mathbb{Q}$ -module of  $\mathbb{Q}G$ ,  $T: \mathbb{Q}G \rightarrow \overline{\mathbb{Q}G}$  for the canonical map, with  $T(\lambda\mu) = T(\mu\lambda)$ ,  $\lambda, \mu \in \mathbb{Q}G$ .

**1.3.** A direct sum decomposition of  $HC_i(\mathbb{Q}G)$ , with terms indexed by the conjugacy classes  $[x]$  in  $G$ , has been given by Burghlea [7] using topological (simplicial) constructions:

$$HC_i(\mathbb{Q}G) = \bigoplus'_{[x]} [H_*(C_x/\langle x \rangle; \mathbb{Q}) \otimes H_*(\mathbb{C}P^\infty; \mathbb{Q})]_i \oplus \bigoplus''_{[x]} H_i(C_x/\langle x \rangle; \mathbb{Q}). \quad (1.3)$$

Here  $\langle x \rangle$  denotes the cyclic subgroup of  $G$  generated by  $x$ , and  $\bigoplus'$  is summation over all  $[x]$  with finite  $\langle x \rangle$ ,  $\bigoplus''$  over all  $[x]$  with infinite  $\langle x \rangle$ .

The methods of [7] also yield the  $\bigoplus$ -decomposition (1.2) of  $HH_i(\mathbb{Q}G)$  and shows that the Connes–Gysin sequence (1.1) decomposes into exact sequences of the same type, one for each  $[x]$  in  $G$ .

We will consider in Section 2 groups of finite dimension  $hd_{\mathbb{Q}}G = n < \infty$  and show that, for certain classes of such groups, one has  $\bigoplus''_{[x]} = 0$  in the stable value

$HC_n(\mathbb{Q}G)$  and  $HC_{n+1}(\mathbb{Q}G)$ ; in other words, *the stable value is concentrated on the conjugacy classes  $[x]$  of elements  $x$  of finite order* – hence for torsion-free groups on the conjugacy class  $[1]$ . This will be done for the classes (a)–(d) listed in the introduction. Immediate consequences (Section 3) concern the character maps from  $K$ -theory of  $\mathbb{Q}G$  to cyclic homology and, in particular, the Hattori–Stallings rank as mentioned in the introduction.

## 2. Groups of finite dimension

**2.1.** “Finite dimension” for groups  $G$  will refer here, unless otherwise specified, to the homology dimension  $hd_{\mathbb{Q}}G$  over  $\mathbb{Q}$ , i.e., with respect to



$\mathbb{Q}G$ -module coefficients. For any subgroup  $S \subset G$ , in particular for the centralizers  $C_x$ , we have  $hd_{\mathbb{Q}}S \leq hd_{\mathbb{Q}}G = n < \infty$ .

In our context we are thus interested in the homology of factor groups  $G/\langle x \rangle$  where  $hd_{\mathbb{Q}}G = n < \infty$  and  $x$  is a central element; actually in homology with trivial  $\mathbb{Q}$ -coefficients only, and in its vanishing above  $n$ . In other words, we are looking at  $thd_{\mathbb{Q}}G/\langle x \rangle$ , the *trivial homology dimension over  $\mathbb{Q}$* ; i.e., defined exactly as  $hd_{\mathbb{Q}}$  but referring to trivial  $\mathbb{Q}$ -module coefficients only. One always has  $thd_{\mathbb{Q}} \leq hd_{\mathbb{Q}}$ . For that type of dimension we recall the following very simple but useful sum formula (Bieri [6]):

**LEMMA 2.1.** *Let  $U$  be a central subgroup of the group  $V$ , and  $W = V/U$ . If both  $thd_{\mathbb{Q}}U$  and  $thd_{\mathbb{Q}}W$  are finite then*

$$thd_{\mathbb{Q}}V = thd_{\mathbb{Q}}U + thd_{\mathbb{Q}}W.$$

A further preliminary remark concerns the case where  $\langle x \rangle$  is finite: then the spectral sequence

$$H_i(G/\langle x \rangle; H_j(\langle x \rangle; \mathbb{Q})) \Rightarrow H_{i+j}(G; \mathbb{Q})$$

shows that  $thd_{\mathbb{Q}}G/\langle x \rangle = thd_{\mathbb{Q}}G$ , hence  $\leq n$ . In all what follows we therefore restrict attention to central elements  $x$  of *infinite* order. In that case the spectral sequence does *not* imply that  $thd_{\mathbb{Q}}G/\langle x \rangle$  is finite; however, if it *is* finite then the sum formula yields

$$thd_{\mathbb{Q}}G/\langle x \rangle \leq n - 1.$$

**2.2. Nilpotent groups.** We recall (Stammbach [11]) that if  $G$  is nilpotent then  $hd_{\mathbb{Q}}G$  is equal to the Hirsch number  $hG$  (the sum of the torsion-free ranks of the factors of any normal series of  $G$  with Abelian factors); this holds, more generally, for solvable groups. We thus assume  $hG = n < \infty$ .

Let  $x \in G$  be a central element of infinite order,  $S$  a finitely generated subgroup of  $G/\langle x \rangle$ , and  $T$  the preimage of  $S$  in  $G$ ,  $T/\langle x \rangle = S$ . Since  $S$  is finitely generated nilpotent it is polycyclic, and therefore  $hd_{\mathbb{Q}}S = hs$  is finite (equal to the number of infinite cyclic factors in a normal series with cyclic factors). The sum formula now yields

$$thd_{\mathbb{Q}}S = thd_{\mathbb{Q}}T - 1 \leq n - 1.$$

$G/\langle x \rangle$  is the direct limit of its finitely generated subgroups  $S$ ; and since homology commutes with direct limits it follows that  $thd_{\mathbb{Q}}G/\langle x \rangle$  is  $\leq n - 1$ :

**THEOREM 2.2.** *Let  $G$  be a nilpotent group of finite dimension  $hd_{\mathbb{Q}}G = n$ . Then one has for any central element  $x \in G$  of infinite order*

$$H_i(G/\langle x \rangle; \mathbb{Q}) = 0 \quad \text{for } i \geq n.$$

**2.3. Torsion-free solvable groups.** Let  $G$  be torsion-free solvable with  $hd_{\mathbb{Q}}G = hG = n < \infty$  (or equivalently, solvable with  $hd_{\mathbb{Z}}G < \infty$ ). We consider the Hirsch–Plotkin radical  $R$  of  $G$ , i.e., the maximal locally nilpotent normal subgroup of  $G$ . For any Abelian subgroup  $S$  of  $G$  the torsion-free rank  $hS = hd_{\mathbb{Q}}S$  is  $\leq n$ . As  $G$  is torsion-free (actually a weaker condition would do) we can apply a theorem of Baer–Heineken [2] which tells that

( $\alpha$ )  $R$  is nilpotent

( $\beta$ )  $G/R$  is finitely generated

( $\gamma$ )  $G/R$  contains an Abelian subgroup  $A$  of finite index.

From (b) and (c) we infer that  $hd_{\mathbb{Q}}G/R = hd_{\mathbb{Q}}A$  is finite, say  $= m$ . If  $x \in G$  is central it must lie in  $R$ , and if it is of infinite order Theorem 2.2 tells that  $thd_{\mathbb{Q}}R/\langle x \rangle = thd_{\mathbb{Q}}R - 1 \leq n - 1$ . From the spectral sequence for  $G/R \cong G/\langle x \rangle / R/\langle x \rangle$ ,

$$H_i(G/R; H_j(R/\langle x \rangle; \mathbb{Q})) \Rightarrow H_{i+j}(G/\langle x \rangle; \mathbb{Q})$$

we see that  $H_k(G/\langle x \rangle; \mathbb{Q}) = 0$  for  $k > m + n - 1$ ; i.e.,  $thd_{\mathbb{Q}}G/\langle x \rangle$  is finite and hence  $\leq n - 1$ .

**THEOREM 2.3.** *Let  $G$  be a torsion-free solvable group of finite dimension  $hd_{\mathbb{Q}}G = n < \infty$ . Then one has for any central element  $x \in G$  of infinite order*

$$H_i(G/\langle x \rangle; \mathbb{Q}) = 0 \quad \text{for } i \geq n.$$

*Remark 2.3'.* The groups  $G$  above admit faithful linear representations over  $\mathbb{Q}$  (cf. [13], p. 25) and thus are included in 2.4 below. However, the proof of the linear embedding starts precisely from the structure properties ( $\alpha$ ), ( $\beta$ ), ( $\gamma$ ); thus the simple direct argument seems preferable.

**2.4. Linear groups.** We now consider a linear group  $G \subset GL_r(F)$ , where  $F$  is a field of characteristic 0, with  $hd_{\mathbb{Q}}G = n < \infty$ . Let  $Z$  be the center of  $G$ ,  $\pi: G \rightarrow G/Z$  the canonical map. Since  $Z$  is closed in the Zariski topology,  $G/Z$  is again a linear group over the same field  $F$ . We are going to apply the R. Alperin–Shalen criterion [1] to finitely generated subgroups  $S$  of  $G/Z$ , in order to prove that they are virtually of finite cohomology dimension.

For this let first  $U$  be a finitely generated unipotent subgroup of  $G/Z$  and put  $\pi^{-1}U = V$ ,  $V/Z = U$ . Then  $U$  being torsion-free finitely generated nilpotent (polycyclic),  $hd_{\mathbb{Q}}U = thd_{\mathbb{Q}}U = hU$  (Hirsch number, see 2.2). By Lemma 2.1,

$$thd_{\mathbb{Q}}Z + thd_{\mathbb{Q}}U = thd_{\mathbb{Q}}V.$$

Now  $hd_{\mathbb{Q}}Z \leq n$  and  $hd_{\mathbb{Q}}V \leq n$ ; let  $m = thd_{\mathbb{Q}}Z$ . Thus  $thd_{\mathbb{Q}}U \leq n - m$ ; i.e., we get a uniform bound for all finitely generated unipotent subgroups of  $G/Z$ , the Hirsch numbers  $hU$  being  $\leq n - m$ .

If  $S$  is any finitely generated subgroup of  $G/Z$ , it follows by [1] that its virtual cohomology dimension, and hence  $hd_{\mathbb{Q}}S$ , is finite. Putting  $\pi^{-1}S = T \subset G$ ,  $T/Z = S$ , Lemma 2.1 tells that

$$thd_{\mathbb{Q}}Z + thd_{\mathbb{Q}}S = thd_{\mathbb{Q}}T \leq n,$$

and thus  $thd_{\mathbb{Q}}S \leq n - m$ . The direct limit argument then yields  $thd_{\mathbb{Q}}G/Z \leq n - m$ .

**THEOREM 2.4.** *Let  $G$  be a linear group of finite dimension  $hd_{\mathbb{Q}}G = n$ , over a field of characteristic 0,  $Z$  is center and  $thd_{\mathbb{Q}}Z = m$ . Then*

$$H_i(G/Z; \mathbb{Q}) = 0 \quad \text{for } i > n - m.$$

We are looking for a similar result concerning  $G/\langle x \rangle$  where  $x$  is a central element of infinite order. Since  $G/Z = G/\langle x \rangle / Z/\langle x \rangle$  we can apply the spectral sequence (with trivial  $\mathbb{Q}$ -module coefficients) together with Theorem 2.2 on  $Z/\langle x \rangle$ . This immediately yields  $H_i(G/\langle x \rangle; \mathbb{Q}) = 0$  for  $i > (n - m) + (m - 1)$ :

**THEOREM 2.4'.** *Let  $G$  be as in Theorem 2.4. and  $x$  a central element of  $G$  of infinite order. Then*

$$H_i(G/\langle x \rangle; \mathbb{Q}) = 0 \quad \text{for } i \geq n.$$

**2.5. Groups of cohomology dimension  $\leq 2$ .** We write as usual  $cd_{\mathbb{Q}}G$  for the cohomology dimension of the group  $G$  over  $\mathbb{Q}$ ; i.e., with respect to all  $\mathbb{Q}G$ -module coefficients. The assumption  $cd_{\mathbb{Q}}G \leq 2$  includes all groups which are virtually of cohomology dimension 2 (over  $\mathbb{Z}$ ), but is more general; it of course implies  $hd_{\mathbb{Q}}G \leq 2$ .

The case  $cd_{\mathbb{Q}} \leq 1$  is easily dealt with: it means that  $G$  contains a free (normal) subgroup  $F$  of finite index. If  $x$  is a central element of infinite order then  $\langle x \rangle \cap F \neq 1$ ; hence  $F$  having non-trivial center must be cyclic  $= \langle c \rangle$ . Then both

$G/\langle x, F \rangle$  and  $\langle x, F \rangle/\langle x \rangle \cong F/F \cap \langle x \rangle$  are finite, and so is  $G/\langle x \rangle$ . Thus  $hd_{\mathbb{Q}} G/\langle x \rangle = 0$ . We thus restrict attention to the case  $cd_{\mathbb{Q}} G = 2$ .

**THEOREM 2.5.** *Let  $G$  be a group with  $cd_{\mathbb{Q}} G = 2$ , and  $x$  a central element of infinite order. Then*

$$H_i(G/\langle x \rangle; \mathbb{Q}) = 0 \quad \text{for } i \geq 2.$$

This is a consequence of various known facts concerning group (co-) homology, finiteness properties, and structure theorems (cf. Bieri [4], Corollary 8.7, and [5]). We give a short outline of the proof for our somewhat more special situation. It suffices to prove the claim for finitely generated  $G$ : If the subgroup  $S \subset G/\langle x \rangle$  is finitely generated so is its preimage  $T \subset G$ , and from the result for  $S = T/\langle x \rangle$  the direct limit argument yields the claim for  $G/\langle x \rangle$ .

As a first step one shows that  $G$  is of type  $FP_{\mathbb{Q}}$ ; i.e., that there exists a finitely generated projective resolution over  $\mathbb{Q}G$  (of length 2 since  $cd_{\mathbb{Q}} G = 2$ )

$$0 \rightarrow P_2 \rightarrow P_1 \rightarrow P_0 \rightarrow \mathbb{Q} \rightarrow 0.$$

To prove this we use R. Strebel's finiteness criterion [11]:  $G$  is of type  $FP_{\mathbb{Q}}$  if and only if  $cd_{\mathbb{Q}} G$  is finite and the canonical map  $H^i(G; \bigoplus \mathbb{Q}G) \rightarrow \bigoplus H^i(G; \mathbb{Q}G)$  is an isomorphism for all  $i$  and all direct sums  $\bigoplus$ . In our case the spectral sequence for  $G/\langle x \rangle$  yields

$$H^1(G; \mathbb{Q}G) \cong H^0(G/\langle x \rangle; H^1(\langle x \rangle; \mathbb{Q}G))$$

and

$$H^2(G; \mathbb{Q}G) \cong H^1(G/\langle x \rangle; H^1(\langle x \rangle; \mathbb{Q}G))$$

(the action of  $G$  on  $H^1(\langle x \rangle; \mathbb{Q}G)$  is induced by the trivial conjugation of  $G$  on  $\langle x \rangle$ ). For the infinite cyclic group  $\langle x \rangle$  one has  $H^i(\langle x \rangle; \mathbb{Q}\langle x \rangle) = 0$  for  $i \neq 1$ , and  $H^1(\langle x \rangle; \mathbb{Q}\langle x \rangle) = \mathbb{Q}$  with trivial action; and  $H^1(\langle x \rangle; \mathbb{Q}G) = 0$  for  $i \neq 1$ ,  $H^1(\langle x \rangle; \mathbb{Q}G) = \mathbb{Q} \otimes_{\mathbb{Q}\langle x \rangle} \mathbb{Q}G \cong \mathbb{Q}(G/\langle x \rangle)$ .

For  $A = \bigoplus \mathbb{Q}G$  we get  $H^1(\langle x \rangle; A) = \bigoplus H^1(\langle x \rangle; \mathbb{Q}G)$ , and since  $G/\langle x \rangle$  is finitely generated, we see that  $G$  fulfills the Strebel criterion for  $i = 1$  and 2; in dimensions  $i \neq 1, 2$  this is trivially the case since  $H^i(G; \bigoplus \mathbb{Q}G) = 0 = \bigoplus H^i(G; \mathbb{Q}G)$ . Thus  $G$  is of type  $FP_{\mathbb{Q}}$ .

As a second step one draws more conclusions from the above formula for  $H^2(G; \mathbb{Q}G)$ . We note that

$$H^2(G; \mathbb{Q}G) = H^1(G/\langle x \rangle; \mathbb{Q}(G/\langle x \rangle)).$$

As  $G$  is of type  $FP_{\mathbb{Q}}$  with  $cdG = 2$ ,  $H^2(G; \mathbb{Q}G)$  is  $\neq 0$  and finitely generated as a right  $\mathbb{Q}G$ -module, and so is  $H^1(G/\langle x \rangle; \mathbb{Q}(G/\langle x \rangle))$  over  $\mathbb{Q}(G/\langle x \rangle)$ . This implies that  $G/\langle x \rangle$  has more than one end and is *accessible*; in other words,  $G/\langle x \rangle$  is the fundamental group of a finite graph of groups with finite edge groups, and with vertex groups  $V$  satisfying  $H^1(V; \mathbb{Q}V) = 0$  (1 or 0 ends).  $V$  is finitely generated, and so is its preimage  $W$  in  $G$ ,  $W/\langle x \rangle = V$ . As before we get  $H^2(W; \mathbb{Q}W) = H^1(V; \mathbb{Q}V)$ ; but now this is  $= 0$ , whence  $cd_{\mathbb{Q}} W = 1$ . The above formula for  $H^1(G; \mathbb{Q}G)$  applied to  $W$  and to  $W/\langle x \rangle = V$  yields  $H^1(W; \mathbb{Q}W) \cong H^0(V; \mathbb{Q}V) \neq 0$ . This implies that  $V$  is finite.

We thus have proved that  $G/\langle x \rangle$  is the fundamental group of a finite graph of finite edge *and* vertex groups. Such a group is well-known to contain a (normal) free subgroup of finite index; from the corresponding spectral sequence we obtain the required result

$$H_i(G/\langle x \rangle; \mathbb{Q}) = 0$$

for  $i \geq 2$ .

**2.6.** From Theorems 2.2, 2.3, 2.4', 2.5 we immediately obtain the result claimed in 1.3 for cyclic group homology:

**COROLLARY 2.6.** *Let  $G$  be a group with  $hd_{\mathbb{Q}} G = n < \infty$  and belonging to one of the classes (a), (b), (c), or (d),  $n = 2$  in the case (d). Then for  $i \geq n$  the cyclic homology  $HC_i(\mathbb{Q}G)$  vanishes on the conjugacy classes of elements of infinite order.*

### 3. Cyclic homology characters of $\mathbb{Q}G$

**3.1.** The Connes character  $Ch'_0$  of  $\mathbb{Q}G$  (cf. Karoubi [8]) is a homomorphism of  $K_0(\mathbb{Q}G)$  to  $HC_{2l}(\mathbb{Q}G)$ ,  $l = 0, 1, 2, \dots$ ; we will write  $Ch^l$  for  $Ch'_0$  since we will not consider here the higher characters  $Ch'_i$  (see however Remark 3.2). The  $Ch^l$  are compatible with the map  $S$  in the Connes–Gysin sequence (1.1), i.e.,

$$\begin{array}{ccc} & & HC_{2l}(\mathbb{Q}G) \\ & \nearrow Ch^l & \downarrow S \\ K_0(\mathbb{Q}G) & \xrightarrow{Ch^{l-1}} & HC_{2l-2}(\mathbb{Q}G) \end{array}$$

is commutative. Corollary 2.6 immediately yields

**THEOREM 3.1.** *Let  $G$  be a group of finite dimension  $hd_{\mathbb{Q}}G = n$  belonging to one of the classes (a), (b), (c) or (d). Then the characters  $Ch^l$ ,  $l = 0, 1, 2, \dots$  all have 0-components in the summands  $\bigoplus_{[x]}''$  corresponding to elements  $x$  of infinite order. In particular, if  $G$  is torsion-free, the  $Ch^l$  are concentrated on the  $[1]$ -summand, i.e. lie in  $[H_*(G; \mathbb{Q}) \otimes H_*(\mathbb{C}P^\infty; \mathbb{Q})]_{2l}$ .*

*Remark 3.2.* A similar result holds, of course, for the higher characters  $Ch_i^l: K_i(\mathbb{Q}G) \rightarrow HC_{2l+i}(\mathbb{Q}G)$ .

**3.2.** A look at the definitions shows that  $Ch^0: K_0(\mathbb{Q}G) \rightarrow HC_0(\mathbb{Q}G)$  is the same as the Hattori–Stallings rank, as follows.

By (1.2)  $HC_0$  is isomorphic to Hochschild homology  $HH_0$ . For any  $\mathbb{Q}$ -algebra  $\Lambda$  the latter, with  $\Lambda$  as bimodule by left- and right-multiplication for coefficients,  $HH_0(\Lambda)$  is well-known to be  $\Lambda/\{\lambda\mu - \mu\lambda\}$ , where  $\{\lambda\mu - \mu\lambda\}$  is the  $\mathbb{Q}$ -sub-module generated by all  $\lambda\mu - \mu\lambda$ ,  $\lambda, \mu \in \Lambda$ . In Remark 1.1 we have written  $\overline{\mathbb{Q}G}$  for  $HH_0(\mathbb{Q}G)$  and  $T: \mathbb{Q}G \rightarrow \overline{\mathbb{Q}G}$  for the canonical map. Similarly, for the matrix algebra  $\mathbb{M}_k(\mathbb{Q}G)$ , we have  $HH_0(\mathbb{M}_k(\mathbb{Q}G)) = \overline{\mathbb{M}_k(\mathbb{Q}G)}$  with  $T: \mathbb{M}_k(\mathbb{Q}G) \rightarrow \overline{\mathbb{M}_k(\mathbb{Q}G)}$ . The trace of matrices  $\text{tr}: \mathbb{M}_k(\mathbb{Q}G) \rightarrow \mathbb{Q}G$  induces an isomorphism  $\text{tr}: \overline{\mathbb{M}_k(\mathbb{Q}G)} \rightarrow \overline{\mathbb{Q}G}$ , and clearly  $T \circ \text{tr} = \text{tr} \circ T: \mathbb{M}_k(\mathbb{Q}G) \rightarrow \overline{\mathbb{Q}G}$ .

Now  $Ch^0$  is defined, on a finitely generated projective  $\mathbb{Q}G$ -module  $P$  representing an element of  $K_0(\mathbb{Q}G)$ , as follows: Let  $p$  be an idempotent matrix  $\in \mathbb{M}_k(\mathbb{Q}G)$ , for suitable  $k$ , describing  $P$  as a direct summand of a free  $\mathbb{Q}G$ -module  $M$ , and put  $Ch^0 p = \text{tr} \circ T(p) \in \overline{\mathbb{Q}G}$ , i.e.,  $= T \circ \text{tr}(p)$ . This is precisely the definition of the Hattori–Stallings rank  $r_p \in \overline{\mathbb{Q}G}$ , independent of choices and of bases in  $M$ . We recall that  $\overline{\mathbb{Q}G}$  is the  $\mathbb{Q}$ -module having the conjugacy classes  $[x]$  as basis.

**THEOREM 3.3.** *For the groups  $G$  of finite dimension  $hd_{\mathbb{Q}}G$  belonging to one of the classes (a), (b), (c), (d), the Hattori–Stallings rank  $r_p$  of a finitely generated projective  $\mathbb{Q}G$ -module  $P$  vanishes on the conjugacy classes of elements of infinite order.*

*Remark 3.4.* The vanishing of the character map  $Ch^l: K_0(\mathbb{Q}G) \rightarrow HC_{2l}(\mathbb{Q}G)$  on the conjugacy classes of elements of infinite order, in particular of the Hattori–Stallings rank  $Ch^0$ , would of course follow from properties much weaker than those established in Section 2 for certain classes of groups. Indeed it suffices that, for a group  $G$  under consideration, some iteration  $S^k: HC_{2l+2k}(\mathbb{Q}G) \rightarrow HC_{2l}(\mathbb{Q}G)$ ,  $1 \leq k \leq \infty$ , of  $S$  in (1.1) is zero on the conjugacy classes of elements of infinite order;  $k = \infty$  refers to the inverse limit. It has been conjectured, for

example, that this is the case for  $k = \infty$  and for all groups having a finite Eilenberg–MacLane complex, cf. [7].

*Note Added in Proof.* The proof of Theorem 2.3, without assuming  $G$  to be torsion-free (and hence also of Theorem 2.2), becomes much simpler if one uses the fact that the Hirsch number of a factor group of  $G$  is less or equal to that of  $G$ , combined with Stammbach's theorem [11].

## REFERENCES

- [1] R. ALPERIN and P. SHALEN, *Linear groups of finite cohomological dimension*, Bull. Am. Math. Soc. (New Series) 4, 339–341; Inventiones Math. 66, 89–98 (1982).
- [2] R. BAER and H. HEINEKEN, *Radical groups of finite Abelian subgroup rank*, Illinois J. of Math. 16, 533–586 (1972).
- [3] H. BASS, *Euler characteristics and characters of discrete groups*, Inventiones Math. 35, 155–196 (1976).
- [4] R. BIERI, *Homological dimension of discrete groups*, Queen Mary College Lecture Notes (1976).
- [5] R. BIERI, *On groups of cohomology dimension 2*, in Topology and Algebra (Enseignement Math. 1978), 55–62.
- [6] R. BIERI, *A connection between the integral homology and the centre of a rational linear group*, Math. Z. 170, 263–266 (1980).
- [7] D. BURGHELEA, *The cyclic homology of the group rings*, Comment. Math. Helv. 60, 354–365 (1985).
- [8] M. KAROUBI, *Homologie cyclique et K-théorie algébrique*, C.R. Acad. Sc. Paris, 297 S.I., 447–450 (1983).
- [9] S. MACLANE, *Homology* (Springer Verlag Berlin 1975), Chapt. X.
- [10] J.-P. SERRE, *Cohomologie des groupes discrets*, Annals of Math. Studies 70 (Princeton University Press 1971).
- [11] U. STAMMBACH, *On the weak homological dimension of the group algebra of solvable groups*, J. London Math. Soc. (2) 2, 567–570 (1970).
- [12] R. STREBEL, *A homological finiteness criterion*, Math. Z. 151, 263–275 (1976).
- [13] B. A. F. WEHRFRITZ, *Infinite linear groups* (Springer Verlag Berlin 1973).

*Mathematics*  
*ETH-Zentrum*  
*CH-8092 Zürich/Switzerland*

Received November 28, 1985