

# On the classification of constant mean curvature tori in $R^3$ .

Autor(en): **Jaggy, Christian**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **69 (1994)**

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-52279>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# On the classification of constant mean curvature tori in $\mathbb{R}^3$

CHRISTIAN JAGGY

## 1. Introduction

Let  $S$  be a compact oriented surface and  $i : S \rightarrow \mathbb{R}^3$  an immersion with constant mean curvature. Hopf [6] investigated such immersions, and for genus  $(S) = 0$  he showed that  $i : S \rightarrow \mathbb{R}^3$  must be an embedding of a round sphere. Conversely, the genus of the surface  $S$  is 0, if  $i$  is an embedding. This statement was proved by Alexandrov [1]. Only a few years ago Wente [10] and Kapouleas [7] proved the existence of constant mean curvature immersions for genus  $(S) = 1$  and genus  $(S) \geq 2$ , respectively. In this work we will only look at constant mean curvature immersions with genus  $(S) = 1$ .

First the relation of hyperelliptic curves and constant mean curvature immersions is sketched. For a rigorous formulation see Bobenko [3].

Let  $u$  be a solution of the elliptic-sinh Gordon equation

$$u_{,w\bar{w}} + \sinh u = 0 \tag{1}$$

on a simply-connected domain  $\Omega \subset \mathbb{C}$ . There is an algorithm that associates an immersion  $i : \Omega \rightarrow \mathbb{R}^3$  to  $u$  with constant mean curvature  $\frac{1}{2}$ . Conversely, every constant mean curvature immersion yields a solution  $u$  of equation (1).

On the other hand we can associate quasi-periodic solutions of equation (1) on  $\mathbb{R}^2$  to hyperelliptic curves

$$X : y^2 = x \prod_{i=1}^{2g} (x - e_i) \tag{2}$$

where the branch points are distinct and satisfy

$$e_{i+g} = \frac{1}{\bar{e}_i}, \quad i = 1, \dots, g. \tag{3}$$

We first have to fix some notation to write down an explicit formula for solutions




Figure 1

of equation (1). In figure (1) a canonical basis  $a_1, \dots, a_g, b_1, \dots, b_g$  of  $H_1(X, \mathbb{Z})$  with intersection numbers

$$a_i b_j = \delta_{ij}, \quad a_i a_j = 0, \quad b_i b_j = 0, \quad i, j = 1, \dots, g \quad (4)$$

is introduced. Let  $\Omega_0$  and  $\Omega_\infty$  be meromorphic differentials on  $X$ , holomorphic outside 0 and  $\infty$ , respectively, which satisfy the conditions

$$\int_{a_i} \Omega_0 = \int_{a_i} \Omega_\infty = 0, \quad i = 1, \dots, g \quad (5)$$

and

$$\begin{aligned} \Omega_0 &\text{ has a pole of second order at } 0, \\ \Omega_\infty &\text{ has a pole of second order at } \infty. \end{aligned} \quad (6)$$

Define the vectors  $\mu_0, \mu_\infty$  by

$$\begin{aligned} \mu_0 &= \left( \int_{b_1} \Omega_0, \dots, \int_{b_g} \Omega_0 \right) \\ \mu_\infty &= \left( \int_{b_1} \Omega_\infty, \dots, \int_{b_g} \Omega_\infty \right), \end{aligned}$$

and for  $\zeta \in \mathbb{C}^g$  put

$$u(\zeta) = 2 \log \frac{\theta\left(\zeta + \left(\frac{1}{2}, \dots, \frac{1}{2}\right)\right)}{\theta(\zeta)}$$

where  $\theta$  is the Riemann theta function of  $X$  for the given homology basis. The function

$$u(\zeta + w\mu_0 + \bar{w}\mu_\infty) \tag{7}$$

is a real quasi-periodic solution of equation (1) for every  $\zeta \in \mathbb{R}^g$ .

The question arises, whether it is possible to choose  $X$  in a way, such that  $X$  yields constant mean curvature tori. The answer to this question was given by Bobenko [4] and Pinkall-Sterling [9].

#### THEOREM 1.1.

(1) *Under the correspondence mentioned above  $X$  yields constant mean curvature tori in  $\mathbb{R}^3$  if and only if*

(a)  $\Omega_\infty$  has a root  $p = (x_0, y_0)$  with  $|x_0| = 1$ ;

(b) *Let  $\gamma$  be a path that connects the two points  $(x_0, y_0)$  and  $(x_0, -y_0)$ . Then the span of the vectors*

$$v_0 = \left( \int_\gamma \Omega_0, \int_{b_1} \Omega_0, \dots, \int_{b_g} \Omega_0 \right)$$

$$v_\infty = \left( \int_\gamma \Omega_\infty, \int_{b_1} \Omega_\infty, \dots, \int_{b_g} \Omega_\infty \right)$$

*in  $\mathbb{C}^{g+1}$  must contain two linearly independent rational vectors. In this case one gets a  $(g - 2)$ -parameter family of constant mean curvature tori.*

(2) *Every constant mean curvature torus arises in such a way.*

It is known that there are no curves satisfying the condition (a) for genus  $(X) = 1$ . Wente found constant mean curvature tori which are known to correspond to curves with genus  $(X) = 2$  or genus  $(X) = 3$ . In 1991 Ercolani–Knörrer–Trubowitz [5] proved the existence of such curves for even genus  $(X)$ . All curves constructed there have the additional property, that the set of branch points is invariant under the map  $x \mapsto 1/x$ . In this paper the existence of curves  $X$  fulfilling the conditions (a) and (b) is proved for genus  $(X)$  arbitrary.

## 2. Preliminaries

The map  $\sigma : X \rightarrow X$

$$(x, y) \mapsto \left( \frac{1}{\bar{x}}, \frac{\left( \prod_{i=1}^{2g} e_i \right)^{1/2} \bar{y}}{\bar{x}^{g+1}} \right)$$

is an antiholomorphic involution of  $X$ . The sign of  $(\prod_{i=1}^{2g} e_i)^{1/2}$  is chosen in such a way, that the points lying over  $S^1$  are fixed points of  $\sigma$ . Then  $\sigma_*$  acts as follows on the cycles:

$$\sigma_*(a_i) = -a_i, \quad i = 1, \dots, g \tag{8}$$

$$\sigma_*(b_i) = b_i + \sum_{j=1}^g \lambda_{ij} a_j, \quad i = 1, \dots, g$$

with  $\lambda_{ij} \in \mathbb{Z}$ ;  $i, j = 1, \dots, g$ , and

$$\gamma - \sigma_* \gamma = \sum_{j=1}^g \mu_j a_j, \tag{9}$$

with  $\mu_j \in \mathbb{Z}$ ;  $j = 1, \dots, g$ .

It is possible to choose  $\Omega_0, \Omega_\infty$  in a way, such that

$$\sigma^* \Omega_0 = \bar{\Omega}_\infty$$

holds. It follows that the vectors  $v_0, v_\infty$  are complex conjugate. The new vectors

$$v := v_0 + v_\infty$$

$$w := i(v_\infty - v_0)$$

are elements of  $\mathbb{R}^{g+1}$ .

Now consider the map  $f : \mathbb{C}^g \rightarrow \mathbb{C} \times Gr(2, \mathbb{R}^{g+1})$

$$(e_1, \dots, e_g) \mapsto (\text{root of } \Omega_\infty, \langle v, w \rangle).$$

$f$  is a multivalued function and one should restrict the domain of definition of  $f$  to the open subset  $U \subset \mathbb{C}^g$ , where all the branch points are distinct.  $Gr(2, \mathbb{R}^{g+1})$

denotes the Grassmannian of 2-dimensional subspaces of  $\mathbb{R}^{g+1}$ . The vectors  $v$  and  $w$  are linearly independent and  $\langle v, w \rangle$  is a welldefined element of  $Gr(2, \mathbb{R}^{g+1})$ .

It is interesting to look at this map, because if one finds a root  $p = (x_0, y_0)$  with  $|x_0| = 1$  and if  $\langle v, w \rangle$  contains two linearly independent rational vectors, the existence of constant mean curvature tori is guaranteed. In section 3 the following theorem will be proved.

**THEOREM 2.1.** *Let  $e = (e_1, \dots, e_g)$  be in  $U$ . Assume that the differentials  $\Omega_0, \Omega_\infty$  on the hyperelliptic curve*

$$X : y^2 = x \prod_{i=1}^{2g} (x - e_i)$$

*fulfill the following conditions:*

- (1)  $\Omega_0, \Omega_\infty$  have a common root  $\alpha$  over  $x = 1$ ,
- (2)  $\Omega_0, \Omega_\infty$  don't have any other common roots,
- (3)  $(\Omega_\infty - \Omega_0)(e_m) \neq 0$  for  $m = 1, \dots, 2g$ , and  $\Omega_\infty - \Omega_0$  has a root of order 1 at  $\alpha$ .

*Then  $df(e)$  is invertible.*

We denote  $X_e$  as the hyperelliptic curve associated to the point  $e \in U$ . Due to this theorem it follows, that arbitrarily close to  $e$  there are points, such that the corresponding curves  $X_e$  fulfill conditions (a) and (b). In section 4 we will finally show

**THEOREM 2.2.** *For every  $g \geq 2$  there are curves  $X_e, e \in U$ , satisfying the conditions (1), (2), (3) above.*

This theorem will be proved by induction on  $g$ .

### 3. Simplification

*Proof of Theorem 2.1.* Since dimensions are equal it is enough to show that  $df(e)$  is injective. The strategy is due to Krichever [8], Bikbaev and Kuksin [2].

Let  $e(\tau), \tau \in \mathbb{R}$ , be an arbitrary differentiable curve passing through  $e$ , such that  $f(e(\tau))$  changes only in order  $\tau^2$ , in other words

$$\begin{pmatrix} v(\tau) \\ w(\tau) \end{pmatrix} = A(\tau) \begin{pmatrix} v(0) \\ w(0) \end{pmatrix} + \mathcal{O}(\tau^2), \quad \text{with } A(0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (10)$$

$$\alpha(\tau) = 1 + \mathcal{O}(\tau^2). \quad (11)$$

We want to conclude that

$$\frac{d}{d\tau} e(\tau) \Big|_{\tau=0} = 0$$

holds. This implies that  $df(e)$  is injective. Put  $B(\tau) := A(\tau)^{-1}$ , clearly

$$B(\tau) \begin{pmatrix} v(\tau) \\ w(\tau) \end{pmatrix} = \begin{pmatrix} v(0) \\ w(0) \end{pmatrix} + \mathcal{O}(\tau^2)$$

and after differentiation

$$\dot{B}(0) \begin{pmatrix} v(0) \\ w(0) \end{pmatrix} + \begin{pmatrix} \dot{v}(0) \\ \dot{w}(0) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \tag{12}$$

These are  $2g + 2$  equations,  $2g$  of them describe relations among period integrals. Define differentials  $\omega_1, \omega_2$  by

$$\begin{pmatrix} \omega_1(\tau) \\ \omega_2(\tau) \end{pmatrix} := B(\tau) \begin{pmatrix} \Omega_0(\tau) + \Omega_\infty(\tau) \\ i(\Omega_\infty(\tau) - \Omega_0(\tau)) \end{pmatrix}. \tag{13}$$

By integration of  $\omega_1(\tau), \omega_2(\tau)$  one get's multivalued meromorphic functions on  $X_{e(\tau)}$ :

$$\Omega_i(P, \tau) := \int_{J(P)}^P \omega_i(\tau), \quad i = 1, 2 \tag{14}$$

where  $J$  denotes the hyperelliptic involution.

LEMMA 3.1. *The functions*

$$\frac{\partial}{\partial \tau} \Omega_i(P, \tau) \Big|_{\tau=0}$$

are single-valued meromorphic functions on  $X_e$ . At the points  $e_1, \dots, e_{2g}, 0, \infty$  they have first order poles. Furthermore there are non-zero complex numbers  $c_1, \dots, c_{2g}$  such that

$$\text{res}_{P=e_m} \left( \frac{\partial}{\partial \tau} \Omega_2(P, \tau) \Big|_{\tau=0} \right) = c_m \frac{\partial}{\partial \tau} e_m \Big|_{\tau=0}, \quad m = 1, \dots, 2g. \tag{15}$$

Due to this lemma it is enough to show that

$$\frac{\partial}{\partial \tau} \Omega_2(P, \tau) \Big|_{\tau=0} \equiv 0.$$

This will prove the theorem. We first prove this lemma, before we continue the proof of the theorem.

*Proof.* To see that the functions  $(\partial/\partial\tau)\Omega_i(P, \tau) |_{\tau=0}$  are single-valued, we have to look at the corresponding  $b$ -periods:

$$\begin{aligned} \frac{\partial}{\partial \tau} \int_{b_j} \omega_1 \Big|_{\tau=0} &= \frac{\partial}{\partial \tau} \int_{b_j} (b_{11}(\Omega_0 + \Omega_\infty) + b_{12}i(\Omega_\infty - \Omega_0)) \Big|_{\tau=0} \\ &= \int_{b_j} (\dot{b}_{11}(0)(\Omega_0 + \Omega_\infty) + \dot{b}_{12}(0)i(\Omega_\infty - \Omega_0) + \dot{\Omega}_0 + \dot{\Omega}_\infty) \\ &= 0. \end{aligned}$$

The last identity is true due to equation (12). The same is true for  $\omega_2$  and the first statement is proved.

Expand  $\omega_i(\tau)$  at  $e_m(\tau)$  in the local coordinate  $(x - e_m(\tau))^{1/2}$ :

$$\omega_i(x, \tau) = \sum_{k=-1}^{\infty} (x - e_m(\tau))^{k/2} x_k^{i,m}(e(\tau)) dx.$$

Put  $P = (x, y)$ , then we get

$$\begin{aligned} \frac{\partial}{\partial \tau} \Omega_i(P, \tau) \Big|_{\tau=0} &= \int_{J(P)} \frac{\partial}{\partial \tau} \omega_i(x, \tau) \Big|_{\tau=0} \\ &= 2 \sum_{k=-1}^{\infty} \left( -(x - e_m(0))^{k/2} \frac{\partial}{\partial \tau} e_m(0) x_k^{i,m}(e(0)) \right. \\ &\quad \left. + \frac{2}{2+k} (x - e_m)^{1+k/2} \frac{\partial}{\partial \tau} x_k^{i,m}(e(\tau)) \Big|_{\tau=0} \right). \end{aligned}$$

It follows that the functions  $(\partial/\partial\tau)\Omega_i(P, \tau) |_{\tau=0}$  have first order poles at the points  $e_1, \dots, e_{2g}$  and the same is true for 0 and  $\infty$  by a similar calculation. Due to the assumption (3) in Theorem 2.1 the claim about the numbers  $c_m$  is obvious.  $\square$



Let's continue the proof of the theorem. Take  $P \in X_e$  with  $\omega_2(P) \neq 0$ . The implicit function theorem yields a curve  $P(\tau)$  with

$$\Omega_2(P(\tau), \tau) = \Omega_2(P, 0) \quad (16)$$

and after differentiation

$$\frac{d}{d\tau} \Omega_2(P(\tau), \tau) \Big|_{\tau=0} = \omega_2(P) \frac{d}{d\tau} P(\tau) \Big|_{\tau=0} + \frac{\partial}{\partial \tau} \Omega_2(P, \tau) \Big|_{\tau=0} = 0. \quad (17)$$

Define a new function

$$\dot{\Omega}_1(P) := \frac{d}{d\tau} \Omega_1(P(\tau), \tau) \Big|_{\tau=0}. \quad (18)$$

The function  $\dot{\Omega}_1$  is welldefined and by the equation (17) above one gets

$$\dot{\Omega}_1(P) = \frac{\partial}{\partial \tau} \Omega_1(P, \tau) \Big|_{\tau=0} - \frac{\partial}{\partial \tau} \Omega_2(P, \tau) \Big|_{\tau=0} \cdot \frac{\omega_1(P)}{\omega_2(P)}. \quad (19)$$

It follows that  $\dot{\Omega}_1$  is a meromorphic function on  $X$ . To finish the proof of the theorem we need the following lemma:

**LEMMA 3.2.** *The functions  $(\partial/\partial\tau)\Omega_i(P, \tau) \Big|_{\tau=0}$  have a root of order 2 at  $\alpha$ .*

*Proof.* By equation (11) the differentials  $(\partial/\partial\tau)\omega_i(P, \tau) \Big|_{\tau=0}$  have a root of order 1 at  $\alpha$ . The functions

$$h_i(P) := \int_{\alpha}^P \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0}$$

have a root or order 2 at  $\alpha$ . Now look at

$$\begin{aligned} \frac{\partial}{\partial \tau} \Omega_i(P, \tau) \Big|_{\tau=0} &= \int_{J(P)}^P \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0} \\ &= \int_{J(P)}^{J(\alpha)} \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0} + \int_{J(\alpha)}^{\alpha} \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0} \\ &\quad + \int_{\alpha}^P \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0}. \end{aligned}$$

With equation (12) one gets

$$\int_{J(\alpha)}^{\alpha} \frac{\partial}{\partial \tau} \omega_i(P, \tau) \Big|_{\tau=0} = 0$$

and this implies

$$\frac{\partial}{\partial \tau} \Omega_i(P, \tau) \Big|_{\tau=0} = 2h_i(P). \quad \square$$

$\dot{\Omega}_1$  has  $2g$  roots at the branch points  $e_1, \dots, e_{2g}$  and another 4 roots over  $x = 1$ . The roots of  $\omega_2$  lying outside the set  $\{\alpha, J(\alpha)\}$  yield  $2g$  poles of  $\dot{\Omega}_1$ , together with the simple poles at 0 and  $\infty$  we see that  $\dot{\Omega}_1$  has at most  $2g + 2$  poles. Consequently  $\dot{\Omega}_1$  is the zero-function and one gets the following equation:

$$\frac{\partial}{\partial \tau} \Omega_2(P, \tau) \Big|_{\tau=0} \cdot \omega_1(P) = \frac{\partial}{\partial \tau} \Omega_1(P, \tau) \Big|_{\tau=0} \cdot \omega_2(P). \quad (20)$$

There are  $2g$  roots of  $\omega_2$  outside the set  $\{\alpha, J(\alpha)\}$ , which can't coincide with roots of  $\omega_1$  due to the assumption (2). These  $2g$  roots of  $\omega_2$  must be roots of  $(\partial/\partial\tau)\Omega_2(P, \tau) \Big|_{\tau=0}$ . Together with the 4 roots lying over  $x = 1$  we conclude that  $(\partial/\partial\tau)\Omega_2(P, \tau) \Big|_{\tau=0}$  has at least  $2g + 4$  roots. But  $(\partial/\partial\tau)\Omega_2(P, \tau) \Big|_{\tau=0}$  has at most  $2g + 2$  poles at the branch points. We get  $(\partial/\partial\tau)\Omega_2(P, \tau) \Big|_{\tau=0} \equiv 0$  and by Lemma 3.1  $(d/d\tau)e(\tau) \Big|_{\tau=0} = 0$  follows. This proves the theorem.  $\square$

#### 4. Induction

Theorem 2.2 will be proved by induction on  $g$ . We will see that a good configuration of branch points for genus  $g$  yields a good configuration of branch points for genus  $g + 1$ . Let's first prepare the induction step.

Take a point  $e = (e_1, \dots, e_g)$  for which the conditions (1), (2), (3) are fulfilled. The corresponding curve  $X_e$  and differentials  $\Omega_0^g, \Omega_\infty^g, \Omega_0^g + \Omega_\infty^g$  look like

$$X_e : y_0^2 = x \prod_{i=1}^g (x - e_i) \left( x - \frac{1}{\bar{e}_i} \right)$$

$$\Omega_0^g = \frac{c_g \prod_{i=1}^g (x - \beta_i)}{xy_0} dx, \quad \beta_1 = 1, \quad c_g \in \mathbb{C},$$

$$\Omega_\infty^g = \frac{\prod_{i=1}^g (x - \alpha_i)}{y_0} dx, \quad \alpha_1 = 1,$$

$$\Omega_\infty^g - \Omega_0^g = \frac{d_g \prod_{i=1}^{g+1} (x - \xi_i)}{xy_0} dx, \quad \xi_1 = 1, \quad d_g \in \mathbb{C}.$$

For  $(e_1, \dots, e_g, a, t) \in U \times S^1 \times (-\varepsilon, \varepsilon)$ ,  $\varepsilon > 0$  we define

$$X_{(e,a,t)} : y^2 = x(x - ae^t)(x - ae^{-t}) \prod_{i=1}^g (x - e_i) \left(x - \frac{1}{\bar{e}_i}\right),$$

and corresponding normalized differentials

$$\Omega_0^{g+1} = \frac{c_{g+1} \prod_{i=1}^{g+1} (x - \beta_i^{g+1})}{xy} dx, \quad c_{g+1} \in \mathbb{C},$$

$$\Omega_\infty^{g+1} = \frac{\prod_{i=1}^{g+1} (x - \alpha_i^{g+1})}{y} dx,$$

$$\Omega_\infty^{g+1} - \Omega_0^{g+1} = \frac{d_{g+1} \prod_{i=1}^{g+2} (x - \xi_i^{g+1})}{xy} dx, \quad d_{g+1} \in \mathbb{C}.$$

Due to the compactness of  $X_{(e,a,t)}$ , the normalization conditions and the residue theorem one has the following equations

$$\begin{aligned} \alpha_i^{g+1}(e, a, 0) &= \alpha_i, & i &= 1, \dots, g, \\ \alpha_{g+1}^{g+1}(e, a, 0) &= a \\ \xi_i^{g+1}(e, a, 0) &= \xi_i, & i &= 1, \dots, g + 1, \\ \xi_{g+2}^{g+1}(e, a, 0) &= a \end{aligned} \tag{21}$$

and

$$\Omega_\infty^{g+1}(e, a, 0) = \Omega_\infty^g. \tag{22}$$

Due to the reduction (21) we delete the superscript  $g + 1$  from  $\alpha_i^{g+1}, \xi_i^{g+1}$ . Now put

$$\alpha_1 = u_1 + iu_2, \quad e_i = x_i + iy_i, \quad i = 1, \dots, g$$

and let's impose the further conditions on  $X_e$

$$(4) \quad \text{rank} \left( \frac{\partial u_r}{\partial x_i \partial y_j} \right) = 2, \quad r = 1, 2,$$

(5) the real part of the meromorphic function

$$k(x) = 1 + x \frac{\frac{\partial \Omega_\infty^g}{\partial x}}{\frac{\Omega_\infty^g}{dx}}$$

doesn't vanish identically on  $S^1$ .

The conditions (4) and (5) are used to prove the following lemma:

LEMMA 4.1. *The map  $h : U \times S^1 \times (-\varepsilon, \varepsilon) \rightarrow \mathbb{C} \times \mathbb{R}$*

$$(e, a, t^2) \mapsto (\alpha_1, |\alpha_{g+1}|)$$

*has maximal rank in a point  $P = (e, a, 0)$ , where  $\text{Re}(k(a)) \neq 0$ .*

REMARK. This lemma together with the property

$$\frac{\partial \xi_{g+2}}{\partial \tau} \Big|_{t=0} = 0$$

yields the existence of curves  $X_{(e,a,t)}$  of genus  $g + 1$ , which satisfy the conditions (1), (2), (3). Taking  $t$  small enough the conditions (4) and (5) are also fulfilled.

*Proof.* Due to the reduction (21) and condition (4) we have

$$\text{rank} \left( \frac{\partial u_r}{\partial x_i \partial y_j} \right) = 2, \quad \left( \frac{\partial |\alpha_{g+1}|}{\partial x_i \partial y_j} \Big|_P \right) = 0, \quad r = 1, 2; \quad i, j = 1, \dots, g.$$

It remains to prove that

$$\frac{\partial}{\partial t^2} |\alpha_{g+1}| \Big|_P = \operatorname{Re} \left( \frac{\partial}{\partial t^2} \alpha_{g+1} \bar{\alpha}_{g+1} \right) \Big|_P \neq 0.$$

For this we will deduce an equation for  $(\partial/\partial t^2)\alpha_{g+1}|_P$ . Differentiation of  $\Omega_\infty^{g+1}$  yields

$$\frac{\partial}{\partial t^2} \Omega_\infty^{g+1} \Big|_P = \frac{\left( -\sum_{i=1}^{g+1} \frac{\partial}{\partial t^2} \alpha_i \Big|_P \frac{1}{x - \alpha_i} \right) \prod_{i=1}^g (x - \alpha_i)}{y_0} dx + \frac{ax \prod_{i=1}^g (x - \alpha_i)}{2(x - a)^2 y_0} dx.$$

Since

$$\operatorname{res}_{x=a} \left( \frac{\partial}{\partial t^2} \Omega_\infty^{g+1} \Big|_P \right) = 0$$

we get the equation

$$\operatorname{res}_{x=a} \left( \frac{\partial}{\partial t^2} \alpha_{g+1} \Big|_P \Omega_\infty^g \right) = \operatorname{res}_{x=a} \left( \frac{ax}{2(x - a)^2} \Omega_\infty^g \right),$$

and

$$\frac{\partial}{\partial t^2} \alpha_{g+1} \Big|_P \cdot \bar{a} = \frac{1}{2} + \frac{1}{2} x \frac{\frac{\partial}{\partial x} \Omega_\infty^g}{\Omega_\infty^g} \Big|_{x=a}.$$

Since  $\operatorname{Re}(k(a)) \neq 0$  we have

$$\frac{\partial}{\partial t^2} |\alpha_{g+1}| \Big|_P \neq 0,$$

and the lemma is proved. □

Finally, we have to prove the existence of curves  $X_e$  of genus  $g = 2$  which satisfy the conditions (1) up to (5). For the beginning of the induction results of Bobenko [4] and Ercolani–Knörrer–Trubowitz [5] are used.




Figure 2

Let  $X_e$  be the hyperelliptic curve (figure (2))

$$X_e : y^2 = x(x - \mu) \left(x - \frac{1}{\bar{\mu}}\right) (x - \bar{\mu}) \left(x - \frac{1}{\mu}\right) \tag{23}$$

with normalized differentials

$$\Omega_0 = \frac{\bar{\alpha}_1 \bar{\alpha}_2 (x - \beta_1) (x - \beta_2)}{xy} dx,$$

$$\Omega_\infty = \frac{(x - \alpha_1) (x - \alpha_2)}{y} dx.$$

Let  $C_1, C_2$  be the elliptic curves

$$C_1 : y^2 = (z - 2)(z - \lambda)(z - \bar{\lambda}), \quad \lambda = \mu + \frac{1}{\mu}$$

$$C_2 : y^2 = (z + 2)(z - \lambda)(z - \bar{\lambda})$$

and

$$\varphi_v = \frac{(z - \zeta_v)}{y} dz$$

meromorphic differentials on  $C_v$  with vanishing  $a$ -periods (see figure (3)).




Figure 3

There are maps  $\tau_v : X_e \rightarrow C_v$ , given by

$$(x, y) \mapsto \left( x + \frac{1}{x}, \frac{x + (-1)^v}{x^2} y \right).$$

The pullback of  $\varphi_v$  with respect to  $\tau_v$  is given by

$$\tau_1^* \varphi_1 = \frac{(x^2 - \zeta_1 x + 1)(x + 1)}{xy} dx,$$

$$\tau_2^* \varphi_2 = \frac{(x^2 - \zeta_2 x + 1)(x - 1)}{xy} dx.$$

Taking the sum and the difference one gets

$$\tau_1^* \varphi_1 + \tau_2^* \varphi_2 = 2\Omega_\infty,$$

$$\tau_1^* \varphi_1 - \tau_2^* \varphi_2 = 2\Omega_0.$$

Introduce new parameters  $r, \theta$  by the equation

$$\lambda = 2 + re^{i\theta}. \tag{24}$$

Now, look at the following lemma:

**LEMMA 4.2.**

(i) *There is a unique  $\theta = \theta_0 \in (0, \pi/2)$ , such that  $\xi_1(r, \theta_0) = 2$  holds for arbitrary  $r$ ,*

(ii) 
$$\frac{\partial \xi_1}{\partial \theta}(r, \theta_0) = \frac{-r}{2 \sin(\theta_0)},$$

(iii) 
$$\xi_2(r, \theta) = 2 + r \cos(\theta) + \mathcal{O}(r^2).$$

*Proof.* Let's make the change of variables  $\xi = z - 2$  and let's define

$$Z(r, \theta) := \xi_1(r, \theta) - 2.$$

The curve  $C_1$  is given by

$$y^2 = \xi(\xi^2 - 2r\xi \cos \theta + r^2)$$

and for the differential  $\varphi_1$  we have

$$\varphi_1 = \frac{\xi - Z(r, \theta)}{y} d\xi.$$

Following Bobenko [4] one has

$$\int_a^\pi \frac{\xi d\xi}{y} = \sqrt{8r} \int_\theta^\pi \frac{\cos t dt}{\sqrt{\cos \theta - \cos t}},$$

and there is a unique  $\theta = \theta_0 \in (0, \pi/2)$  for which

$$\int_\theta^\pi \frac{\cos t dt}{\sqrt{\cos \theta - \cos t}} = 0.$$

Consequently, we have the equation

$$Z(r, \theta) = 0 \Leftrightarrow \theta = \theta_0.$$

To prove (ii) we first observe that  $Z(r, \theta) = rZ(1, \theta)$ . Differentiation of  $\varphi_1$  yields

$$\frac{\partial}{\partial \theta} \varphi_1(1, \theta) \Big|_{\theta=\theta_0} = \left( -\frac{\partial Z}{\partial \theta}(1, \theta) \right) \Big|_{\theta=\theta_0} \cdot \frac{d\xi}{y} - \sin \theta_0 \frac{\xi^3 d\xi}{y^3} \quad (25)$$

and

$$d\left( \frac{-\xi^2 \cos \theta_0 + \xi}{y} \right) = -\sin^2 \theta_0 \frac{\xi^3 d\xi}{y^3} - \frac{1}{2} \cos \theta_0 \frac{\xi d\xi}{y} + \frac{1}{2} \frac{d\xi}{y}. \quad (26)$$

Due to

$$\int_a^\pi \frac{\partial}{\partial \theta} \varphi_1(1, \theta) \Big|_{\theta=\theta_0} = 0$$



equation (25) gives rise to

$$\frac{\partial Z}{\partial \theta}(1, \theta) \Big|_{\theta=\theta_0} \cdot \int_a \frac{d\xi}{y} = -\sin \theta_0 \int_a \frac{\xi^3 d\xi}{y^3}.$$

Integration of equation (26) yields

$$-\sin \theta_0 \int_a \frac{\xi^3 d\xi}{y^3} = \frac{1 \cos \theta_0}{2 \sin \theta_0} \int_a \frac{\xi d\xi}{y} - \frac{1}{2 \sin \theta_0} \int_a \frac{d\xi}{y}.$$

The first expression on the right is zero and we get

$$\frac{\partial Z}{\partial \theta}(1, \theta) \Big|_{\theta=\theta_0} = -\frac{1}{2 \sin \theta_0},$$

which proves (ii).

The curve  $C_2$  is given by

$$y^2 = (\xi + 4)(\xi^2 - 2r\xi \cos \theta + r^2)$$

and the differential  $\varphi_2$  looks like

$$\varphi_2 = \frac{\xi - (\xi_2 - 2)}{y} d\xi.$$

Put

$$Q(r, \theta) := \frac{1}{2\pi i} \int_a \frac{d\xi}{y},$$

and we have

$$\begin{aligned} Q(0, \theta) &= \operatorname{res}_{\xi=0} \left( \frac{d\xi}{\xi \sqrt{\xi+4}} \right) = \frac{1}{2}, \\ \frac{\partial}{\partial r} Q(r, \theta) \Big|_{r=0} &= \operatorname{res}_{\xi=0} \left( \frac{\partial}{\partial r} \frac{d\xi}{y} \Big|_{r=0} \right) \\ &= \operatorname{res}_{\xi=0} \left( \frac{\cos \theta d\xi}{\xi^2 \sqrt{\xi+4}} \right) = -\frac{1}{16} \cos \theta. \end{aligned}$$

Consequently,

$$Q(r, \theta) = \frac{1}{2} - \frac{1}{16} r \cos \theta + \mathcal{O}(r^2). \quad (27)$$

Similarly we put

$$P(r, \theta) := \frac{1}{2\pi i} \int_a \frac{\xi d\xi}{y},$$

and this yields

$$P(r, \theta) = \frac{1}{2} r \cos \theta + \mathcal{O}(r^2). \quad (28)$$

Since the integral of  $\varphi_2$  over  $a$  is identically zero, (iii) follows from the equations (27) and (28).  $\square$

We use this lemma to prove the final step:

**PROPOSITION 4.3.** *There are curves  $X_e$  of genus  $g = 2$  which satisfy the conditions (1),  $\dots$ , (5).*

*Proof.* For  $\theta = \theta_0$  the differential  $\varphi_1$  has a root over  $z = 2$ . Put  $\zeta_1 = 2$ . Then  $\Omega_0$  and  $\Omega_\infty$  have a common root  $\alpha$  over  $x = 1$  and condition (1) is fulfilled.

For condition (2) we have to look at  $\alpha_2$  and  $\beta_2$ . They satisfy the equations

$$\zeta_2 \beta_2 = 2, \quad 2\alpha_2 = \zeta_2.$$

Suppose  $\alpha_2 = \beta_2$  holds, then we have  $\zeta_2^2 = 4$ , but for  $\zeta_2$  we know

$$\zeta_2(r, \theta) = 2 + r \cos \theta + \mathcal{O}(r^2).$$

For condition (3) we have to examine the roots of  $\Omega_\infty - \Omega_0 = \tau_2^* \varphi_2$ . Due to the equation above for  $\zeta_2$  the roots of the polynomial

$$p(x) = (x^2 - \zeta_2 x + 1)(x - 1)$$

don't lie in the branch points and  $\Omega_\infty - \Omega_0$  has a root of order 1 at  $\alpha$ . For small  $r$  the conditions (1), (2), (3) are satisfied.

Now look at the condition (4). We want to show that the matrix

$$\begin{pmatrix} \frac{\partial u_r}{\partial x_i \partial y_j} \end{pmatrix}$$

with  $e_1 = \mu$  and  $e_2 = \bar{\mu}$  has rank 2. If we rotate the configuration of branch points around the origin, also  $\alpha_1$  is rotated. Moreover, if we move  $\theta$  for fixed  $r$ , the root  $\alpha_1$  can only move on the real axis. Now look at the equations

$$\alpha_1 + \alpha_2 = \frac{1}{2} (\zeta_1 + \zeta_2),$$

$$\alpha_1 \alpha_2 = \frac{1}{2} (\zeta_2 - \zeta_1) + 1.$$

Suppose we have

$$\left. \frac{d\alpha_1}{d\theta} \right|_{\theta = \theta_0} = 0,$$

then we can conclude

$$\left. \frac{d\zeta_1}{d\theta} \right|_{\theta = \theta_0} = 0,$$

but

$$\left. \frac{d\zeta_1}{d\theta} \right|_{\theta = \theta_0} = \frac{-r}{2 \sin \theta_0}.$$

So, the assumption was false and we get the desired result.

For condition (5) we take the limit  $r \rightarrow 0$  and we get  $k(a) = 1/2$  (using the identities  $\mu = 1, \alpha_2 = 1$ ). Thus the proof of the theorem is complete.  $\square$

## REFERENCES

- [1] ALEXANDROV, *Uniqueness theorems for surfaces in the large*, Transl., Ser. II., Am. Math. Soc. 21 (1962), 412–416.
- [2] BIKBAEV, R. and KUKSIN, S., *On the parametrization of finite-gap solutions by frequency vector and wave-number vector and a theorem of I. Krichever*, Lett. Math. Phys. 28 (1993), 115–122.
- [3] BOBENKO, A., *Constant mean curvature surfaces and integrable systems*, Russian Math. Surveys 46:4 (1991), 1–45.

- [4] BOBENKO, A., *All constant mean curvature tori in  $\mathbb{R}^3$ ,  $S^3$ ,  $\mathbb{H}^3$  in terms of theta-functions*, Math. Ann. 290 (1991), 209–245.
- [5] ERCOLANI, N., KNÖRRER, H. and TRUBOWITZ, E., *Hyperelliptic curves that generate constant mean curvature tori in  $\mathbb{R}^3$* , Integrable Systems (The Verdier Memorial Conference), Birkhäuser Progress in Mathematics 115, 81–114.
- [6] HOPF, H., *Differential geometry in the large*, Lect. Notes Math. 1000 (1983).
- [7] KAPOULEAS, N., *Constant mean curvature surfaces in euclidean three-space*, Bull. Am. Math. Soc. 17:2 (1987), 318–320.
- [8] KRICHEVER, I., *Perturbation theory in periodic problems for two-dimensional systems*, Sov. Sci. C. Math. Phys. 9 (1991), 1–101.
- [9] PINKALL, U. and STERLING, I., *On the classification of constant mean curvature tori*, Ann. Math. 130 (1989), 407–451.
- [10] WENTE, H., *Counterexample to a conjecture of Hopf*, Pacific J. Math. 121 (1986), 193–246.

*Department of Mathematics*  
*ETH-Zurich*  
*Switzerland*

Received February 2, 1994