

# An example of an immersed complete genus one minimal surface in $\mathbb{R}^3$ with two convex ends

Autor(en): **Nelli, Barbara**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **73 (1998)**

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-55104>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## An example of an immersed complete genus one minimal surface in $\mathbb{R}^3$ with two convex ends

Barbara Nelli

**Abstract.** We prove the existence of a compact genus one immersed minimal surface  $M$ , whose boundary is the union of two immersed locally convex curves lying in parallel planes.  $M$  is a part of a complete minimal surface with two finite total curvature ends.

**Mathematics Subject Classification (1991).** 53A10, 53C42.

**Keywords.** Minimal surface, convex boundary, Weierstrass representation, elliptic functions.

### 1. Introduction

In 1978 Meeks conjectured that a connected minimal surface bounded by two convex curves in two parallel planes is topologically an annulus; hence it has genus zero. The conjecture has never been proved and the most general result, due to Schoen, is the following.

Let  $\Gamma = \Gamma_1 \cup \Gamma_2$  be any boundary consisting of two Jordan curves in parallel planes; assume that  $\Gamma$  is invariant by reflection through two planes  $P_1, P_2$  orthogonal to the planes of the  $\Gamma_i$  and that both  $P_1$  and  $P_2$  divide  $\Gamma$  into pieces which are graphs with locally bounded slope over the dividing plane. Then any minimal surface spanning  $\Gamma$  is topologically an annulus and is an embedded surface meeting each parallel plane between the planes of the  $\Gamma_i$  in smooth Jordan curves.

In particular, if  $\Gamma_1$  and  $\Gamma_2$  are circles such that the line joining their centers is perpendicular to the planes in which they lie, then  $M$  is a catenoid (cf. [Sc]).

In 1991, Meeks and White studied the space of minimal annuli bounded by convex curves in parallel planes (cf. [MW]).

In this paper we prove the existence of a compact genus one immersed minimal surface  $M$ , whose boundary is the union of two immersed locally convex curves lying in parallel planes. In fact  $M$  is a part of a complete minimal surface with two finite total curvature ends.

The method we use to construct our surface is the following.

It is well known that a minimal surface of genus  $g$  and  $k$  ends can be described

by its Weierstrass representation, that is a triple  $\{\overline{R} \setminus [p_1, \dots, p_k], \omega = f dz, g\}$ , where  $\overline{R}$  is a compact Riemann surface of genus  $g$ ,  $p_1, \dots, p_k$  are points in  $\overline{R}$ ,  $\omega$  is a holomorphic differential on  $R$  and  $g$  is a meromorphic function on  $R$ .

In our setting  $\overline{R}$  is a torus, so we can choose  $f$  and  $g$  to be elliptic functions. For references about the use of elliptic functions in the Weierstrass representation, see [A], [A1], [C], [C1], [R].

I would like to thank Professor Harold Rosenberg for his continual encouragement and advice.

## 2. Statement of results

Consider the lattice  $L(1, i)$  on  $\mathbb{C}$  generated by 1 and  $i$  and let  $T^2$  be the torus  $\mathbb{C}/L(1, i)$ . Let  $\pi : \mathbb{C} \rightarrow T^2$  be the standard projection to the quotient and set  $p_0 = \pi(0)$ ,  $p_1 = \pi(\frac{1}{2})$ ,  $p_2 = \pi(\frac{1+i}{2})$  and  $p_3 = \pi(\frac{i}{2})$ . Finally, let  $\wp$  be the Weierstrass function associated to the lattice  $L(1, i)$  and  $\wp'$  its derivative.

**Theorem 2.1.** *Let  $f, g : T^2 \setminus \{p_0, p_2\} \rightarrow \mathbb{C}$  be the two meromorphic functions defined by*

$$f = \wp^2 \quad g = \frac{\alpha \wp'}{\wp^3}$$

where  $\alpha$  is a real constant depending only on  $L(1, i)$  and  $\wp$ .

Then  $\{T^2 \setminus [p_0, p_2], f dz, g\}$  is the Weierstrass representation of a complete genus one immersed minimal surface  $M$  with finite total curvature.

**Remark 2.2.** The ends of  $M$  cannot be embedded. In fact, if a complete finite total curvature minimal surface has two embedded ends, it is a catenoid (cf. [Sc]).

The functions  $f$  and  $g$  extend meromorphically to  $T^2$  and we have  $g(p_0) = 0$  and  $g(p_2) = \infty$ . Hence the limit normal vector at both ends of  $M$  is vertical. Then we have the following result.

**Theorem 2.3.** *There exists a positive constant  $c \in \mathbb{R}$  such that  $M \cap \{|x_3| \leq c\}$  is a compact genus one immersed minimal surface having the property that each of the boundary curves  $M \cap \{x_3 = \pm c\}$  is a compact locally convex immersed curve.*

## 3. Proof of the theorems

We list some useful classical properties of the function  $\wp$  (cf. [B], [WW]).

By abuse of notation, we often identify points of  $\mathbb{C}$  with points of  $T^2$ . Let  $'$  be the differentiation with respect to the variable  $z \in \mathbb{C}$ .

(i)  $\wp$  is even and  $\wp'$  is odd. We have  $\wp(z), \wp'(z) \in \mathbb{R}$  when  $z \in \mathbb{R}, \wp(p_1) = e_1 \in \mathbb{R}_+^*, \wp(p_2) = 0$  and  $\wp(p_3) = -e_1$ .

The following identities hold:

(ii)  $(\wp')^2 = 4\wp(\wp^2 - e_1^2), \wp'' = 2(3\wp^2 - e_1^2)$ .

(iii)  $\wp(z + p_1) = \frac{e_1(\wp(z) + e_1)}{\wp(z) - e_1}, \wp(z + p_3) = \frac{e_1(\wp(z) - e_1)}{\wp(z) + e_1}, \wp(z + p_2) = -\frac{e_1^2}{\wp(z)}$ .

(iv)  $\wp'(z + p_2) = e_1^2 \frac{\wp'(z)}{\wp(z)^2}$ .

(v)  $\wp(iz) = -\wp(z), \wp'(iz) = i\wp'(z)$ .

(vi) The local expansion of  $\wp$  and  $\wp'$  around  $p_o$  is

$$\wp(z) = \frac{1}{z^2} + \frac{e_1^2}{5}z^2 + O(z^6),$$

$$\wp'(z) = -\frac{2}{z^3} + \frac{2e_1^2}{5}z + O(z^5).$$

*Proof of Theorem 2.1.* It is sufficient to prove that the following conditions are satisfied.

(A)  $z$  is a pole of order  $m$  of  $g \iff z$  is a zero of order  $2m$  of  $f$ .

(B)  $\int_\gamma (1 + |g|^2)|f| = \infty$  for every divergent path  $\gamma$  in  $M$ .

(C)  $\text{Re} \int_\gamma fg = 0$  and  $\int_\gamma fg^2 = \overline{\int_\gamma f}$  for every closed path in  $M$ .

Zeros and poles of  $f, g, fg, fg^2$  in a fundamental region are as in figure 1.




Figure 1.

The function  $g$  does not have poles in  $T^2 \setminus \{p_o, p_2\}$ , hence condition (A) is satisfied.

The expression of the metric on  $M$  in terms of  $\varphi$  is

$$ds = \left( 1 + \alpha^2 \frac{|\varphi'|^2}{|\varphi|^6} \right) |\varphi|^2$$

hence the metric is complete at the ends and condition (B) is satisfied.

We must verify (C) on paths that are not homologous to 0 in  $T^2 \setminus \{p_o, p_2\}$ , i.e. paths around  $p_o$  and  $p_2$  and paths that generate the homology of  $T^2$ . Denote by  $\alpha(p_o)$  and  $\alpha(p_2)$  any closed path around  $p_o$  and  $p_2$  respectively, and by  $\gamma_1$  and  $\gamma_2$  the following paths generating the homology of  $T^2$  :

$$\gamma_1(t) = \frac{i}{4} + t \quad t \in [0, 1]$$

$$\gamma_2(t) = \frac{1}{4} + it \quad t \in [0, 1]$$

The functions  $f$  and  $fg^2$  are even, so they have no residue at  $p_o$ , i.e.

$$\int_{\alpha(p_o)} fg^2 = \int_{\alpha(p_o)} f = 0$$

Furthermore

$$\operatorname{Re} \int_{\alpha(p_o)} fg = \operatorname{Re} \int_{\alpha(p_o)} \frac{\alpha \varphi'}{\varphi} = \operatorname{Re} \left[ \operatorname{Res}_{p_o} \left( 2\pi i \alpha \frac{\varphi'}{\varphi} \right) \right]$$

By the local expansion of  $\varphi$  and  $\varphi'$  around 0 we have that  $\operatorname{Res}_{p_o} \left( 2\pi i \alpha \frac{\varphi'}{\varphi} \right) = -4\pi i \alpha$ , hence for  $\alpha \in \mathbb{R}$  we have

$$\operatorname{Re} \int_{\alpha(p_o)} fg = 0$$

By (iii) and (iv) we have

$$f(z + p_2) = \frac{e_1^4}{\varphi^2(z)},$$

$$fg^2(z + p_2) = \frac{\alpha^2}{e_1^4} (\varphi'(z))^2.$$

Hence  $f(z + p_2)$  and  $fg^2(z + p_2)$  are even functions of  $z$  and this gives

$$\int_{\alpha(p_2)} fg^2 = \int_{\alpha(p_2)} f = 0.$$

By (iii) and (iv) we have

$$fg(z + p_2) = -\alpha \frac{\wp'(z)}{\wp(z)}.$$

Hence, by the computation above, for  $\alpha \in \mathbb{R}$  we have

$$\operatorname{Re} \int_{\alpha(p_2)} fg = 0.$$

Now we verify (C) over  $\gamma_1$  and  $\gamma_2$ . We have

$$\operatorname{Re} \int_{\gamma_i} fg = \operatorname{Re} \int_{\gamma_i} \alpha \frac{\wp'}{\wp} = \alpha [\ln |\wp|]_{\gamma_i(0)}^{\gamma_i(1)} = 0$$

by periodicity of  $\wp$ , as  $\alpha$  is real.

Integral of  $f$  over  $\gamma_1$ : by Cauchy theorem and periodicity we can move  $\gamma_1$  up to the segment from  $p_3$  to  $p_3 + 1$ , hence

$$\int_{\gamma_1} f = \int_0^1 f(p_3 + t) dt = \int_0^1 e_1^2 \frac{(\wp(t) - e_1)^2}{(\wp(t) + e_1)^2} dt$$

where the last equality is given by (iii).

Integral of  $f$  over  $\gamma_2$ : we can move  $\gamma_2$  to the vertical segment from  $p_1$  to  $p_1 + i$ , hence by (iii) and (iv)

$$\int_{\gamma_2} f = \int_0^1 f(p_1 + t) i dt = i \int_0^1 e_1^2 \frac{(\wp(t) - e_1)^2}{(\wp(t) + e_1)^2} dt.$$

Integral of  $fg^2$  over  $\gamma_1$ : we can move  $\gamma_1$  down to the real segment from  $p_o$  to  $p_o + 1$ , hence

$$\int_{\gamma_1} fg^2 = \int_0^1 f(t) g^2(t) dt = \int_0^1 \alpha^2 \frac{\wp'(t)^2}{\wp(t)^4} dt.$$

Integral of  $fg^2$  over  $\gamma_2$ : we can move  $\gamma_2$  to the vertical segment from  $p_o$  to  $p_o + i$ , hence

$$\int_{\gamma_2} fg^2 = \int_0^1 f(it) g^2(it) i dt = -i \int_0^1 \alpha^2 \frac{\wp'(t)^2}{\wp(t)^4} dt.$$

Then  $\alpha$  must satisfy

$$\alpha^2 \int_0^1 \frac{\wp'(t)^2}{\wp(t)^4} dt = \int_0^1 e_1^2 \frac{(\wp(t) - e_1)^2}{(\wp(t) + e_1)^2} dt.$$

If  $t \in \mathbb{R}$  we have  $\varphi(t), \varphi'(t) \in \mathbb{R}$ , hence the two integrals involved in the definition of  $\alpha$  are positive real numbers. Furthermore they are convergent, so  $\alpha \in \mathbb{R}$ .

Since  $g$  and  $f$  extend meromorphically to  $T^2$ ,  $M$  has finite total curvature.  $\square$

Before proving Theorem 2.3 we need the following lemma.

**Lemma 3.1.** *Consider a minimal surface  $M$  with Weierstrass representation given by  $\{f dz, g\}$  such that the vector corresponding to  $g(0)$  is parallel to the  $x_3$ -axis. Then the planar curvature of the intersection curves of  $M$  with the horizontal planes is*

$$k = \frac{1}{|f^2 g|(1 + |g|^2)} \operatorname{Re} \left( \frac{\overline{f g} g'}{g} \right).$$

*Proof.* Let  $\theta = \arg g$  and  $s$  be the arc length of the curve  $M \cap \{x_3 = c\}$ ; then  $k(s) = \frac{d\theta}{ds}$ . As  $\arg g = \operatorname{Im}(\ln g)$ , we have

$$k(s) = \frac{d \operatorname{Im} \ln g}{ds} = \operatorname{Im} \left( \frac{d \ln g}{dz} \frac{dz}{ds} \right) = \operatorname{Im} \left( \frac{g'}{g} \frac{dz}{ds} \right).$$

By the Weierstrass representation we have

$$x_3 = \operatorname{Re} \int f g.$$

Hence, on the curve  $M \cap \{x_3 = c\}$ ,  $\frac{dz}{ds}$  must satisfy

$$0 = \frac{d}{ds} \operatorname{Re} \int f g = \frac{1}{2} \operatorname{Re} \left( f g \frac{dz}{ds} \right).$$

By a straightforward computation we obtain

$$\frac{dz}{ds} = \frac{i}{(1 + |g|^2)|f|} \frac{\overline{f g}}{|f g|}.$$

Then

$$k = \operatorname{Im} \left( \frac{i}{(1 + |g|^2)|f|} \frac{\overline{f g}}{|f g|} \frac{g'}{g} \right) = \frac{1}{|f^2 g|(1 + |g|^2)} \operatorname{Re} \left( \frac{\overline{f g} g'}{g} \right).$$

$\square$

*Proof of Theorem 2.3.* The third coordinate of  $M$  is given by

$$x_3 = \operatorname{Re} \int f g = \operatorname{Re} \int \alpha \frac{\varphi'}{\varphi} = \alpha \ln |\varphi|,$$

since  $\alpha$  is real. Then, any level curve is given by  $|\wp| = c$  and next to the ends this is a compact immersed curve with only one component.

By a straightforward computation, we obtain

$$g'(z) = 2\alpha \left[ \frac{5e_1^2 - 3\wp(z)^2}{\wp(z)^3} \right],$$

$$\frac{g'(z)}{g(z)} = \frac{2(5e_1^2 - 3\wp(z)^2)}{\wp'(z)},$$

$$\overline{f(z)g(z)} = \overline{\alpha} \frac{\overline{\wp'(z)}}{\wp(z)}.$$

By using the expansion of  $\wp$  and  $\wp'$  at  $p_o$  we have

$$\overline{f(z)g(z)} \sim -2\frac{\overline{\alpha}}{\bar{z}},$$

$$\frac{g'(z)}{g(z)} \sim \frac{3}{z},$$

where  $\sim$  denotes equality between the principal parts of the functions in a neighborhood of zero. Hence the sign of the curvature of the level curve next to the end  $p_o$  is the same as the sign of

$$\operatorname{Re}\left(\frac{-6\overline{\alpha}}{\bar{z}z}\right) = -\frac{6\alpha}{|z|^2},$$

$\alpha$  being real.

We use the equality

$$\overline{f(z+p_2)g(z+p_2)} = -\overline{f(z)g(z)}$$

and the fact that in a neighborhood of zero we have

$$\frac{g'(z+p_2)}{g(z+p_2)} = \frac{2(5\wp(z)^2 - 3e_1^2)}{\wp'(z)} \sim -\frac{5}{z},$$

to conclude that the sign of the curvature of the level curve next to the end  $p_2$  is the same as the sign of

$$\operatorname{Re}\left(\frac{-10\overline{\alpha}}{\bar{z}z}\right) = -\frac{10\alpha}{|z|^2}$$

since  $\alpha$  is real.

Thus, if we choose a negative  $\alpha$ , the level curves are locally convex next to the two ends of  $M$ .  $\square$



## References

- [A] F. Abi-Khuzam, A One Parameter Family of Minimal Surfaces with Four Planar Ends, *Intern. Jour. of Math.* **6** (2) (1995), 149–159.
- [A1] F. Abi-Khuzam, Jacobian Elliptic Functions and Minimal Surfaces, *Proc. Amer. Math. Soc.* **123** (12) (1995), 3837–3849.
- [B] P. Byrd, M. Friedman, *Handbook of Elliptic Integrals for Engineers and Scientists*, Springer-Verlag 1971.
- [C] C. Costa, Example of a Complete Minimal Immersion in  $\mathbb{R}^3$  of Genus One and Three Embedded Ends, *Bol. Soc. Bras. Mat.* **15** (1) (1984), 47–54.
- [C1] C. Costa, Complete Minimal Surfaces in  $\mathbb{R}^3$  of Genus One and Four Planar Embedded Ends, *Proc. Amer. Math. Soc.* **119** (4) (1993), 1279–1287.
- [M] W. Meeks III, *Lectures on Plateau's problem*, Inst. Mat. Pura Apl., Rio de Janeiro 1978.
- [MW] W. Meeks III, B. White, Minimal Surfaces Bounded by Convex Curves in Parallel Planes, *Comment. Math. Helv.* **66** (1991), 263–278.
- [R] B. Riemann, Über die Fläche vom kleinsten Inhalt bei gegebener Begrenzung, *Abh. Königl. d. Wiss Göttingen, Mathem. Cl.* **13** (1867), 3–52.
- [Sc] R. Schoen, Uniqueness, Symmetry, and Embeddedness of Minimal Surfaces, *Jour. Diff. Geom.* **18** (1983), 791–809.
- [WW] E. Whittaker, G. Watson, *A Course of Modern Analysis*, Cambridge University Press, 1995.

Barbara Nelli  
Dipartimento di Matematica  
Università di Pisa  
Via Buonarroti 2  
Pisa - Italy  
e-mail: nelli@dm.unipi.it

(Received: June 30, 1997)