

NK0 and NK1 of the groups C4 and D4 : addendum to “Lower algebraic K-theory of hyperbolic 3-simplex reflection groups” by J.-F. Lafont and I. J. Ortiz

Autor(en): **Weibel, Charles**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **84 (2009)**

PDF erstellt am: **23.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-99122>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

NK_0 and NK_1 of the groups C_4 and D_4 **Addendum to “Lower algebraic K -theory of hyperbolic 3-simplex reflection groups” by J.-F. Lafont and I. J. Ortiz**

Charles Weibel

Abstract. In this addendum to [LO] we explicitly compute the Bass Nil-groups $NK_i(\mathbb{Z}[C_4])$ for $i = 0, 1$ and $NK_0(\mathbb{Z}[D_4])$. We also show that $NK_1(\mathbb{Z}[D_4])$ is not trivial. Here C_4 denotes the cyclic group of order 4 and D_4 is the dihedral group of order 8.

Mathematics Subject Classification (2000). 19A31, 19B28, 19D35, 18F25, 16E20.

Keywords. Lower algebraic K -theory, hyperbolic reflection group, Bass Nil-groups.

In [LO], Lafont and Ortiz computed the lower algebraic K -theory of the integral group ring of all 32 hyperbolic 3-simplex reflection groups (see [LO, Tables 6–7]). For 25 of these integral group rings, their computation was completely explicit. For the remaining 7 examples, the expression for some of the K -groups involved the Bass Nil-groups NK_0 and NK_1 associated to D_4 (the dihedral group of order 8).

In [L05], Lück computed the lower algebraic K -theory of the integral group ring of the semi-direct product of the three-dimensional discrete Heisenberg group by C_4 (the cyclic group of order 4). These computations involved the Bass Nil-groups NK_0 and NK_1 associated to C_4 (see [L05, Corollary 3.9]).

In this addendum we compute the Bass Nil-groups

$$NK_n(\mathbb{Z}G) = \ker\{K_n(\mathbb{Z}G[x]) \xrightarrow{x \mapsto 0} K_n(\mathbb{Z}G)\},$$

where G is D_2 , C_4 or D_4 , and $n = 0, 1$. We will use these calculations to complement the calculations of [L05] and [LO] in 1.5 and 2.9 below.

Our calculation will keep track of the additional structure on the groups $NK_n(A)$ given by the Verschiebung and Frobenius operators, V_m and F_m , as well as the continuous module structure over the ring $W(\mathbb{Z})$ of big Witt vectors; the additive group of $W(\mathbb{Z})$ is the abelian group $(1 + x\mathbb{Z}[[x]])^\times$. (See [We80] for more details.) In fact, it is a module over the slightly larger Cartier algebra consisting of row-and-column

finite sums $\sum V_m[a_{mn}]F_n$, where V_m and F_m are the Verschiebung and Frobenius operators, and the $[a]$ are the homotheties operators for $a \in \mathbb{Z}$; see [DW93] as well as Remarks 1.2.1 and 2.4 below. Some of the identities satisfied by these operators include: $V_mV_n = V_{mn}$, $F_mF_n = F_{mn}$, $F_mV_m = m$, $[a]V_m = V_m[a^m]$ and $F_m[a] = [a^m]F_m$.

It is convenient to write V for the continuous $W(\mathbb{F}_2)$ -module $x\mathbb{F}_2[x]$, which, as an abelian group, is just a countable direct sum of copies of $\mathbb{F}_2 = \mathbb{Z}/2$ on generators x^i , $i > 0$. The module structure on V is determined by: $V_m(x^n) = x^{mn}$; $[a]x^n = a^n x^n$; $F_m(x^n) = 0$ if $(m, n) = 1$ ($m > 1$) and $F_d(x^n) = d x^{n/d}$ when $d | n$.

1. The groups C_2 , D_2 and C_4

For the cyclic group $C_2 = \langle \sigma \rangle$ of order two, consider the Rim square:

$$\begin{array}{ccc}
 \mathbb{Z}[C_2] & \xrightarrow{\sigma \mapsto +1} & \mathbb{Z} \\
 \sigma \mapsto -1 \downarrow & & \downarrow q \\
 \mathbb{Z} & \xrightarrow{q} & \mathbb{F}_2
 \end{array}
 \quad \text{or, equivalently,} \quad
 \begin{array}{ccc}
 \mathbb{Z}[C_2] & \xrightarrow{\sigma \mapsto (1,-1)} & \mathbb{Z} \times \mathbb{Z} \\
 \downarrow & & \downarrow q \times q \\
 \mathbb{F}_2 & \xrightarrow{\Delta} & \mathbb{F}_2 \times \mathbb{F}_2
 \end{array}
 \quad (1)$$

from which we immediately get $NK_0(\mathbb{Z}[C_2]) = NK_1(\mathbb{Z}[C_2]) = 0$ as in [Bas68, XII.10.6] and [Mi71, 6.4]. From Guin–Loday–Keune [GL80], [Kou81], the double relative group $NK_2(\mathbb{Z}[C_2], \sigma + 1, \sigma - 1)$ is isomorphic to V , with the Dennis–Stein symbol $\langle x^n(\sigma - 1), \sigma + 1 \rangle$ corresponding to $x^n \in V$. We also have a diagram

$$\begin{array}{ccccccc}
 & & NK_2(\mathbb{Z}[C_2], \sigma + 1, \sigma - 1) \cong V & & & & \\
 & & \downarrow \cong & & & & \\
 0 = NK_3(\mathbb{Z}) & \longrightarrow & NK_2(\mathbb{Z}[C_2], \sigma + 1) & \xrightarrow{\cong} & NK_2(\mathbb{Z}[C_2]) & \longrightarrow & NK_2(\mathbb{Z}) = 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 = NK_2(\mathbb{Z}, 2) & \longrightarrow & NK_2(\mathbb{Z}) = 0 & &
 \end{array}$$

Thus we obtain:

Theorem 1.1. $NK_2(\mathbb{Z}[C_2]) \cong V$ with $\langle x^n(\sigma - 1), \sigma + 1 \rangle$ corresponding to $x^n \in V$.

We now turn to the group $D_2 = C_2 \times C_2$. First we need a calculation. Let $\Phi(V)$ denote the subgroup (and Cartier submodule) $x^2\mathbb{F}_2[x^2]$ of V , and write Ω_R for the Kähler differentials of R , so that $\Omega_{\mathbb{F}_2[x]} \cong \mathbb{F}_2[x]dx$. By abuse, we will write $\mathbb{F}_2[\varepsilon]$ for the 2-dimensional algebra $\mathbb{F}_2[\varepsilon]/(\varepsilon^2)$.

Lemma 1.2. *The map $q: \mathbb{Z}[C_2] \rightarrow \mathbb{F}_2[C_2] \cong \mathbb{F}_2[\varepsilon]$ in (1) induces an exact sequence*

$$0 \rightarrow \Phi(V) \rightarrow NK_2(\mathbb{Z}[C_2]) \xrightarrow{q} NK_2(\mathbb{F}_2[\varepsilon]) \xrightarrow{D} \Omega_{\mathbb{F}_2[x]} \rightarrow 0.$$

Proof. Van der Kallen computed $NK_2(\mathbb{F}_2[\varepsilon])$ in [vdK71, Exemple 3]: there is a split short exact sequence

$$0 \rightarrow V/\Phi(V) \xrightarrow{F} NK_2(\mathbb{F}_2[\varepsilon]) \xrightarrow{D} \Omega_{\mathbb{F}_2[x]} \rightarrow 0, \tag{2}$$

where $F(x^n) = \langle x^n \varepsilon, \varepsilon \rangle$ and $D(\langle f\varepsilon, g + g'\varepsilon \rangle) = f dg$. The map $NK_2(\mathbb{Z}[C_2]) \xrightarrow{q} NK_2(\mathbb{F}_2[\varepsilon])$ sends $\langle x^n(\sigma + 1), \sigma - 1 \rangle$ to $F(x^n) = \langle x^n \varepsilon, \varepsilon \rangle$. By Theorem 1.1 and (2), this map has kernel $\Phi(V)$ and image $F(V/\Phi(V))$. \square

Remark 1.2.1. Although $\Omega_{\mathbb{F}_2[x]}$ is isomorphic to V as an abelian group, it has a different $W(\mathbb{F}_2)$ -module structure. This is determined by the formulas in $\Omega_{\mathbb{F}_2[x]}$:

$$V_m(x^{n-1} dx) = mx^{mn-1} dx, \quad F_m(x^{n-1} dx) = \begin{cases} x^{n/m-1} dx & \text{if } m \mid n, \\ 0 & \text{else.} \end{cases}$$

Grunewald has pointed out that since $\Omega_{\mathbb{F}_2[x]}$ is not finitely generated as a module over the \mathbb{F}_2 -Cartier algebra (of row-and-column finite sums $\sum V_m[a_{mn}]F_n$), or over the subalgebra $W(\mathbb{F}_2)$), neither are $NK_2(\mathbb{F}_2[\varepsilon])$ or (by 1.3 below) $NK_1(\mathbb{Z}[D_2])$.

Theorem 1.3. *For $D_2 = C_2 \times C_2$, $NK_0(\mathbb{Z}[D_2]) \cong V$, $NK_1(\mathbb{Z}[D_2]) \cong \Omega_{\mathbb{F}_2[x]}$ and the image of the map $NK_2(\mathbb{Z}[D_2]) \rightarrow NK_2(\mathbb{Z}[C_2])^2 \cong V^2$ is $\Phi(V) \times V$.*

Proof. We tensor (1) with $\mathbb{Z}[C_2]$. Since $\mathbb{F}_2[C_2] \cong \mathbb{F}_2[\varepsilon]$, $\varepsilon^2 = 0$, and $NK_1(\mathbb{F}_2[C_2]) \cong (1 + x\varepsilon\mathbb{F}_2[x])^\times \cong V$, then the Mayer–Vietoris sequence in [Mi71, Theorem 6.4] for the NK -functor,

$$\begin{aligned} NK_2(\mathbb{Z}[D_2]) &\rightarrow (NK_2(\mathbb{Z}[C_2]))^2 \xrightarrow{q \times q} NK_2(\mathbb{F}_2[\varepsilon]) \rightarrow NK_1(\mathbb{Z}[D_2]) \\ &\rightarrow (NK_1(\mathbb{Z}[C_2]))^2 \rightarrow NK_1(\mathbb{F}_2[\varepsilon]) \xrightarrow{\cong} NK_0(\mathbb{Z}[D_2]) \rightarrow NK_0(\mathbb{Z}[C_2]), \end{aligned} \tag{3}$$

quickly gives $NK_0(\mathbb{Z}[D_2]) \cong NK_1(\mathbb{F}_2[\varepsilon]) \cong V$. By Lemma 1.2, the initial portion of (3) yields the calculation of $NK_1(\mathbb{Z}[D_2])$ and the asserted surjection $NK_2(\mathbb{Z}[D_2]) \twoheadrightarrow \Phi(V) \times V$. \square

Remark. The kernel K of the map $NK_2(\mathbb{Z}[D_2]) \rightarrow V^2$ in Theorem 1.3 has a subgroup generated by the double relative group $NK_2(\mathbb{Z}[D_2], \sigma_1 + 1, \sigma_1 - 1)$, which is isomorphic to $\mathbb{F}_2[\varepsilon] \otimes V$ on the symbols $\langle x^n(a + b\sigma_2)(\sigma_1 + 1), \sigma_2 - 1 \rangle$, where σ_1, σ_2 are the generators of $D_2 = C_2 \times C_2$. The quotient of K by this subgroup is generated by the image of $NK_3(\mathbb{F}_2[\varepsilon])$, a group which I do not know.

The analysis for the cyclic group C_4 of order 4 on generator σ is similar, using the Rim square

$$\begin{array}{ccc}
 \mathbb{Z}[C_4] & \xrightarrow{\sigma \mapsto i} & \mathbb{Z}[i] \\
 \sigma^2 \mapsto 1 \downarrow & & \downarrow i \mapsto 1 + \varepsilon \\
 \mathbb{Z}[C_2] & \xrightarrow{q} & \mathbb{F}_2[\varepsilon].
 \end{array} \tag{4}$$

Theorem 1.4. $NK_1(\mathbb{Z}[C_4]) \cong \Omega_{\mathbb{F}_2[x]}$ and $NK_0(\mathbb{Z}[C_4]) \cong V$.

Proof. Since $\mathbb{Z}[i]$ is a regular ring, the Mayer–Vietoris sequence for (4) reduces to

$$\begin{aligned}
 NK_2(\mathbb{Z}[C_4]) &\xrightarrow{p_2} NK_2(\mathbb{Z}[C_2]) \xrightarrow{q} NK_2(\mathbb{F}_2[\varepsilon]) \rightarrow NK_1(\mathbb{Z}[C_4]) \\
 &\rightarrow NK_1(\mathbb{Z}[C_2]) \rightarrow NK_1(\mathbb{F}_2[\varepsilon]) \xrightarrow{\cong} NK_0(\mathbb{Z}[C_4]) \rightarrow NK_0(\mathbb{Z}[C_2]).
 \end{aligned}$$

The isomorphism marked in this sequence follows from Theorem 1.1. By Lemma 1.2, the image of the first map p_2 is $\Phi(V)$ and the cokernel of the map q is $\Omega_{\mathbb{F}_2[\varepsilon]}$. \square

Remark. The proof provides a surjection $NK_2(\mathbb{Z}[C_4]) \xrightarrow{p_2} \Phi(V)$. The kernel of p_2 contains the image E of the double relative group $NK_2(\mathbb{Z}[C_4], \sigma^2 + 1, \sigma^2 - 1)$, which is isomorphic to $\mathbb{F}_2[\varepsilon] \otimes V$ on symbols $\langle \sigma^2 + 1, x^n(\sigma^2 - 1) \rangle$. The quotient $\ker(p_2)/E$ is generated by the image of $NK_3(\mathbb{F}_2[\varepsilon])$, which I do not know.

Here is an application of this calculation. Let Hei denote the *three-dimensional discrete Heisenberg group*, which is the subgroup of $GL(3, \mathbb{Z})$ consisting of upper triangular integral matrices with ones along the diagonal. Consider the action of the cyclic group C_4 given by

$$\begin{pmatrix} 1 & x & y \\ 0 & 1 & z \\ 0 & 0 & 1 \end{pmatrix} \mapsto \begin{pmatrix} 1 & -z & y - xz \\ 0 & 1 & x \\ 0 & 0 & 1 \end{pmatrix}.$$

Combining Theorem 1.4 with [L05, 3.9], the lower K -theory of the group $\text{Hei} \rtimes C_4$ is given in the following proposition.

Proposition 1.5. *We have*

$$\text{Wh}_n(\text{Hei} \rtimes C_4) = \begin{cases} \bigoplus_{\infty} \mathbb{Z}/2, & n = 0, 1, \\ 0, & n \leq -1. \end{cases}$$

2. The dihedral group D_4

Before moving on to the group ring of D_4 , we need some facts about the double relative groups $K_1(A, B, I)$ when $A \rightarrow B$ is an injection. These groups were described in [GW83, 0.2] as follows:

$$K_1(A, B, I) \cong (B/A) \otimes (I/I^2) / \{b \otimes cz + c \otimes zb - bc \otimes z\} \quad (b, c \in B, z \in I). \quad (5)$$

Moreover, by [GW83, 3.12 and 4.1], the map $K_1(A, B, I) \rightarrow K_1(A, I)$ sends the class of $b \otimes z$ to the class of the matrix $\begin{pmatrix} 1-zb & z \\ -bzb & 1+bz \end{pmatrix}$.

Lemma 2.1. *Suppose that $A \rightarrow B$ is a ring homomorphism mapping an ideal I of A isomorphically onto an ideal of B . Then the double relative group satisfies*

$$K_1(A[x], B[x], I[x]) \cong K_1(A, B, I) \otimes \mathbb{Z}[x],$$

and

$$NK_1(A, B, I) \cong K_1(A, B, I) \otimes x\mathbb{Z}[x].$$

Proof. Because x is central in $B[x]$, the formulas are immediate from (5). □

We will be specifically interested in the twisted group ring $A = \mathbb{Z}[i] \rtimes C_2$, where $C_2 = \langle \tau \rangle$ acts on $\mathbb{Z}[i]$ by $\tau i \tau^{-1} = -i$. It injects into the matrix ring $B = M_2(\mathbb{Z})$ by the map $\phi: A \rightarrow B$ defined by $\phi(i) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $\phi(\tau) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. The ideal $I = (2, 1 + \tau)A$ maps isomorphically to $2B$, and $A/I \cong \mathbb{F}_2[\varepsilon_1]$, where $\varepsilon_1 = 1 + i$ and $\varepsilon_1^2 = 0$. Hence we have the following cartesian square:

$$\begin{array}{ccc} A & \xrightarrow{\phi} & M_2(\mathbb{Z}) \\ \text{mod } I \downarrow & & \downarrow \text{mod } 2B \\ \mathbb{F}_2[\varepsilon_1] & \longrightarrow & M_2(\mathbb{F}_2). \end{array} \quad (6)$$

To calculate $NK_1(A)$, we use the following double relative calculation.

Lemma 2.2. *The double relative group $K_1(A, B, 2B)$ of (6) is isomorphic to \mathbb{F}_2 , and $NK_1(A, B, 2B) \cong V$. The map $V \cong NK_1(A, B, 2B) \rightarrow NK_1(A)$ sends $x^n \in V$ to the class of the unit $1 + x^n i(1 + \tau)$ of $A[x]$.*

Proof. Since $\dim(B/A) = 2$ and $\dim(I/I^2) = 4$, the group $(B/A) \otimes (I/I^2)$ has 8 generators and 64 relations; a basis of B/A is $\{e_{11}, e_{12}\}$ and the $2e_{ij}$ span I/I^2 . By inspection of the relations in (5) we see that the map $B/I \otimes I/I^2 \rightarrow \mathbb{F}_2$ sending $e_{ij} \otimes 2e_{kl}$ to $\delta_{il} + \delta_{jk}$ sends $A/I \otimes I/I^2$ and all the relations in (5) to zero, and

sends $e_{11} \otimes 2e_{12}$ to 1. Thus it induces a surjection $K_1(A, B, I) \rightarrow \mathbb{F}_2$. We claim that this is an isomorphism.

The relations for $(b, c, z) = (e_{11}, e_{11}, 2e_{11}), (e_{11}, e_{11}, 2e_{22}), (e_{11}, e_{12}, 2e_{12})$ and $(e_{11}, e_{21}, 2e_{21})$ in (5) yield the relations

$$0 = e_{11} \otimes 2e_{11} = e_{11} \otimes 2e_{22} = e_{12} \otimes 2e_{12} = e_{21} \otimes 2e_{21}.$$

The relations for $(e_{12}, e_{11}, 2e_{11}), (b, c, z) = (e_{11}, e_{21}, 2e_{11})$ and $(e_{12}, e_{12}, 2e_{21})$ in (5) yield the relations

$$0 = e_{11} \otimes 2e_{12} - e_{12} \otimes 2e_{11} = e_{21} \otimes 2e_{11} - e_{11} \otimes 2e_{21} = e_{12} \otimes 2e_{22} - e_{12} \otimes 2e_{11}.$$

This verifies the claim, proving that $K_1(A, B, 2B) \cong \mathbb{F}_2$.

Finally, the map $NK_1(A, B, 2B) \rightarrow NK_1(A, I)$ sends the class of $x^n e_{11} \otimes 2e_{12}$ to the class of the matrix $\begin{pmatrix} 1-2x^n e_{12} e_{11} & 2x^n e_{12} \\ -x^{2n} e_{11} e_{12} e_{11} & 1+2x^n e_{11} e_{12} \end{pmatrix} = \begin{pmatrix} 1 & x^n i(1+\tau) \\ 0 & 1+x^n i(1+\tau) \end{pmatrix}$, which is the class of $1 + x^n i(1 + \tau)$ in $NK_1(A)$. □

Remark. The elements $u = i + \tau$ and $v = i(1 + \tau)$ of A satisfy $u^2 = v^2 = 0$, and are distinct in $A/2A = \mathbb{F}_2[i, \tau]$. Hence the units $(1 + x^m u)(1 + x^n v)$ of $A[x]$ generate a subgroup of $NK_1(A)$ isomorphic to V^2 , which injects into $NK_1(\mathbb{F}_2[i, \tau]) \cong V^3$. (Since $\mathbb{F}_2[i, \tau] = \mathbb{F}_2[u, v]/(u^2, v^2)$, the other copy of V in $NK_1(\mathbb{F}_2[i, \tau])$ is the subgroup generated by all $(1 + x^n uv)$.)

Proposition 2.3. $NK_0(A) = 0$ and $NK_1(A) \cong V^2$ on the units $(1 + x^m u)(1 + x^n v)$.

The maps $A \rightarrow \mathbb{F}_2[i, \tau] \cong \mathbb{F}_2[\varepsilon_1, \varepsilon_2]$ and $\Omega_{\mathbb{F}_2[x]} \xrightarrow{\delta} NK_2(\mathbb{F}_2[\varepsilon_1, \varepsilon_2])$ sending x^n to $\langle x^{n-1} \varepsilon_1 \varepsilon_2, x \rangle$ induce a surjection $NK_2(A) \times \Omega_{\mathbb{F}_2[x]} \rightarrow NK_2(\mathbb{F}_2[\varepsilon_1, \varepsilon_2])$.

Proof. Consider the Mayer–Vietoris sequence of the square (6). Since $B = M_2(\mathbb{Z})$ and $B/I = M_2(\mathbb{F}_2)$ are regular rings, $NK_n(B) = NK_n(B/I) = 0$ and hence $NK_n(B, I) = 0$ for all n . We immediately get that $NK_n(A, B, I) \cong NK_n(A, I)$, that the Mayer–Vietoris sequence reduces to $NK_0(A) \cong NK_0(A/I) = 0$, and that there is an exact sequence

$$NK_2(A) \rightarrow NK_2(\mathbb{F}_2[\varepsilon_1]) \rightarrow NK_1(A, B, I) \rightarrow NK_1(A) \rightarrow NK_1(\mathbb{F}_2[\varepsilon_1]) \rightarrow 0.$$

By Lemma 2.2 and the remark preceding it, this yields the calculation of $NK_1(A)$.

Now $\pi: A \rightarrow \mathbb{F}_2[\varepsilon_1, \varepsilon_2]$ satisfies $\pi(u) = \varepsilon_1 + \varepsilon_2$, $\pi(v) = \varepsilon_1 + \varepsilon_1 \varepsilon_2$ and $\pi(uv) = \varepsilon_1 \varepsilon_2$, so we may write $\mathbb{F}_2[\varepsilon_1, \varepsilon_2] \cong \mathbb{F}_2[\bar{u}, \bar{v}]/(\bar{u}^2, \bar{v}^2)$. By [vdK71], the group $NK_2(\mathbb{F}_2[\bar{u}, \bar{v}])$ is isomorphic to the direct sum of $NK_2(\mathbb{F}_2[\bar{u}])$, $NK_2(\mathbb{F}_2[\bar{v}])$ and a group with the following generators:

$$\langle x^n \bar{u}, \bar{v} \rangle, \quad \langle x^n \bar{u} \bar{v}, \bar{u} \rangle, \quad \langle x^n \bar{u} \bar{v}, \bar{v} \rangle \quad \text{and} \quad \langle x^{n-1} \bar{u} \bar{v}, x \rangle.$$

Since $u^2 = v^2 = 0$ in A , all these symbols lift to Dennis–Stein symbols in $NK_2(A)$ except possibly the symbols $\langle x^{n-1}\bar{u}\bar{v}, x \rangle$. But these symbols are hit by the image of $\Omega_{\mathbb{F}_2[x]}$ under δ . □

Remark. $\delta: \Omega_{\mathbb{F}_2[x]} \rightarrow NK_2(\mathbb{F}_2[\varepsilon_1, \varepsilon_2])$ is a homomorphism by the Dennis–Stein identity $\langle f, x \rangle \langle g, x \rangle = \langle f + g - fgx, x \rangle$ with $fg = 0$; see [GL80, p. 184]. It is a morphism of \mathbb{F}_2 -Cartier modules since $V_m \langle x^{n-1}\varepsilon_1\varepsilon_2, x \rangle = m \langle x^{mn-1}\varepsilon_1\varepsilon_2, x \rangle = \delta(V_m(x^n))$ and (by [St80, 2.1])

$$F_m \langle x^{n-1}\varepsilon_1\varepsilon_2, x \rangle = \begin{cases} \langle x^{n/m-1}\varepsilon_1\varepsilon_2, x \rangle, & m \mid n \\ r \langle x^{n-1}(\varepsilon_1\varepsilon_2)^m, x \rangle - s \langle x^n(\varepsilon_1\varepsilon_2)^{m-1}, \varepsilon_1\varepsilon_2 \rangle = 0, & rm + sn = 1. \end{cases}$$

Our analysis of D_4 will involve the units of the ring $\mathbb{Z}/4[x][C_2]$.

Example 2.4. Consider the modular group ring $B = \mathbb{Z}/4[C_2] = \mathbb{Z}/4[e]/(e^2 - 2e)$, with $e = 1 - \tau$. The ideals $2eB$ of B and $eB/2eB$ of $B/2eB$ are isomorphic to \mathbb{F}_2 , so both $NK_1(B, 2e)$ and $NK_1(B/2e, e)$ are isomorphic to V and the group $NK_1(B, e)$, identified with the abelian group $(1 + xeB[x])^\times$, is a nontrivial extension:

$$0 \rightarrow V \rightarrow NK_1(B, e) \rightarrow V \rightarrow 0.$$

As an abelian group, $NK_1(B, e)$ is the direct sum of a countably infinite free $\mathbb{Z}/4$ -module on the $(1 + ex^m)$ ($m = 1, 2, \dots$) and a countably infinite free $\mathbb{Z}/2$ -module on the $(1 + 2ex^{2i-1})$ ($i = 1, 2, \dots$). As a module over the $\mathbb{Z}/4$ -Cartier algebra (generated by the operators V_m, F_m and homothety [2]), $NK_1(B, e)$ is cyclic on generator $u = 1 + ex$; $V_m(u) = 1 + ex^m$ and $V_m[2](u) = 1 + 2ex^m$.

Finally, we are in position to analyze $NK_0(\mathbb{Z}[D_4])$. The sharp exponent 4 for $NK_0(\mathbb{Z}[D_4])$ in Theorem 2.5 is a slight improvement on the bound in [CP02]. It is convenient to write D_4 as the semidirect product of C_4 (on σ) with the cyclic group $C_2 = \{1, \tau\}$, with relation $\tau\sigma\tau = \sigma^{-1}$.

Theorem 2.5. The group $NK_0(\mathbb{Z}[D_4])$ is isomorphic to the cyclic Cartier module $NK_1(\mathbb{Z}/4[C_2], 1 - \tau)$, described in Example 2.4. As a group, it is the direct sum of a countably infinite free $\mathbb{Z}/4$ -module and a countably infinite free $\mathbb{Z}/2$ -module.

Proof. We can map $\mathbb{Z}[D_4]$ to the twisted ring $A = \mathbb{Z}[i] \rtimes C_2$ occurring in (6) above, sending σ to i . Combining this with the natural surjection onto the subring $\mathbb{Z}[D_2]$ of $\mathbb{Z}[C_2] \times \mathbb{Z}[C_2]$, we get a ring map $\mathbb{Z}[D_4] \rightarrow A \times \mathbb{Z}[C_2] \times \mathbb{Z}[C_2]$. The ideal $I = (4, 2 - 2\sigma, \sigma^2 - 1)\mathbb{Z}[D_4]$ has $B_0 = \mathbb{Z}[D_4]/I = \mathbb{Z}/4[D_2]/(2 - 2\sigma)$, and is

isomorphic to the ideal $2A \times (4) \times (4)$ of $A \times \mathbb{Z}[C_2] \times \mathbb{Z}[C_2]$. Consider the following cartesian square:

$$\begin{array}{ccc}
 \mathbb{Z}[D_4] & \xrightarrow[\tau \mapsto (\tau, \tau, \tau)]{\sigma \mapsto (i, 1, -1)} & A \times \mathbb{Z}[C_2] \times \mathbb{Z}[C_2] \\
 \downarrow & & \downarrow \pi \\
 B_0 = \mathbb{Z}/4[D_2]/(2 - 2\sigma) & \xrightarrow{q = (q_0, q_+, q_-)} & \mathbb{F}_2[D_2] \times B \times B.
 \end{array} \tag{7}$$

The kernel of the split surjection $q_+ : B_0 \rightarrow B = \mathbb{Z}/4[C_2]$ is the 2-dimensional ideal $J = (1 - \sigma)B_0$. This implies that $NK_1(B_0) = NK_1(B) \oplus NK_1(B_0, J)$. Because $NK_1(\mathbb{Z}[C_2]) = NK_0(\mathbb{Z}[C_2]) = NK_0(A) = 0$ (by 2.3), the Mayer–Vietoris sequence associated to (7) ends

$$NK_1(A) \times NK_1(B_0, J) \xrightarrow{\eta} NK_1(\mathbb{F}_2[D_2]) \times NK_1(B) \rightarrow NK_0(\mathbb{Z}[D_4]) \rightarrow 0. \tag{8}$$

The displayed map η is given by the matrix $\begin{pmatrix} \pi & 0 \\ q_0 & q_- \end{pmatrix}$. It is easy to see that $NK_1(B_0, J)$ is isomorphic to V^2 on the terms $(1 + (1 - \sigma)x^m)$ and $(1 + (1 - \sigma)\tau x^n)$. An elementary calculation using the isomorphism $NK_1(A) \cong V^2$ of 2.3 shows that η is an injection, sending the module $NK_1(A) \times NK_1(B_0, J) \cong V^4$ isomorphically onto the subgroup $NK_1(\mathbb{F}_2[D_2]) \times NK_1(\mathbb{Z}/4)$. Since $NK_1(B) = NK_1(\mathbb{Z}/4) \oplus NK_1(B, eB)$, $e = 1 - \tau$, it follows that the induced map $NK_1(B, e) \rightarrow NK_0(\mathbb{Z}[D_4])$ is an isomorphism. □

To begin the calculation of $NK_1(\mathbb{Z}[D_4])$, we extend the Mayer–Vietoris sequence (8) associated to (7) to the left. This is possible by the following observation: since B_0 maps onto each of the three ring factors on the lower right of (7), the presentation (5) shows that the double relative K_1 obstruction vanishes. Because η is an injection in (8), the continuation of the Mayer–Vietoris sequence yields the exact sequence

$$(*) \xrightarrow{\begin{pmatrix} \pi & 0 \\ q_0 & q_- \end{pmatrix}} NK_2(\mathbb{F}_2[D_2]) \times NK_2(B) \rightarrow NK_1(\mathbb{Z}[D_4]) \rightarrow 0, \tag{9}$$

where (*) denotes $NK_2(A) \times NK_2(\mathbb{Z}[C_2])^2 \times NK_2(B_0, J)$.

Definition 2.6. The map $\cup[x] : NK_0(\mathbb{Z}[D_4]) \rightarrow NK_1(\mathbb{Z}[D_4])$ is obtained by composing the isomorphism $NK_1(B, e) \cong NK_0(\mathbb{Z}[D_4])$ of Theorem 2.5 with the canonical map $NK_2(B, e) \rightarrow NK_2(B) \rightarrow NK_1(\mathbb{Z}[D_4])$ of (9).

Remark. There is also a canonical map $NK_1(B, e) \rightarrow NK_2(B, e)$ sending the unit $1 - aex$ to $\langle ae, x \rangle$; the composition with $NK_2(B, e) \subset K_2(B[x, x^{-1}], e)$ is given by $1 - aex \mapsto \{1 - aex, x\}$ (multiplication by the class of x in $K_1(\mathbb{Z}[x, x^{-1}])$).

The analogous maps from $V \cong NK_1(B, 2e)$ and $V \cong NK_1(B/2eB, e)$ to $\Omega_{\mathbb{F}_2[x]} \cong NK_2(B, 2e)$ and $\Omega_{\mathbb{F}_2[x]} \cong NK_2(B/2eB, e)$ are compatible with the divided power map $[d]: V \rightarrow \Omega_{\mathbb{F}_2[x]}$ sending x^n to $x^{n-1} dx$. Note that $[d]$ is an isomorphism of abelian groups but is not a morphism of \mathbb{F}_2 -Cartier modules.

Theorem 2.7. *The map $\cup[x]: NK_0(\mathbb{Z}[D_4]) \rightarrow NK_1(\mathbb{Z}[D_4])$ in Definition 2.6 is a surjection. Hence the group $NK_1(\mathbb{Z}[D_4])$ has exponent 2 or 4, and there is a commutative diagram whose rows are exact:*

$$\begin{array}{ccccccc}
 0 & \longrightarrow & V & \longrightarrow & NK_0(\mathbb{Z}[D_4]) & \longrightarrow & V \longrightarrow 0 \\
 & & \cong \downarrow [d] & & \downarrow \cup[x] & & \cong \downarrow [d] \\
 & & \Omega_{\mathbb{F}_2[x]} & \longrightarrow & NK_1(\mathbb{Z}[D_4]) & \longrightarrow & \Omega_{\mathbb{F}_2[x]} \longrightarrow 0.
 \end{array}$$

Proof. A diagram chase on (9) shows that $NK_1(\mathbb{Z}[D_4])$ is an extension of the cokernel of $NK_2(\mathbb{Z}[C_2]) \times NK_2(B_0, J) \rightarrow NK_2(B)$ by a quotient of the cokernel of $NK_2(A) \rightarrow NK_2(\mathbb{F}_2[D_2])$. These cokernels are both $\Omega_{\mathbb{F}_2[x]}$, by Proposition 2.3 and Lemma 2.8 below, yielding the bottom row of the theorem. The map $\cup[x]$ sends the element corresponding to $1 - x^n ae \in NK_1(B, e)$ to the element corresponding to $\langle x^{n-1} ae, x \rangle \in NK_2(B, e)$, so the diagram in the theorem commutes by inspection. \square

Lemma 2.8. *The cokernel of the map $NK_2(\mathbb{Z}[C_2]) \times NK_2(B_0, J) \rightarrow NK_2(B)$ in (9) is $\Omega_{\mathbb{F}_2[x]}$, on symbols $\langle x^{n-1} e, x \rangle$.*

Proof. The kernel of the map $q_-: B_0 \rightarrow B$ is the ideal $J' = (1 + \sigma)B_0$. Because $J \cap J' = 0$ in B_0 , the double relative group $NK_2(B_0, J, J')$ is isomorphic to $\mathbb{F}_2[C_2][x]$ on symbols $\langle x^m(1 + \sigma), (1 - \sigma) \rangle$ and $\langle x^m \tau(1 + \sigma), (1 - \sigma) \rangle$ by [GL80], [Kou81]. Since $J \cong 2B$, we have an exact sequence

$$\mathbb{F}_2[C_2][x] \rightarrow NK_2(B_0, J) \xrightarrow{q_-} NK_2(B, 2B) \rightarrow 0. \tag{10}$$

Combining this with the ideal sequence for $2B \subset B$ shows that the cokernel of $NK_2(B_0, J) \rightarrow NK_2(B)$ is $NK_2(B/2B)$. Since $B/2B \cong \mathbb{F}_2[C_2]$, the lemma now follows from Lemma 1.2. \square

Inserting the calculations of Theorems 2.5 and 2.7 into Tables 6–7 in [LO], we obtain the following result.

Theorem 2.9. *Let Γ be one of the following hyperbolic 3-simplex reflection groups: $[(3, 4, 3, 6)]$, $[4, 3^{[3]}]$, $[4, 3, 6]$, $[(3^3, 4)]$, $[4, 3, 5]$, $[(3, 4)^{[2]}]$, $[(3, 4, 3, 5)]$. Then the lower algebraic K -theory of the groups Γ is given by the following table:*

Γ	$K_{-1} \neq 0$	$\tilde{K}_0 \neq 0$	$\text{Wh} \neq 0$
$[(3, 4, 3, 6)]$	\mathbb{Z}^3	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0$	Nil_1
$[4, 3^{[3]}]$	\mathbb{Z}^3	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$\text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$
$[4, 3, 6]$	\mathbb{Z}^4	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$\text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$
$[(3^3, 4)]$	\mathbb{Z}^2	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$\text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$
$[4, 3, 5]$	\mathbb{Z}^4	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$\mathbb{Z}^3 \oplus \text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$
$[(3, 4)^{[2]}]$	\mathbb{Z}^4	$(\mathbb{Z}/4)^4 \oplus 2\text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$2\text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$
$[(3, 4, 3, 5)]$	\mathbb{Z}^6	$(\mathbb{Z}/4)^2 \oplus \text{Nil}_0 \oplus \bigoplus_{\infty} \mathbb{Z}/2$	$\mathbb{Z}^3 \oplus \text{Nil}_1 \oplus \bigoplus_{\infty} \mathbb{Z}/2$

In this table, $\text{Nil}_0 = NK_0(\mathbb{Z}[D_4])$ is the direct sum of a countably infinite free $\mathbb{Z}/4$ -module and a countably infinite free $\mathbb{Z}/2$ -module, and $\text{Nil}_1 = NK_1(\mathbb{Z}[D_4])$ is a countably infinite torsion group of exponent 2 or 4.

References

- [Bas68] H. Bass, *Algebraic K-theory*. W. A. Benjamin, New York 1968. [Zbl 0174.30302](#)
[MR 0249491](#)
- [CP02] F. Connolly and S. Parasidis, On the exponent of the cokernel of the forget-control map on K_0 -groups. *Fund. Math.* **172** (2002), 201–216. [Zbl 0992.57022](#) [MR 1898685](#)
- [DW93] B. Dayton and C. Weibel, Module structures on the Hochschild and Cyclic Homology of Graded Rings. In *Algebraic K-theory and Algebraic Topology*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 407, Kluwer, Dordrecht 1993, 63–90. [Zbl 0900.16015](#)
[MR 1367293](#)
- [F87] F. T. Farrell, A remark on K_0 of crystallographic groups. *Topology Appl.* **26** (1987), 97–99. [Zbl 0624.18004](#) [MR 0893807](#)
- [GW83] S. Geller and C. Weibel, $K_1(A, B, I)$. *J. Reine Angew. Math.* **342** (1983), 12–34. [Zbl 0503.18009](#) [MR 0703484](#)
- [GL80] D. Guin-Waléry, and J-L. Loday, Obstruction à l’excision en K -théorie algébrique. In *Algebraic K-theory* (Evanston 1980), Lecture Notes in Math. 854, Springer-Verlag, Berlin 1981, 179–216. [Zbl 0461.18007](#) [MR 0618305](#)
- [Keu81] F. Keune, Doubly relative K -theory and the relative K_3 . *J. Pure Appl. Algebra* **20** (1981), 39–53. [Zbl 0473.18009](#) [MR 0596152](#)
- [LO] J. F. Lafont, and I. J. Ortiz, Lower algebraic K -theory of hyperbolic 3-simplex reflection groups. *Comment. Math. Helv.* **84** (2009), 297–337.

- [L05] W. Lück, K - and L -theory of the semi-direct product of the discrete 3-dimensional Heisenberg group by $\mathbb{Z}/4$. *Geom. Topol.* **9** (2005), 1639–1676 [Zbl 1073.19004](#)
[MR 2175154](#)
- [Mi71] J. Milnor, *Introduction to algebraic K-theory*. Ann. of Math. Stud. 72, Princeton University Press, Princeton, N.J., 1971. [Zbl 0237.18005](#) [MR 0349811](#)
- [St80] J. Stienstra, On K_2 and K_3 of truncated polynomial rings. In *Algebraic K-theory, Evanston 1980*, Lecture Notes in Math. 854, Springer-Verlag, Berlin 1981, 409–455. [Zbl 0463.13007](#) [MR 0618315](#)
- [vdK71] W. van der Kallen, Le K_2 des nombres duaux. *C. R. Acad. Sci. Paris Sér. A-B* **273** (1971), 1204–1207. [Zbl 0225.13006](#) [MR 0291158](#)
- [We80] C. A. Weibel, Mayer-Vietoris sequences and module structures on NK_* . In *Algebraic K-theory, Evanston 1980*, Lecture Notes in Math. 854, Springer-Verlag, Berlin 1981, 466–493. [Zbl 0487.18012](#) [MR 0618317](#)

Received February 17, 2008

Charles Weibel, Department of Mathematics, Rutgers University, Piscataway, NJ 08854, U.S.A.

E-mail: weibel@math.rutgers.edu