

# Note on a diophantine equation

Autor(en): **Williams, H.C.**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **25 (1970)**

Heft 6

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-27360>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

a distance. Can this be done with as few as  $O(n^{1/2})$  points; or with  $O(n^{1/3})$  points in one dimension?

Another open problem [1] is given any  $n$  points in the plane (not necessarily lattice points) [or in  $d$  dimensions], how many can one select so that the distances which are determined are all distinct? P. ERDÖS and R. K. GUY, Budapest

REFERENCES

[1] ERDÖS, P., *Néhány geometriai problémáról* (in Hungarian), *Mat. Lapok* 8, 86–92 (1957); *M. R.* 20, 6056 (1959).  
 [2] ERDÖS, P. and TURAN, P., *On a Problem of Sidon in Additive Number Theory and Some Related Problems*, *J. London Math. Soc.* 16, 212–215 (1941); *M. R.* 3, 270 (1942).  
 [3] HARDY, G. H. and WRIGHT, E. M., *Introduction to the Theory of Numbers*, 4th ed. (Oxford 1960).  
 [4] LANDAU, E., *Handbuch der Lehre von der Verteilung der Primzahlen* (Leipzig 1909), II, 643.  
 [5] LINDSTRÖM, B., *An Inequality for  $B_2$ -sequences*, *J. Combinatorial Theory* 6, 211–212 (1969).  
 [6] SINGER, J., *A Theorem in Finite Projective Geometry and Some Applications to Number Theory*, *Trans. Amer. Math. Soc.* 43, 377–385 (1938).

### Note on a Diophantine Equation

SCHINZEL and SIERPIŃSKI [1] have given the general solution of the diophantine equation

$$(x^2 - 1)(y^2 - 1) = \left[ \left( \frac{x - y}{2} \right)^2 - 1 \right]^2,$$

and SZYMICZEK [2] has given the general solution of

$$(x^2 - z^2)(y^2 - z^2) = \left[ \left( \frac{y - x}{2} \right)^2 - z^2 \right]^2.$$

The purpose of this paper is to obtain a complete solution of the diophantine equation

$$(x^2 + a)(y^2 + a) = \left[ a \left( \frac{y - x}{2b} \right)^2 + b^2 \right]^2, \tag{1}$$

where  $a$  and  $b$  are any two given integers.

Let  $X = x - y$ ,  $Y = x + y$ ; then  $X \equiv Y \pmod{2}$  and (1) becomes

$$b^4 (X^2 + 2XY + Y^2 + 4a)(X^2 - 2XY + Y^2 + 4a) = (aX^2 + 4b^4)^2.$$

This equation reduces to

$$b^4 ((Y^2 - X^2)^2 + 8a(Y^2 - X^2) + 16a^2) = (aX^2 + 4b^4)^2 - 16ab^4X^2$$

and we have

$$b^2 (Y^2 - X^2 + 4a) = \pm (aX^2 - 4b^4).$$

At this point it becomes necessary to find the complete solutions of

$$b^2 (Y^2 - X^2 + 4a) = + (aX^2 - 4b^4) \quad (2)$$

and

$$b^2 (Y^2 - X^2 + 4a) = - (aX^2 - 4b^4) . \quad (3)$$

We shall first solve (2).

Let  $(a, b^2) = \gamma^2 \delta$ , where  $\delta$  has no square factors; then  $a = \gamma^2 \delta \alpha$  and  $b = \gamma \delta \beta$ , where  $(\beta \delta, \alpha) = 1$ .

From (2), we must have

$$\delta \beta^2 \mid \alpha X^2$$

or, equivalently,  $X = \delta \beta Z$ , for some integer  $Z$ . On substituting this value for  $X$  and re-arranging the terms in (2), we obtain the equation

$$Y^2 - (\delta \beta^2 + \alpha) \delta Z^2 = -4 \gamma^2 (\delta^2 \beta^2 + \delta \alpha) . \quad (4)$$

If we let

$$c_1^2 d_1 = \delta \beta^2 + \alpha , \quad (5)$$

where  $d_1$  has no square factors, then  $(d_1, \delta) = 1$  and therefore,  $Y = c_1 d_1 \delta W$ , for some integer  $W$ . Equation (4) reduces to

$$Z^2 - d_1 \delta W^2 = 4 \gamma^2 . \quad (6)$$

Thus the problem of obtaining all solutions of (2) reduces to the solved problem of finding all solutions of (6). It may be shown similarly that if we let

$$c_2^2 d_2 = \delta \beta^2 - \alpha ,$$

where  $d_2$  has no square factor, then the problem of obtaining all solutions of (3) reduces to the problem of obtaining all solutions of

$$z^2 - d_2 \delta w^2 = -4 \gamma^2 .$$

It is evident that part of the solutions of (1) must be of the form

$$x = (\delta \beta Z + c_1 d_1 \delta W)/2 , \quad y = (c_1 d_1 \delta W - \delta \beta Z)/2 ,$$

where  $(Z, W)$  is a solution of (6). If  $2 \mid \delta$  or  $2 \mid d_1$ , it is clear that  $x$  and  $y$  will both be integers. If  $2 \nmid \delta$  and  $2 \nmid d_1$ , then, by (6),  $Z$  and  $W$  will be of the same parity. If they are both even, then  $x$  and  $y$  are integers; if they are both odd, we must have, in order that  $x$  and  $y$  be integers, that  $\beta \equiv c_1 \pmod{2}$ . But, from (5),

$$c_1 \equiv \beta + \alpha \pmod{2} ;$$

thus, it is necessary that  $\alpha \equiv 0 \pmod{2}$ .

Hence, if  $2 \mid (d_1 \delta \alpha)$ ,  $(c_1 d_1 \delta W \pm \delta \beta Z)/2$  are integers. If  $2 \nmid (d_1 \delta \alpha)$ , then  $(c_1 d_1 \delta W \pm \delta \beta Z)/2$  will be integers if and only if  $2 \mid W$  and  $2 \mid Z$ .

This same discussion may be applied to what must be the forms of the remaining solutions of (1), that is,

$$x = (\delta \beta z + c_2 d_2 \delta w)/2 , \quad y = (c_2 d_2 \delta w - \delta \beta z)/2 .$$

Thus we have proved the following

*Theorem.* If  $(a, b^2) = \gamma^2 \delta$ , where  $\delta$  has no square factors, and

$$c^2 d = \delta \beta^2 \pm \alpha,$$

where

$$\beta = \frac{b}{\gamma \delta},$$

and

$$\alpha = \frac{a}{\gamma^2 \delta},$$

and  $d$  has no square factors, then any solution of (1) must be of the form

$$x = (c d \delta W + \delta \beta Z)/2, \quad y = (c d \delta W - \delta \beta Z)/2,$$

where  $(Z, W)$  is a solution of

$$Z^2 - d \delta W^2 = \pm 4 \gamma^2$$

and  $2 \mid (\alpha \delta d)$ ; if  $2 \nmid (\alpha \delta d)$ , then any solution of (1) must be of the form

$$x = c d \delta W + \delta \beta Z, \quad y = c d \delta W - \delta \beta Z,$$

where  $(Z, W)$  is a solution of

$$Z^2 - d \delta W^2 = \pm \gamma^2.$$

This theorem has the following

*Corollary.* The equation

$$(x^2 + a)(y^2 + a) = (a z^2 + b^2)^2, \quad (7)$$

where  $a$  and  $b$  are integers, has integer solutions given by

$$x = b u + c d v, \quad y = b u - c d v, \quad z = u,$$

provided

$$b^2 \pm a = c^2 d,$$

where  $d$  has no square factors, and  $u$  and  $v$  are given by

$$u^2 - d v^2 = \pm 1.$$

It is not known to the author whether there are values of  $a$  and  $b$  for which the above result provides all the solutions to the equation (7); however, the equation (7) with  $a = b = 1$ , which was originally suggested to the author by L. J. MORDELL, has no solutions other than those predicted by the foregoing method for all integers  $x$  and  $y$ , where  $1 \leq x \leq 300$  and  $1 \leq y \leq x$ .

H. C. WILLIAMS, University of Manitoba

#### REFERENCES

- [1] A. SCHINZEL et W. SIERPIŃSKI, *Sur l'équation diophantienne*  $(x^2 - 1)(y^2 - 1) = [((y - x)/2)^2 - 1]^2$ , *El. Math.* 18, 132-133 (1963).  
 [2] K. SZYMICZEK, *On a Diophantine Equation*, *El. Math.* 22, 37-38 (1967).