

A Desarguesian dual for Nagel's middlespoint

Autor(en): **Eddy, Roland H.**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **44 (1989)**

Heft 3

PDF erstellt am: **22.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-41613>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A Desarguesian dual for Nagel's middlespoint

1. In a paper published in 1836, C. H. Nagel [4] defines the “middlespoint” (Mittenpunkt) of a given triangle ABC in the following manner:

Let S_A, S_B, S_C be the midpoints of BC, CA, AB respectively and I_a, I_b, I_c the centres of the excircles, then the lines $S_A I_a, S_B I_b, S_C I_c$ concur at M , the middlespoint of ABC ,

see also [1]. The name is probably derived from the fact that the point is constructed using “middles”, namely, *centres* of circles and *midpoints* of line segments. In this paper, we derive a dual (line) for this remarkable, but seemingly little known, point and show how this new line relates to some known geometry of the triangle.

2. Desargues's two-triangle theorem in the plane states that if triangles ABC and $A_1 B_1 C_1$ are perspective from a point L , they are perspective from a line l , i.e. if $AA_1 \cap BB_1 \cap CC_1 = L$, then $(AB \cap A_1 B_1) \cup (BC \cap B_1 C_1) \cup (CA \cap C_1 A_1) = l$. Clearly, the converse of this theorem is also its dual, hence, for purposes of this paper, we refer to L and l as “Desarguesian duals”. Also, we will have occasion to make reference to the special case when the triangle $A_1 B_1 C_1$ is inscribed in ABC , i.e. A_1 is on BC , etc. In this instance, L is called the trilinear pole of l and, dually, l is the trilinear polar of L , see [2].

3. In order to facilitate the arguments, we shall use a system of homogeneous coordinates called “trilinear” or “normal”. In order to avoid a possible confusion with trilinear poles and polars, we shall use the term “normal” throughout. In this system, the coordinates (x, y, z) of a point L in the plane of ABC are proportional to the signed distances d_a, d_b, d_c of L from the sides of the triangle of reference ABC , where, obviously, $A = (1, 0, 0)$, $B = (0, 1, 0)$, $C = (0, 0, 1)$. The distance d_a , for example, is positive if L and the unit point $I = (1, 1, 1)$, the incentre, are on the same side of $a = BC$ and negative otherwise. For instance, the excentre I_A of the excircle opposite vertex A has coordinates $(-1, 1, 1)$, or its projective equivalent, $(1, -1, -1)$.

We now derive the normal line coordinates of l , the trilinear polar of L , that is, if a line t has equation $ux + vy + wz = 0$ then $t = [u, v, w]$ is its normal representation. Also, for the remainder of the paper, we shall denote A_1 by A_L , etc., thus $A_L = (0, y, z)$, $B_L = (x, 0, z)$, $C_L = (x, y, 0)$ and, consequently, $A_L B_L = [yz, xz, -xy] = \left[\frac{1}{x}, \frac{1}{y}, -\frac{1}{z} \right]$, $xyz \neq 0$. Now $C_L'^{[1*]} = AB \cap A_L B_L = (x, -y, 0)$ and, similarly $B_L' = (-x, 0, z)$, $A_L' = (0, y, -z)$; hence the coordinates of $l = A_L' B_L' C_L'$ readily follow. Since this result does not seem to appear in the available literature, we state it as a proposition.

Proposition 1. If a point $L = (x, y, z)$ is in the plane of, but not incident with, a given triangle of reference ABC then, l , its trilinear polar, has coordinates $\left[\frac{1}{x}, \frac{1}{y}, \frac{1}{z} \right]$.

4. The coordinates of the middlespoint M are also readily obtained. The coordinates of the centroid S are easily seen to be of the form $\left(\frac{1}{\sin \alpha}, \frac{1}{\sin \beta}, \frac{1}{\sin \gamma} \right)$, where α, β, γ are

the measures of the vertex angles at A, B, C respectively, however, since $a = 2R \sin \alpha$, where R is the circumradius of ABC , it is more convenient to write $S = \left(\frac{1}{a}, \frac{1}{b}, \frac{1}{c}\right)$.

Now $I_a S_A = [(b-c), b, -c]$, $I_b S_B = [-a, (c-a), c]$, $I_c S_C = [a, -b, (a-b)]$, consequently, $M = (b+c-a, c+a-b, a+b-c)$. Again, we state this result in the form of a proposition.

Proposition 2. The normal coordinates of the midpoint M of a given triangle ABC are given by $M = (s-a, s-b, s-c)$, where $s = \frac{a+b+c}{2}$.

5. Since the triangles $I_A I_B I_C$ and $S_A S_B S_C$ are perspective from the midpoint M , they are, by Desargues's theorem, perspective from a line m , the Desarguesian dual of M , which we shall call the "middlesline" (mittenlinie) of ABC . Following the procedures above, it is an elementary exercise to show that the coordinates of m are $[a(s-a), b(s-b), c(s-c)]$, the details of which we leave as an exercise for the reader. We now state and prove a related result.

Proposition 3. The middlesline is the trilinear polar of the Gergonne point G of the given triangle ABC .

Proof: Since G_A, G_B, G_C are the points of contact of the incircle with the sides of ABC , $BG_A = s-b$ and $G_A C = s-c$, hence $G_A = (0, c(s-c), b(s-b))$ with similiar expressions for G_B, G_C . It now follows that the coordinates of G are $\left(\frac{1}{a(s-a)}, \frac{1}{b(s-b)}, \frac{1}{c(s-c)}\right)$ and by proposition 1, its trilinear polar is $[a(s-a), b(s-b), c(s-c)] = m$ as claimed.

Roland H. Eddy, Memorial University of Newfoundland,
St. John's, Newfoundland, Canada

REFERENCES

- 1 Baptist P.: Über Nagelsche Punktepaare, *Mathematische Semesterberichte* 1, 118–126 (1988).
- 2 Coxeter H. S. M.: *The Real Projective Plane*, 2nd ed., Cambridge University Press, 1961.
- 3 Johnson R. A.: *Advanced Euclidean Geometry (Modern Geometry)*, Dover, NY (1969).
- 4 Nagel C. H. v.: *Untersuchungen über die wichtigsten zum Dreiecke gehörenden Kreise*, Leipzig (1836).
- 5 Sommerville D. M. Y.: *Analytical Conics*, G. Bell and Sons Ltd., London (1956).
- 6 Springer C. E.: *Geometry and Analysis of Projective Spaces*, W. H. Freeman, San Francisco (1964).

NOTE

[1*] The "prime" notation is particularly useful here since C_L and C'_L are related by the harmonic conjugacy involution, see [2].