

A short combinatorial proof of derangement identity

Autor(en): **Martinjak, Ivica / Stani, Dajana**

Objektyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **73 (2018)**

Heft 1

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-760276>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A short combinatorial proof of derangement identity

Ivica Martinjak and Dajana Stanić

Ivica Martinjak was awarded his doctoral degree from the University of Zagreb. He is currently employed at the Croatian Academy of Science and Art as postdoctoral researcher and holds a position of Assistant Professor of Mathematics at the University of Osijek. His main fields of research are algebraic and enumerative combinatorics, combinatorial geometry and combinatorial number theory.

Dajana Stanić is a final year graduate student in mathematics at the Department of Mathematics, University of Osijek. In 2014 she finished the undergraduate programme in mathematics at the same university with a thesis in number theory. Her mathematical interests are in financial mathematics, statistics and probabilistic number theory.

1 Introduction

Having a permutation $\sigma \in S_n$, $\sigma : [n] \rightarrow [n]$ where $[n] := \{1, 2, \dots, n\}$, it is said that $k \in [n]$ is a *fixed point* if it is mapped to itself, $\sigma(k) = k$. Permutations without fixed points are of particular interest and are usually called *derangements*. We let D_n denote the number of derangements of the set $[n]$, $D_n = |S_n^{(0)}|$,

$$S_n^{(0)} := \{\sigma \in S_n : \sigma(k) \neq k, k = 1, \dots, n\}.$$

Derangements are usually introduced in the context of the inclusion-exclusion principle

Die Subfakultät $!n = n! \sum_{k=0}^n \frac{(-1)^k}{k!} = \left\lfloor \frac{n!+1}{e} \right\rfloor$ gibt die Anzahl der Derangements, d.h. der fixpunktfreien Permutationen von n unterscheidbaren Objekten an. Die Anzahl der Permutationen dieser Objekte mit genau r Fixpunkten wird als Rencontres-Zahl bezeichnet. Es gibt eine Vielzahl interessanter kombinatorischer Identitäten, bei denen die Subfakultät und Rencontres-Zahlen eine Rolle spielen. Am bekanntesten ist vermutlich das Inklusions-Exklusionsprinzip. In der vorliegenden Arbeit betrachten die Autoren gewichtete Summen von Subfakultäten. Diese lassen sich zwar auch algebraisch beweisen, hier werden jedoch elegante kombinatorische Abzählargumente für die Herleitung benützt.

[3, 5, 10], since this principle is used to provide an interpretation of D_n as a *subfactorial*,

$$D_n = n! \sum_{k=0}^n \frac{(-1)^k}{k!}. \quad (1)$$

The numbers $D_0, D_1, D_2, \dots, D_n, \dots$ form the recursive sequence $(D_n)_{n \geq 0}$ defined by the recurrence formula

$$D_n = (n-1)(D_{n-1} + D_{n-2}) \quad (2)$$

and initial terms $D_0 = 1, D_1 = 0$ [7]. There is a counting argument to prove this. Let the number k be mapped by σ to the number $j, j = 1, \dots, k-1, k+1, \dots, n$. Note that there are $(n-1)$ such permutations σ . Now, we separate the set of permutations σ into two disjoint sets \mathcal{A} and \mathcal{B} , such that

$$\begin{aligned} \mathcal{A} &:= \{\sigma \in S_n^{(0)} : \sigma(j) \neq k, \sigma(k) = j\} \\ \mathcal{B} &:= \{\sigma \in S_n^{(0)} : \sigma(j) = k, \sigma(k) = j\}. \end{aligned}$$

This means that

$$D_n = (n-1)(|\mathcal{A}| + |\mathcal{B}|).$$

The set \mathcal{A} counts D_{n-1} elements while the set \mathcal{B} counts D_{n-2} elements. The fact that the number k in this reasoning is chosen without loss of generality, completes the proof of (2). There is another recurrence for the sequence $(D_n)_{n \geq 0}$,

$$D_n = nD_{n-1} + (-1)^n. \quad (3)$$

Namely, set $\delta_n = D_n - nD_{n-1}$ for every $n \geq 1$. Then $\delta_1 = -1$, and formula (2) implies that one has for every $n \geq 2$

$$\delta_n = D_n - nD_{n-1} = (n-1)D_{n-2} - D_{n-1} = -\delta_{n-1},$$

hence one gets immediately $\delta_n = (-1)^n$, which proves (3).

When we iteratively apply recurrence (3) to the derangement number on the r.h.s. of this relation we get

$$nD_{n-1} + (-1)^n = n[(n-1)D_{n-2} + (-1)^{n-1}] + (-1)^n$$

which finally results with

$$n(n-1)(n-2) \cdots 3(-1)^2 + n(n-1)(n-2) \cdots 4(-1)^3 + \cdots + (-1)^n. \quad (4)$$

on the r.h.s. of (3), which completes the proof of (1).

A few identities for the sequence $(D_n)_{n \geq 0}$ are known [4, 6, 8]. In [4] Deutsche and Elizalde give a nice identity

$$D_n = \sum_{k=2}^n (k-1) \binom{n}{k} D_{n-k}. \quad (5)$$

Recently, Bhatnagar presents families of identities for some sequences including the shifted derangement numbers [1], deriving it using an Euler identity [2]. In what follows we demonstrate a combinatorial proof for that derangement identity, with weighted sum.

2 A pair of weighted sums for derangements

We define the *rencontres number* $D_n(r)$ as the number of permutations $\sigma \in S_n$ having exactly r fixed points. Thus, $D_n(0) = D_n$. For a given $r \in \mathbb{N}$, we define the sequence $D_0(r), D_1(r), \dots, D_n(r), \dots$, denoted by $(D_n(r))_{n \geq r}$.

Applying an analogue counting argument that we used when proving relation (2), one can represent rencontres numbers by the derangement numbers,

$$D_n(r) = \binom{n}{r} D_{n-r}. \quad (6)$$

On the other hand, relation (6) follows immediately from the fact that fixed points here are r -combinations over the set of n elements.

A few other notable properties of the rencontres numbers are also known. It follows from (3) that $D_n - D_n(1) = (-1)^n$ for every $n \geq 1$. According to the definition of rencontres numbers, the sum of the n th row in the array of numbers $(D_n(r))_{n \geq r}$ is equal to $n!$,

$$n! = \sum_{k=0}^n D_n(k). \quad (7)$$

Moreover, identity (6) shows that D_n can be interpreted as a weighted sum of rencontres numbers in the n th row of the array, by means of relation (5),

$$D_n = \sum_{k=2}^n (k-1) D_n(k). \quad (8)$$

The number $D_n/(n-1)$ is also a weighted sum of previous consecutive derangement numbers. For example, $24 + 12D_2 + 4D_3 + D_4 = \frac{D_6}{5}$. In general we have

$$n! + \sum_{k=2}^n \frac{n!}{k!} D_k = \frac{D_{n+2}}{n+1}, \quad (9)$$

as follows from Theorem 1.

Theorem 1. For $n \in \mathbb{N}$ and the sequence of derangement numbers $(D_n)_{n \geq 0}$ we have

$$1 + \sum_{k=1}^n \frac{D_k}{k!} = \frac{D_{n+2}}{(n+1)!}. \quad (10)$$

Proof. Within a derangement σ , the number k , $k = 1, \dots, n$ can be mapped to any j , $j = 1, \dots, k-1, k+1, \dots, n$. We let \mathcal{A}_n denote the set of derangements with $\sigma(k) = j$, where $j \neq k$,

$$\mathcal{A}_n := \{\sigma \in S_n^{(0)} : \sigma(k) = j\}.$$

Obviously, cardinality of the set \mathcal{A}_n is independent of j , $j \neq k$. More precisely,

$$|\mathcal{A}_n| = \frac{D_n}{n-1}.$$

Furthermore, we separate the set \mathcal{A}_n into two disjoint sets of derangements, sets \mathcal{B}_n and \mathcal{C}_n ,

$$\begin{aligned}\mathcal{B}_n &:= \{\sigma \in \mathcal{A}_n : \sigma(j) = k\} \\ \mathcal{C}_n &:= \{\sigma \in \mathcal{A}_n : \sigma(j) \neq k\}.\end{aligned}$$

Obviously, the set \mathcal{B}_n counts D_{n-2} elements. For derangements in \mathcal{C}_n there are now $(n-2)$ equivalent ways to map j (excluding j and k), as Figure 1 illustrates. Thus, we have

$$|\mathcal{C}_n| = (n-2)|\mathcal{A}_{n-1}|,$$

which gives the recurrence relation

$$|\mathcal{A}_n| = D_{n-2} + (n-2)|\mathcal{A}_{n-1}|. \quad (11)$$

After repeating usage of (11) we get identity (9) which completes the proof. \square




Figure 1 In case of derangements in the set \mathcal{C}_n there are $(n-2)$ equivalent ways to map j .

In order to prove Theorem 1 algebraically, we apply recurrence (2) to get

$$\begin{aligned}\frac{D_{n+2}}{(n+1)!} &= \frac{(n+1)(D_{n+1} + D_n)}{(n+1)!} = \frac{D_{n+1}}{n!} + \frac{D_n}{n!} \\ &= \frac{n(D_n + D_{n-1})}{n!} + \frac{D_n}{n!} = \frac{D_n}{(n-1)!} + \frac{D_{n-1}}{(n-1)!} + \frac{D_n}{n!} \\ &= 1 + \frac{D_1}{1!} + \dots + \frac{D_n}{n!} = 1 + \sum_{k=1}^n \frac{D_k}{k!}.\end{aligned}$$

Theorem 2. For $n \in \mathbb{N}$ and the sequence of derangement numbers $(D_n)_{n \geq 0}$ we have

$$1 + \sum_{k=1}^n \frac{(-1)^k D_{k+3}}{k+2} = (-1)^n D_{n+2}. \quad (12)$$

Proof. By applying recurrence (2) we have

$$\begin{aligned}\sum_{k=0}^n \frac{(-1)^k D_{k+3}}{k+2} &= \frac{2(D_2 + D_1)}{2} - \frac{3(D_3 + D_2)}{3} + \dots + (-1)^n \frac{(n+2)(D_{n+2} + D_{n+1})}{n+2} \\ &= (D_2 + D_1) - (D_3 + D_2) + \dots + (-1)^n (D_{n+2} + D_{n+1}) \\ &= (-1)^n D_{n+2}\end{aligned}$$

which completes the proof. \square

Once having Theorem 1, substitution of (6) in identity (10) gives the generalization (13).

$$1 + \sum_{k=1}^n \frac{D_{k+r}(r)}{k! \binom{k+r}{r}} = \frac{D_{n+r+2}(r)}{(n+1)! \binom{n+r+2}{r}}. \quad (13)$$

The identity (14) follows by substitution of (6) in (12),

$$1 + \sum_{k=1}^n \frac{(-1)^k D_{k+r+3}(r)}{(k+2) \binom{k+r+3}{r}} = \frac{(-1)^n D_{n+r+2}(r)}{\binom{n+r+2}{r}}. \quad (14)$$

Note that the terms in identity (14) are always integers, which can be seen as a consequence of recurrence relation (2).

Acknowledgement

The authors thank the anonymous referee for valuable suggestions that improved the final version of the paper.

References

- [1] G. Bhatnagar, Analogues of a Fibonacci–Lucas Identity, *Fibonacci Quart.* 54 (2) (2016), 166–171.
- [2] G. Bhatnagar, In praise of an elementary identity of Euler, *Electronic J. Combinatorics*, 18 (2), #P13, (2011) 44 pp.
- [3] M. Bona, *A Walk Through Combinatorics*, World Scientific, 2011.
- [4] E. Deutsch, S. Elizalde, The largest and the smallest fixed points of permutations, *European J. Combin.* 31 (2010), 1404–1409.
- [5] R. Graham, D. Knuth, O. Patashnik, *Concrete Mathematics*, Addison-Wesley, 1994.
- [6] M. Hassani, Derangements and Applications, *J. Integer Seq.* 6 (2003), article 03.1.2.
- [7] The On-line Encyclopedia of Integer Sequences, A000166, oeis.org
- [8] M. Petkovšek, H.S. Wilf, D. Zeilberger, *A = B*, Taylor and Francis, 1996.
- [9] E. Sandifer, *How Euler Did It*, Mathematical Association of America, 2007.
- [10] R. Stanley, *Enumerative Combinatorics, Volume 1*, Cambridge University Press, 2011.

Ivica Martinjak
 Faculty of Science
 University of Zagreb
 Bijenička cesta 32
 HR-10000 Zagreb, Croatia
 e-mail: imartinjak@phy.hr

Dajana Stanić
 Department of Mathematics
 University of Osijek
 Trg Ljudevita Gaja 6
 HR-31000 Osijek, Croatia
 e-mail: dstanic@mathos.hr