

**Zeitschrift:** Entomologica Basiliensia  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 14 (1991)

**Artikel:** Description of a new Aphaonus species (Coleoptera, Carabidae)  
**Autor:** Zamotajlov, A. S.  
**DOI:** <https://doi.org/10.5169/seals-980589>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 03.02.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Description of a new *Aphaonus* species (Coleoptera, Carabidae)

by A.S. Zamotajlov

**Abstract:** A new carabid beetles species, *Aphaonus miroshnikovi* n. sp. from Abkhazia, Caucasus, is described.

**Key words:** Coleoptera Carabidae – Caucasus – *Aphaonus* – description – new species.

According to KRYZHANOVSKIJ & ABDURAKHMANOV (1983), the genus *Aphaonus* Reitt. comprises now 6 species, distributed in the western part of the Caucasus Major. Below is given a description of one further species of the genus.

### ***Aphaonus miroshnikovi* n. sp.**

Figs. 1 – 5.

Habitus (Fig. 1). Length 15.6 – 17.2 mm. Width 3.9 – 4.4 mm. Body black, polished.

Head broad, frontal furrows prominent, evenly divergent posteriad, without foveae at the middle, genae faintly prominent, neck broad, posttemporal tubercles rather small, eyes prominent, about two-thirds times as long as genae; mentum tooth rather short, bifid.

Pronotum longitudinal, 1.00 – 1.15 times as long as wide, moderately convex, strongly contracted posteriorly, widest at about apical third; apex somewhat emarginate, lateral sides narrowly reflexed, moderately arcuate in front, very weakly sinuated behind, base sinuated, hind angles almost rectangular, prominent, extended laterad; anterior transverse impression shallow, although more or less prominent, median line deep, not reaching both extremities, basal foveae rather deep; lateral margins with 1 – 3 anterior marginal setae (2 specimens have 1 seta on one side and 1 on another, 7 specimens – 2 on one side and 1 on another, and 1 specimen – 3 on one side and 2 on another) and 1 posterior one at both sides.

Elytra oblong-oval, rather narrow, 1.91 – 2.10 times as long as wide, sides subparallel, shoulders rounded but prominent, forming obtuse angles; striae moderately prominent, scutellary striole distinct, interspaces subconvex; interspace 3 with 1 pore adjoining stria 2 at two-thirds, marginal series sharply interrupted, of two groups of pores, basal one composed of 5 – 6 pores and apical one of 4 – 6 pores.




Fig. 1: *Aphaonus miroshnikovi* n. sp., general view.

Figs 2-5: *Aphaonus miroshnikovi* n. sp.: 2, aedeagus, left lateral view. 3, idem, apex in dorsal view. 4, left paramere, 5, right paramere. Scale: 3 mm.

Metatrochantera about three-fourths times as long as metafemora in males and about two-thirds in females. Ventral side of tarsal segment 5 absolutely not pubescent.

Aedeagus (Figs 2,3) rather weakly bending, of the normal for the bulk of known species (except for *A. koenigianus* Tschitsch. and *A. tichonis* Sem.) shape, with more or less prominent protuberance at ventral side. Left paramere (Fig. 4) wide, right one (Fig. 5) fairly slender, curving, both normal in form and size.

Holotype: ♂ (ZIL = Zoological Institute Acad. Sci., Leningrad), Caucasus, Abkhazia, Bzybian Mt. Range, environs of Khuap, 1500 m. 14. V – 8. VII. 1990, A. Zamotajlov & A. Miroshnikov, Paratypes: 2 ♂ & 1 ♂ (NHMB = Natural History Museum, Basel); 5 ♂ 3 ♂ (in the author's coll.), same locality, together with holotype.

The new species resembles habitually smaller specimens of *A. compressus* Rost or, especially, larger ones of *A. cylindriformis* Reitt., although it is somewhat more robust, shape and proportions of pronotum are closer to those of *A. pseudopercus* Reitt., ventral side of the ultimate tarsal segments always without even minute pubescence, lateral margins of pronotum usually with more than 1 setiferous pores in apical half (at least at one side). Judging from the structure of aedeagus, the new species is closely related to species of both sections I and III after SEMENOW (1899). From *A. arcanus* (Kurn.), which also occurs at the Bzybian Mt. Range, it differs in many characters (KURNAKOV, 1962 & KRYZHANOVSKIJ, 1989).

Type specimens were collected in the beech forest.

This species is named after my friend and colleague Dr. A. I. Miroshnikov of Krasnodar.

## References

- KRYZHANOVSKIJ, O. L. (1989): *New and little known species of the tribe Pterostichini (Coleoptera, Carabidae) from the Caucasus*. Proc. Zool. Inst. Leningrad 208: 14 – 22 [in Russian].
- KRYZHANOVSKIJ, O. L. & ABDURAKHMANOV G. M. (1983): *New and little known species of the supertribe Pterostichitae (Coleoptera, Carabidae) from the Caucasus*. Entomol. obozr. 62 (3): 529 – 537 [in Russian].
- KURNAKOV, V. N. (1962): *Contribution à la faune des Carabiques du Caucase. IV. Nouveaux Carabiques du Caucase*. Rev. Franç. d'Entomol. 29 (1): 32 – 43.
- SEmenow, A. (1899): *Recensio monographica specierum subgeneris Aphaonus Rtt.* (Coleoptera, Carabidae). Horae Soc. entomol. Ross. 32: 478 – 503.

Author's address:  
Dr. A. S. Zamotajlov  
ul. Kalinina, 13, korp. 43, kv. 27  
Krasnodar, 350044, USSR