

**Zeitschrift:** Entomologica Basiliensia et Collectionis Frey  
**Herausgeber:** Naturhistorisches Museum Basel, Entomologische Sammlungen  
**Band:** 30 (2008)

**Artikel:** A contribution to the Dermestidae (Coleoptera) from the Natural History Museum Basel  
**Autor:** Háva, Jíí  
**DOI:** <https://doi.org/10.5169/seals-981048>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 08.02.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A contribution to the Dermestidae (Coleoptera) from the Natural History Museum Basel

by Jiří Háva

**Abstract.** *Dermestes* (*Dermestes*) *amoenus* sp.nov. from China: Sichuan and Hubei is described, illustrated and compared with related species. The following new faunistic records are presented: *Dermestes* (*Dermestinus*) *nan* Háva, 2002 (Thailand: Lansang); *Thaumaglossa rufocapillata* Redtenbacher, 1867 (Philippines); *Thaumaglossa hilleri* Reitter, 1881 (India: Meghalaya, Sikkim); *Thaumaglossa wittmeri* Háva, 2006 (India: Sikkim); *Evorinea indica* (Arrow, 1915) (India: Sikkim, Uttar Pradesh); *Orphinus hartmanni* Háva, 2001 (India: Meghalaya).

**Key words.** Coleoptera – Dermestidae – taxonomy – distribution – new species – Palearctic region – Oriental region

### Introduction

The Dermestidae is a family of beetles that contains about 1300 species and subspecies worldwide (HÁVA 2003, 2008). In the course of determination of material belonging to the collection of Natural History Museum, Basel, Switzerland, a new species and 8 new records of Dermestidae were revealed.

### Materials and Methods

The sizes of beetles and of their body parts can be useful in species recognition and thus the following measurements were made:

total length (TL) . . . . . linear distance from anterior margin of pronotum  
to apex of elytra  
pronotal length (PL) . . . . . maximum length measured from anterior margin  
to posterior margin  
pronotal width (PW) . . . . . maximum linear transverse distance  
elytral length (EL) . . . . . linear distance from shoulder to apex of elytron  
elytral width (EW) . . . . . maximum linear transverse distance

The following abbreviations refer to the collections in which the examined material is deposited:

JHAC . . . . . Private Entomological Laboratory & Collection, Jiří Háva,  
Prague, Czech Republic  
NHMB . . . . . Naturhistorisches Museum, Basel, Switzerland

## SUBFAMILY DERMESTINAE

## Tribe Dermestini

*Dermestes (Dermestes) amoenus* sp.nov.

**Type material.** Holotype (female): “China, Sichuan, Mt. Emei, 600–1050 m, 5–19.v.1989, Lad. Bocák lgt.” (NHMB). Paratype (1 female): “China, Hubei, Dabie Shan, 31.1N 115.8E, Wujiashan forest park, 22–23.v.2005” (JHAC). Specimens of the species described here are provided with a red, printed label with text as follows: “HOLOTYPE [or PARATYPE, as appropriate] *Dermestes (Dermestes) amoenus* sp.n. Jiří Háva det. 2008”. Male unknown.

**Description of holotype.** Whole body brown-black, elongated, slightly convex.

Head with long, brown-yellow pubescence, punctation dense, maxillary and labial palpi brown. Antennae brown, with eleven antennomeres. The final three antennomeres form a brown antennal club covered in brown pubescence.

Pronotum more closely punctured and more densely setaceous than head. Pronotal disc moderately convex, its pubescence long, made up of brown-yellow setae, hind margins of pronotum with long, yellow setae. Two small depressions near basal margin of pronotum. Scutellum triangular, covered in long, brown pubescence.

Elytra densely and strongly punctured; each elytron with 10 striae; posterior margin near epipleuron with long, dense, brown-yellow pubescence; apices of elytra gradually rounded, with dense, short, brown pubescence. Legs black-brown with yellow pubescence.

Ventral part of body densely covered with brown-yellow pubescence. Prosternum with long, brown pubescence. All sternites brown and with short, dense, yellow pubescence. First visible abdominal sternite with distinct lateral depressions (Fig. 1). Sternites III and IV without median tuft of yellow setae; sternites I–V with small latero-median depression.


Body measurements (mm): TL 8.7, PL 1.8, PW 3.1, EL 7.1, EW 3.3.

Variability. Paratype female. Body measurements (mm): TL 9.1, PL 2.1, PW 3.3, EL 7.3, EW 3.5. Whole body more dark brown, pubescence on dorsal and ventral surfaces more brown.

**Distribution.** China: Sichuan, Hubei.

**Derivatio nominis.** Latin adjective “*amoenus*” (= attractive, beautiful).

**Differential diagnosis.** *Dermestes (Dermestes) amoenus* sp.nov. belongs to the *D. bicolor* species group defined by KALÍK & OHBAYASHI (1982); this group, part of the nominotypical subgenus, differs from other groups in the very visible lateral striation on elytra; the group of eight species presented is distributed in the Holarctic region, southern Africa and Brazil; the new species very similar in habitus to *D. (D.) laosensis* Háva, 2004 (Laos) and *D. (D.) schneideri* Háva, 2002 (Mongolia) but differs from them in the form of the distinct lateral depressions of the first visible abdominal sternite (Figs 2, 3).



**Figs 1–3.** Lateral margin of the first visible abdominal sternite with distinct lateral depressions: 1, *Dermestes (Dermestes) amoenus* sp.nov.; 2, *D. (D.) laosensis* Háva, 2004; 3, *D. (D.) schneideri* Háva, 2002.

### *Dermestes (Dermestinus) nan* Háva, 2002

**Material examined.** “Thailand, Lansang n.p. 500 m, 16°48′N 98°57′E, 18–24.iv.1991, David Král lgt.”, 4 exx, J. Háva det. (3 NHMB, 1 JHAC).

**Distribution.** Species known from Thailand (Chiang Mai and Nan), new for Lansang.

## SUBFAMILY TRINODINAE

### Tribe Trinodini

### *Evorinea indica* (Arrow, 1915)

**Material examined.** “India, Darjeeling D., Bhakta B., Tista, 200 m, 26.iv.1985” 1 ex., J. Háva det., (NHMB); “India U.P., Barkot, 1100–1200 m, 5–12.vi.1981, M. Brancucci leg.” 1 ex., J. Háva det. (NHMB).

**Distribution.** Species known from China (Guangdong), India (Madhya Pradesh, Maharashtra), Japan and Taiwan. New records for India: Sikkim, Uttar Pradesh.

## SUBFAMILY MEGATOMINAE

## Tribe Megatomini

*Orphinus hartmanni* Háva, 2001

**Material examined.** “NE India, Meghalaya, 3 km E Tura, 1150 m, 25°30'N 90°14'E, 4.v.1999, Dembický & Pacholátko leg.” 1 ex., J. Háva det. (NHMB).

**Distribution.** Species known from Bhutan, India, Sikkim and Nepal. New record for India: Meghalaya.

*Thaumaglossa hilleri* Reitter, 1881

**Material examined.** “NE India, Meghalaya, 3 km E Tura, 1150 m, 25°30'N 90°14'E, 4.v.1999, Dembický & Pacholátko leg.” 1 male, J. Háva det. (NHMB); “India, Bhakta B., Darjeeling D., Sakyong, 1000 m, 10–11.iv.1983” 1 female, J. Háva det. (NHMB).

**Distribution.** Species known from China, India (Uttar Pradesh), Nepal, Japan, Philippines and Taiwan. New records for India: Meghalaya, Sikkim.

*Thaumaglossa rufocapillata* Redtenbacher, 1867

**Material examined.** “Philippines, 1600 m, Mindanao, 30 km W of Maramag, 28–30.dec.1990, Bolm lgt.” 2 exx, J. Háva det. (NHMB).

**Distribution.** Species known from Germany (intr.), Netherlands (intr.), Cameroon, Congo, Kenya, Madagascar, Nigeria, Tanzania, Zimbabwe, China, India, Indonesia, Japan, Korea, Laos, Malaysia, Nepal, Taiwan, Thailand and Vietnam. New record for Philippines.

*Thaumaglossa wittmeri* Háva, 2006

**Material examined.** “India, Darjeeling D., Bhakta B., Sindepung, 23.iv.1987” 1 ex., J. Háva det. (NHMB); “India, Bhakta B., Darjeeling D., Sakyong, 1000 m, 10–11.iv.1983” 4 exx, J. Háva det. (NHMB).

**Distribution.** Species known from Nepal. New record for India: Sikkim.

## Acknowledgements

I am very much obliged to Dr. Michel Brancucci (NHMB) for the opportunity to study this interesting material, for his valuable comments and for reading the manuscript.

### References

- HÁVA J. (2003): *World Catalogue of the Dermestidae (Coleoptera)*. Studie a zprávy Okresního muzea Praha-východ, **Supplementum 1**: 1–196.
- HÁVA J. (2008): *World Catalogue of Dermestidae*. <http://www.dermestidae.wz.cz>.
- KALÍK V. & OHBAYASHI N. (1982): *A new Dermestid Beetle from Asia (Coleoptera)*. *Entomol. Rev. Japan* **37**: 65–67.

#### Author's address:

Jiří Háva  
Private Entomological Laboratory and Collection  
Únětice u Prahy 37  
CZ-252 62 Praha-západ  
CZECH REPUBLIC  
E-mail: [jh.dermestidae@volny.cz](mailto:jh.dermestidae@volny.cz)

