

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 8 (1962)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** SURVEY OF COBORDISM THEORY  
**Autor:** Milnor, J.

### Bibliographie

**DOI:** <https://doi.org/10.5169/seals-37949>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 18.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$\Omega_k = H_k(\mathcal{D}_o)$  to  $H_k(\mathcal{C}_o)$ . Thom, Rohlin and Švarč have shown that Pontrjagin classes can be defined for combinatorial manifolds. Therefore we have:

**THEOREM 3'.** — *The homomorphism  $\Omega_k \rightarrow H_k(\mathcal{C}_o)$  has kernel zero.*

However examples show that this homomorphism is not onto. The reader is referred to [13, 18].

Another interesting possibility would be to look at the class of compact homology manifolds.

Returning to the differentiable case, interesting cobordism groups can be obtained by restricting the connectivities of the manifolds involved. As an extreme case we can consider only differentiable manifolds which are either homotopy spheres or homotopy cells. The resulting cobordism groups are closely related to the problem of classifying differentiable structures on spheres. The reader is referred to Milnor [8] and Smale [14].

As a final, quite different, example consider differentiable imbeddings of the circle  $S^1$  in the 3-sphere  $S^3$ . Such an object (a knot) is said to *bound* if it can be extended to a differentiable imbedding of the disk  $D^2$  in the disk  $D^4$ . The resulting cobordism group has been studied by Fox and Milnor [5]. This group is not finitely generated.

#### REFERENCES

- [1] ATIYAH, M., Bordism and cobordism, Proc. Cambridge Phil. Soc., 57 (1961), 200-208.
- [2] AVERBUH, B. G., Algebraic structure of internal homology groups. Doklady Akad. Nauk S.S.S.R., 125 (1959), 11-14.
- [3] BROWN, M., Locally flat embeddings of topological manifolds. *A.M.S. Notices*, 7 (1960), 939-940.
- [4] DOLD, A., Erzeugende der Thomschen Algebra *N. Math. Zeits.*, 65 (1956), 25-35.
- [5] FOX, R. H. and J. MILNOR, Singularities of 2-spheres in 4-space and equivalence of knots. *Bull. Amer. Math. Soc.*, 63 (1957), 406.
- [6] HIRZEBRUCH, F., *Neue topologische Methoden in der algebraischen Geometrie*. Springer (Berlin), 1956.
- [7] —— Komplexe Mannigfaltigkeiten. *Proceedings Int. Congr. Math.*, 1958 (1960), 119-136.

- [8] MILNOR, J., Sommes de variétés différentiables et structures différentiables des sphères. *Bull. Soc. Math. France*, 87 (1959), 439-444.
- [9] —— On the cobordism ring  $\Omega^*$  and a complex analogue, I. *American Journ. of Math.*, 82 (1960), 505-521. [See also *A.M.S. Notices*, 5 (1958), 457.]
- [10] PONTRJAGIN, L. S., Characteristic cycles on differentiable manifolds. *Mat. Sbornik*, 21 (1947), 233-284.
- [11] —— Smooth manifolds and their applications in homotopy theory. *A.M.S. Translations*, 11 (1959), 1-114.
- [12] ROHLIN, V. A., A 3-dimensional manifold is the boundary of a 4-dimensional manifold. (Russian.) *Doklady Akad. Nauk S.S.R.*, 81 (1951), 355. [See also *Doklady*, v. 84, p. 221,, v. 89, p. 789 and v. 119, p. 876.]
- [13] —— and A. S. ŠVARČ, Combinatorial invariance of the Pontrjagin classes. (Russian.) *Doklady Akad. Nauk S.S.R.*, 114 (1957), 490-493.
- [14] SMALE, S., Generalized Poincaré's conjecture in dimensions greater than four. *Annals of Math.*, 74 (1961), 391-406.
- [15] STEENROD, N., *The topology of fibre bundles*. Princeton, 1951.
- [16] THOM, R., Espaces fibrés en sphères et carrés de Steenrod. *Ann. Ecole Norm. Sup.*, 69 (1952), 109-181.
- [17] —— Quelques propriétés globales des variétés différentiables. *Comment. Math. Helv.*, 28 (1954), 17-86.
- [18] —— Les classes caractéristiques de Pontrjagin des variétés triangulées. *Symposium Internacional de Topologia Algebraica, Mexico* (1958), pp. 259-272.
- [19] WALL, C. T. C., Determination of the cobordism ring. *Annals of Math.*, 72 (1960), 292-311.
- [20] WHITEHEAD, J. H. C., On  $C^1$ -complexes. *Annals of Math.*, 41 (1940), 809-824.
- [21] CONNER, P. E. and E. E. FLOYD, Differentiable periodic maps, Technical Note 13, Dept. of Math., Univ. of Virginia, 1961.
- [22] NOVIKOV, S. P., Some problems in the topology of manifolds connected with the theory of Thom spaces. *Dokl. Akad. Nauk SSSR* 132 (1960), 1031-1034 (Russian); translated as *Soviet Math. Dokl.* 1, 717-720.

Princeton University.