

# THE COHERENCE OF DIRECT IMAGES

Autor(en): **Grauert, H.**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-42345>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

**Vide-leer-empty**

# THE COHERENCE OF DIRECT IMAGES

by H. GRAUERT

## INTRODUCTION

The coherence of the direct images of coherent sheaves was treated in the paper [1]: H. Grauert: Ein Theorem der analytischen Garbentheorie und die Modulräume komplexer Strukturen (*Pub. Math. IHES* 1960, pp. 5-64, corrections 1963). This paper deals with the most general case and its technique is very difficult. The main point in the proof is the Hauptlemma on page 47. Here a proof of this Hauptlemma in the case of regular families of compact complex manifolds and locally free analytic sheaves is given. Although this special case is easier than the general, the ideas are practically the same. Therefore these lecture notes of some talks given by H. Grauert, Helsinki 1967, may lead to an understanding of the general proof. In these notes only the Hauptlemma is proved. The proof of coherence is omitted. This part is more formal and can be done like in [1] on p. 55. See [1] for applications of the theorem.

A detailed presentation of the proof in the general case is given also by Knorr [2].

## COHOMOLOGY THEORY

In this paper we use Čech cohomology. We shall briefly show how this cohomology is defined. In the following discussion  $X$  denotes a connected complex analytic manifold,  $\mathcal{O}$  is the sheaf of germs of holomorphic functions and  $S$  a sheaf of  $\mathcal{O}$ -modules. Let  $\mathfrak{U} = \{ U_i \}_{i \in J}$  be an open covering of  $X$ . We put  $U_{i_0 \dots i_l} = U_{i_0} \cap \dots \cap U_{i_l}$ . We consider cochains of order  $l$  with values in  $S$ . Let us put  $C^l(\mathfrak{U}, S) = \{ \xi \}$  where  $\xi$  denotes a full collection of crosssections  $\xi_{i_0 \dots i_l}$  over all  $U_{i_0 \dots i_l}$ . We always assume that  $\xi_{i_0 \dots i_l}$  is anticommutative in its indices. In the system  $\{ C^l(\mathfrak{U}, S) \}$  we have the

usual coboundary map  $\delta : C^l(\mathfrak{U}, S) \rightarrow C^{l+1}(\mathfrak{U}, S)$  which makes the system a complex. We put  $Z^l(\mathfrak{U}, S) = \text{Ker } \delta \subset C^l(\mathfrak{U}, S)$  and  $B^l(\mathfrak{U}, S) = \delta(C^{l-1}(\mathfrak{U}, S))$ . The  $l$ -th cohomology group  $H^l(\mathfrak{U}, S)$  with respect to the open covering  $\mathfrak{U}$  is  $Z^l(\mathfrak{U}, S)/B^l(\mathfrak{U}, S)$ . An open covering  $\mathfrak{B} = \{V_\nu\}_{\nu \in N}$  is finer than an open covering  $\mathfrak{U} = \{U_i\}_{i \in J}$  if there exists an index map  $\tau : N \rightarrow J$  such that  $V_\nu \subset U_{\tau(\nu)}$  for  $\nu \in N$ . It follows that an element of  $\Gamma(U_{\tau(\nu_0)} \dots \tau(\nu_l), S)$  can be restricted to a continuous cross-section over  $V_{\nu_0} \dots \nu_l$ . In this way we get a map  $\tau^* : C^l(\mathfrak{U}, S) \rightarrow C^l(\mathfrak{B}, S)$ . The following diagram is commutative:

$$\begin{array}{ccc} & \tau^* & \\ C^l(\mathfrak{U}, S) & \rightarrow & C^l(\mathfrak{B}, S) \\ \delta \downarrow & & \delta \downarrow \\ & \tau^* & \\ C^{l+1}(\mathfrak{U}, S) & \rightarrow & C^{l+1}(\mathfrak{B}, S) \end{array}$$

It follows that we have a map  $\tau^* : Z^l(\mathfrak{U}, S) \rightarrow Z^l(\mathfrak{B}, S)$ . Let us put  $Z^l(X, S) = \bigcup_{\mathfrak{U}} Z^l(\mathfrak{U}, S)$ , where  $\mathfrak{U}$  runs over all open coverings of  $X$ . In

$Z^l(X, S)$  we can introduce an equivalence relation  $\approx$  as follows: Let  $\xi_1 \in Z^l(\mathfrak{U}, S)$  and  $\xi_2 \in Z^l(\mathfrak{U}_1, S)$ . We put  $\xi_1 \approx \xi_2$  iff there exists  $\mathfrak{U}_2$  such that  $\mathfrak{U}_2$  is finer than  $\mathfrak{U}$  and  $\mathfrak{U}_1$  and  $\xi_1|_{\mathfrak{U}_2} - \xi_2|_{\mathfrak{U}_2} \in B^l(\mathfrak{U}_2, S)$ . Here we have put  $\xi_\nu|_{\mathfrak{U}_2} = \tau_\nu^*(\xi_\nu)$  where  $\tau_\nu^*$  comes from an index map  $\tau_\nu : \mathfrak{U}_2 \rightarrow \mathfrak{U}, \mathfrak{U}_1$ . It is easy to check that the equivalence relation defined on  $Z^l(X, S)$  is independent of the index maps. Now  $H^l(X, S)$  is the set of equivalence classes in  $Z^l(X, S)$ . Because  $C^l(\mathfrak{U}, S)$  is a module over the ring  $I(X)$  of holomorphic functions on  $X$  it follows that  $H^l(\mathfrak{U}, S)$  and  $H^l(X, S)$  are modules over  $I(X)$ . We have a natural homomorphism  $H^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$ . Let now  $X' \subset X$  be an open subset. Then  $X'$  is a complex analytic manifold. We put  $S' = S|_{X'}$  and  $\mathfrak{U}' = \mathfrak{U} \cap X' = \{U_i \cap X'\}$  and obtain an open covering of  $X'$ . The restriction of cross-sections gives a homomorphism  $\gamma : C^l(\mathfrak{U}, S) \rightarrow C^l(\mathfrak{U}', S')$  which commutes with  $\delta$  and any index map  $\tau$ . Thus we obtain restriction homomorphisms:  $H^l(\mathfrak{U}, S) \rightarrow H^l(\mathfrak{U}', S')$  and  $H^l(X, S) \rightarrow H^l(X', S')$ .

### STEIN MANIFOLDS

A complex analytic manifold  $X$  is a Stein manifold if: 1)  $X$  is holomorphically convex, i.e. if  $D = (x_\nu)_1^\infty$  is an infinite discrete set, then there exists  $f \in I(X)$  such that  $|f(D)| = \sup_\nu |f(x_\nu)|$  is infinite. 2)  $X$  can be



spread holomorphically, i.e. for any  $x \in X$  there exists  $f_1 \dots f_N \in I(X)$  such that  $x$  is an isolated common zero of  $f_1 \dots f_N$ .

Let  $X$  be a complex analytic manifold. A Stein covering  $\mathfrak{U} = \{U_i\}_{i \in J}$  of  $X$  is an open covering of  $X$  such that every  $U_i$  is Stein. We shall often use the following result:

*Leray's Theorem:* If  $\mathfrak{U}$  is a Stein covering of  $X$  then  $H^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$  is an isomorphism for every coherent analytic sheaf  $S$ .

The isomorphism between  $H^l(\mathfrak{U}, S)$  and  $H^l(X, S)$  means the following: If  $\underline{\xi} \in H^l(X, S)$  there exists  $\xi \in Z^l(\mathfrak{U}, S)$  such that  $\xi$  maps into  $\underline{\xi}$  under the natural homomorphism  $Z^l(\mathfrak{U}, S) \rightarrow H^l(X, S)$  and moreover if  $\underline{\xi} \in Z^l(\mathfrak{U}, S)$  is mapped into zero in  $H^l(X, S)$  there exist  $\eta \in C^{l-1}(\mathfrak{U}, S)$  such that  $\xi = \delta\eta$  in  $C^l(\mathfrak{U}, S)$ .

#### DIRECT IMAGES OF SHEAVES

Let  $X$  and  $Y$  be complex analytic manifolds. Let  $\psi : X \rightarrow Y$  be a holomorphic map and let  $S$  be an analytic sheaf on  $X$ . Now  $X$  is fibered by the fibers  $X(y) = \psi^{-1}(y)$  for  $y \in Y$ . Let  $U$  be an open neighborhood of a point  $y \in Y$ , then  $V = \psi^{-1}(U)$  is an open set in  $X$ . Hence  $V$  is a complex analytic manifold and the restriction of  $S$  to  $V$  gives an analytic sheaf on  $V$ . We can now define  $H^l(V, S)$ . Let us put  $H_y^l = \bigcup_U H^l(\psi^{-1}(U), S)$

where  $U$  runs over all open neighborhoods of  $y$  in  $Y$ . In  $H_y^l$  we introduce an equivalence relation as follows:  $\xi_1 \in H^l(\psi^{-1}(U_1), S)$  and  $\xi_2 \in H^l(\psi^{-1}(U_2), S)$  are equivalent iff there exists  $U = U(y)$  in  $Y$  such that  $U \subset U_1 \cap U_2$  and  $\xi_1|_{\psi^{-1}(U)} = \xi_2|_{\psi^{-1}(U)}$  in  $H^l(\psi^{-1}(U), S)$ . We let  $\psi_{(l)}(S)_{(y)}$  denote the set of equivalence classes in  $H_y^l$ . The equivalence class generated by  $\xi \in H^l(\psi^{-1}(U), S)$  is denoted by  $\xi_y$ . The set  $\psi_{(l)}(S)_{(y)}$  is called the set of germs of cohomology classes of dimension  $l$  along the fiber  $X(y)$ . Now  $\psi_{(l)}(S)_{(y)}$  is an  $\mathcal{O}_{y,Y}$ -module. For if  $g_y \in \mathcal{O}_{y,Y}$  we have a representative  $g \in I(U)$  for some open neighborhood  $U$  of  $y$ . Then  $g \circ \psi \in I(\psi^{-1}(U))$ . If  $\xi_y \in \psi_{(l)}(S)_{(y)}$  and  $U$  is sufficiently small we can find a representative  $\xi \in H^l(\psi^{-1}(U), S)$  for  $\xi_y$ . Then we put  $g_y \cdot \xi_y = ((g \circ \psi)\xi)_y$ . Now we form  $\psi_{(l)}(S) = \bigcup_{y \in Y} \psi_{(l)}(S)_{(y)}$  where we introduce a sheaf topology.

A base of the open sets are  $\{\xi_y : y \in U\}$  for  $\xi \in H^l(\psi^{-1}(U), S)$ . If  $\xi \in H^l(X, S)$  then the map  $y \rightarrow \xi_y$  is a cross-section in  $\psi_{(l)}(S)$ . We call it the direct image of  $\xi$  and denote it by  $\psi_{(l)}(\xi)$ . The sheaf  $\psi_{(l)}(S)$  is the direct

image sheaf of  $S$  of dimension  $l$ . Our main problem is to decide whether  $\psi_{(l)}(S)$  is a coherent analytic sheaf of  $\mathcal{O}_Y$ -modules if  $S$  is a coherent analytic sheaf on  $X$ .

A VERY SPECIAL CASE

We shall consider a special case where our main problem is easily solved. Let  $X_0$  be a compact analytic manifold of pure dimension  $m - n$ . We put  $E^n(\rho_0) = \{ (t_1 \dots t_n) \in \mathbf{C}^n ; |t_i| < \rho_i^0 \}$ . Here  $\rho_0 = (\rho_1^0 \dots \rho_n^0)$  is a fixed  $n$ -tuple of strictly positive numbers. Let  $X = E^n(\rho_0) \times X_0$  and  $X(\rho) = E^n(\rho) \times X_0$  for  $\rho \leq \rho_0$ . We see that  $X$  is an analytic manifold of pure dimension  $m$ . Let  $\psi : X \rightarrow E^n(\rho_0)$  be the projection map. Now  $X$  is fibered by the fibers  $\psi^{-1}(t) = X(t) = \{t\} \times X_0 \cong X_0$  for  $t \in E^n(\rho_0)$ . We take the sheaf  $S$  to be  $S = (q\mathcal{C})_X$ . With these notations we can state the following.

*Theorem:* The direct image sheaf  $\psi_{(l)}((q\mathcal{C})_X)$  is a coherent sheaf of  $\mathcal{C}_{E^n(\rho_0)}$ -modules for every  $l \geq 0$ .

*Proof.* Because  $X_0$  is a compact analytic manifold we can find a finite Stein covering  $\mathfrak{U} = \{ U_1 \dots U_{l_*} \}$  of  $X_0$ . Let us put  $\hat{U}_i = E^n(\rho_0) \times U_i$ , then we see that  $\hat{\mathfrak{U}} = \{ \hat{U}_1 \dots \hat{U}_{l_*} \}$  is a Stein covering of  $X$ . Let  $\hat{\xi} = \{ \hat{\xi}_{i_0 \dots i_l} \} \in C^l(\hat{\mathfrak{U}}, (q\mathcal{C})_X)$ . Now  $\hat{\xi}_{i_0 \dots i_l}$  is a  $q$ -tuple of holomorphic functions on  $E^n(\rho_0) \times U_{i_0 \dots i_l}$ . Hence  $\hat{\xi}_{i_0 \dots i_l}$  admits a Taylor series of the form  $\hat{\xi}_{i_0 \dots i_l} = \sum_{|v|=0}^{\infty} \xi_{i_0 \dots i_l}^{(v)} (t/\rho_0)^v$  where  $v = (v_1, \dots, v_n)$ ,  $|v| = v_1 + \dots + v_n$  and  $(t/\rho)^v = (t_1/\rho_1)^{v_1} \dots (t_n/\rho_n)^{v_n}$ . The uniqueness of a Taylor series shows that  $\{ \xi_{i_0 \dots i_l}^{(v)} \}$  is an alternating cochain over  $\mathfrak{U}$ . Putting  $\xi_{(v)} = \{ \xi_{i_0 \dots i_l}^{(v)} \} \in C^l(\mathfrak{U}, (q\mathcal{C})_X)$  we may write  $\hat{\xi} = \sum \xi_{(v)} (t/\rho)^v$ . Introducing the map  $(v) : \hat{\xi} \rightarrow \xi_{(v)}$  we get a commutative diagram of the form:

$$\begin{array}{ccc} C^l(\hat{\mathfrak{U}}, (q\mathcal{C})_X) & \xrightarrow{\delta} & C^{l+1}(\hat{\mathfrak{U}}, (q\mathcal{C})_X) \\ (v)\downarrow & & \downarrow(v) \\ C^l(\mathfrak{U}, (q\mathcal{C})_{X_0}) & \xrightarrow{\delta} & C^{l+1}(\mathfrak{U}, (q\mathcal{C})_{X_0}). \end{array}$$

We now need a *theorem of Cartan-Serre*: Let  $X_0$  be a compact analytic manifold. Then, for any coherent analytic sheaf  $S$  the set  $H^p(X_0, S)$  is a finite dimensional vector space for all  $p \geq 0$ .

Using this theorem we conclude that  $H^l(X_0, (q\mathcal{O})_{X_0})$  has a finite base  $\mathfrak{b}_1 \dots \mathfrak{b}_r$ . By Leray's theorem we also have  $H^l(\mathfrak{U}, (q\mathcal{O})_{X_0}) \cong H^l(X_0, (q\mathcal{O})_{X_0})$ . Hence we can find  $\mathfrak{b}_1 \dots \mathfrak{b}_r \in Z^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$  such that  $\mathfrak{b}_v$  maps into  $\mathfrak{b}_v$  under the natural homomorphism  $Z^l(\mathfrak{U}, (q\mathcal{O})_{X_0}) \rightarrow H^l(X_0, (q\mathcal{O})_{X_0})$ . We now introduce a pseudonorm in  $C^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$  as follows:

*Norm definition.* Let  $\eta \in C^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$ . Then we put  $\|\eta\| = \sup_{(\iota_0 \dots \iota_l)} \|\eta_{\iota_0 \dots \iota_l}\|$  and  $\|\eta_{\iota_0 \dots \iota_l}\| = \max_{1 \leq \varrho \leq l} \sup |\eta_{\varrho}(U_{\iota_0 \dots \iota_l})|$ , where,  $\eta_{\iota_0 \dots \iota_l} = (\eta_1, \dots, \eta_q)$ . Notice that it may happen that  $\|\eta\| = +\infty$ . Let  $\mathfrak{B} = \{V_1 \dots V_{l^*}\}$  be an open covering of  $X_0$ . The covering  $\mathfrak{B}$  is much finer than  $\mathfrak{U} = \{U_1 \dots U_{l^*}\}$  if  $V_\iota \subset \subset U_\iota$  holds for every  $\iota$ . We write  $\mathfrak{B} \ll \mathfrak{U}$  in that case. Let us now choose Stein coverings  $\mathfrak{B}_1$  and  $\mathfrak{B}$  such that  $\mathfrak{B}_1 \ll \mathfrak{B} \ll \mathfrak{U}$ . In  $C^l(\mathfrak{B}, (q\mathcal{O})_{X_0})$  and  $C^l(\mathfrak{B}_1, (q\mathcal{O})_{X_0})$  we introduce a pseudonorm just as in  $C^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$ . If  $\xi \in C^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$  we have defined  $\xi|_{\mathfrak{B}} \in C^l(\mathfrak{B}, (q\mathcal{O})_{X_0})$ . It follows that  $\|\xi|_{\mathfrak{B}}\| < \infty$  because  $V_{\iota_0 \dots \iota_l} \subset \subset U_{\iota_0 \dots \iota_l}$ . Let us now choose  $\xi \in Z^l(\mathfrak{B}, (q\mathcal{O})_{X_0})$ . Since  $\mathfrak{b}_1 \dots \mathfrak{b}_r \in Z^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$  constitute a base of  $H^l(X_0, (q\mathcal{O})_{X_0})$  it follows from Leray's theorem that  $\xi = \sum a_v \mathfrak{b}_v|_{\mathfrak{B}} + \delta\eta$  where  $a_v \in C^1$  and  $\eta \in C^{l-1}(\mathfrak{B}, (q\mathcal{O})_{X_0})$ . Now we need the following.

*Lemma:* There exists a constant  $K$  such that  $|a_v| \leq K \|\xi\|$  and  $\|\eta|_{\mathfrak{B}_1}\| \leq K \|\xi\|$ .

The proof follows because by the Banach theorem the map  $(a_1, \dots, a_r, \eta) \rightarrow \xi$  of the Fréchet spaces  $C^r \times C^{l-1}(\mathfrak{B}, (q\mathcal{O})_{X_0})$  onto  $Z^l(\mathfrak{B}, (q\mathcal{O})_{X_0})$  is open.

Let  $\xi \in C^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$ . We can extend each  $\xi_{\iota_0 \dots \iota_l} \in qI(U_{\iota_0 \dots \iota_l})$  constantly over  $\hat{U}_{\iota_0 \dots \iota_l} = E^n(\rho_0) \times U_{\iota_0 \dots \iota_l}$ . We get  $\hat{\xi} \in Z^l(\hat{\mathfrak{U}}, (q\mathcal{O})_X)$  obtained from  $\xi$  by a constant extension. In particular we extend  $\mathfrak{b}_1 \dots \mathfrak{b}_r$  constantly to  $\hat{\mathfrak{b}}_1 \dots \hat{\mathfrak{b}}_r \in Z^l(\hat{\mathfrak{U}}, (q\mathcal{O})_X)$ . Let  $\hat{\mathfrak{b}}_1 \dots \hat{\mathfrak{b}}_r$  be the images of  $\hat{\mathfrak{b}}_1 \dots \hat{\mathfrak{b}}_r$  in the direct image sheaf  $\psi_{(l)}((q\mathcal{O})_X)$ . Let now  $\xi_0 \in \psi_{(l)}((q\mathcal{O})_X)_{(0)}$  where 0 is the origin of  $E^n(\rho_0)$ . By definition we can find  $\xi \in H^l(X(\rho_1), q\mathcal{O})$  with  $0 < \rho_1 \leq \rho_0$  which maps into  $\xi_0$ . Now  $\hat{\mathfrak{U}}(\rho_1) = \{E^n(\rho_1) \times U_\iota\}$  is a Stein covering of  $X(\rho_1)$ . Hence Leray's theorem shows that we can find  $\hat{\xi} \in Z^l(\hat{\mathfrak{U}}(\rho_1), (q\mathcal{O})_X)$  such that  $\hat{\xi}$  maps into  $\xi_0$ . Let us write  $\hat{\xi} = \sum \xi_{(v)}(t/\rho_1)^v$  where  $\xi_{(v)} \in Z^l(\mathfrak{U}, (q\mathcal{O})_{X_0})$ . Let us also choose  $0 < \rho_2 < \rho_1$  and consider  $\hat{\xi}|_{\mathfrak{B}(\rho_2)} = \hat{\xi}_1 \in Z^l(\hat{\mathfrak{B}}(\rho_2), (q\mathcal{O})_X)$ . Let us write  $\hat{\xi}_1 = \sum \xi_{(v)}^*(t/\rho_2)^v$ . Obviously we get  $\xi_{(v)}^* = (\rho_2/\rho_1)^v \xi_{(v)}|_{\mathfrak{B}}$ . It follows easily that  $\sup_v \|\xi_{(v)}^*\| < \infty$ .

The previous lemma shows that  $\xi_{(v)}^* = \sum a_{v\lambda} b_\lambda + \delta\eta_v$  where  $\eta_v \in C^{l-1}(\mathfrak{B})$  with  $\|\eta_v|_{\mathfrak{B}_1}\| \leq K \|\xi_{(v)}^*\|$  and  $|a_{v\lambda}| \leq K \|\xi_{(v)}^*\|$ . Let us put  $a_\lambda = \sum a_{v\lambda} (t/\rho_2)^v$  and  $\hat{\eta} = \sum \eta_v (t/\rho_2)^v$ . We see that  $\hat{\eta} \in C^{l-1}(\hat{\mathfrak{B}}_1(\rho_2))$  and  $a_\lambda \in I(E^n(\rho_2))$ . An easy computation gives  $\hat{\xi}_1|_{\hat{\mathfrak{B}}_1(\rho_2)} = \sum a_\lambda \hat{b}_\lambda|_{\hat{\mathfrak{B}}_1(\rho_2)} + \delta\hat{\eta}$ . It follows by definition that  $\xi_0 = \sum a_\lambda \hat{b}_\lambda$ . We have now proved that  $\hat{b}_1 \dots \hat{b}_r$  generate  $\psi_{(l)}((q\mathcal{O})_X)$  at the origin. It follows in the same way that  $\hat{b}_1 \dots \hat{b}_r$  generate  $\psi_{(l)}((q\mathcal{O})_X)$  for every  $t \in E^n(\rho_0)$  because it is enough to do everything in a polydisc around  $t$ . Now we also prove that the sheaf  $\psi_{(l)}((q\mathcal{O})_X)$  is free, i.e. there are no relations between  $\hat{b}_1 \dots \hat{b}_r$  at any point. Say for example that  $a_1 \hat{b}_1 + \dots + a_r \hat{b}_r = 0$  at  $\psi_{(l)}((q\mathcal{O})_X)_{(0)}$  where  $a_i$  are germs of analytic functions at the origin in  $E^n(\rho_0)$ . Hence  $\tilde{a}_1 \hat{b}_1 + \dots + \tilde{a}_r \hat{b}_r = 0$  in  $H^l(X(\rho), (q\mathcal{O})_X)$  for some  $\rho > 0$  with  $\tilde{a}_i \in I(E^n(\rho))$ . It follows that  $\sum \tilde{a}_v \hat{b}_v = \delta \hat{\xi}$  in  $X(\rho)$  for some  $\hat{\xi} \in C^{l-1}(\hat{\mathcal{U}}(\rho), (q\mathcal{O})_X)$ . Take a point  $t \in E^n(\rho)$  where some  $\tilde{a}_v \neq 0$ . Now we see that on  $\{t\} \times X_0$  we have  $\tilde{a}_1(t) b_1 + \dots + \tilde{a}_r(t) b_r = \delta \hat{\xi}|_{\{t\} \times X_0} \in C^{l-1}(\mathcal{U}, (q\mathcal{O})_{X_0})$ . This gives a contradiction to the fact that  $b_1 \dots b_r$  are a base of  $H^l(X_0, (q\mathcal{O})_{X_0})$ .

### MEASURE CHARTS

Let  $X$  be a connected complex analytic manifold of dimension  $m$ . Let  $F$  be a holomorphic vector bundle of rank  $q$  on  $X$  and  $\mathbf{F}$  the sheaf of holomorphic crosssections in  $F$ . This sheaf is locally free. A regular proper holomorphic map  $\psi: X \rightarrow E^n$  is given. Let us put  $X_0 = \psi^{-1}(0)$ . Now  $X_0$  is a compact analytic manifold of dimension  $m - n$ . We now introduce special open coverings around  $X_0$  in  $X$ .

*Definition.* A measure chart  $\mathcal{W} = (\hat{W}, \Phi, \Theta, \rho)$  is a quadruple satisfying the conditions:

- 1)  $\hat{W} \subset X$  is open and  $W = \hat{W} \cap X_0$  is Stein.
- 2)  $\Phi: \hat{W} \rightarrow E^n(\rho) \times W$  is a biholomorphic map such that the following diagram is commutative:

$$\begin{array}{ccc} \hat{W} & \xrightarrow{\Phi} & E^n(\rho) \times W \\ \psi \searrow & & \swarrow \pi \\ & & E^n(\rho). \end{array}$$

Here  $\pi$  is the projection map.

3)  $\Theta: F|_{\hat{W}} \rightarrow \hat{W} \times \mathbf{C}^q$  is a trivialization of  $F$  on  $\hat{W}$ .

If  $\mathcal{W}$  is a given measure chart on  $X$  we can identify the sheaf  $(\hat{W}, F|_{\hat{W}})$  of  $\mathcal{C}_X$ -modules with the sheaf  $(W \times E^n(\rho), q\mathcal{O})$  using  $\Phi$  and  $\Theta$ . If  $U \subset W$  is open and  $\rho' \leq \rho$  we put  $\hat{U}(\rho') = \Phi^{-1}(U \times E^n(\rho'))$ . Hence if  $\hat{s} \in \Gamma(\hat{U}(\rho'), F)$  we can identify  $\hat{s}$  with an element of  $\Gamma(U \times E^n(\rho'), q\mathcal{O})$ . We shall simply denote this element of  $\Gamma(U \times E^n(\rho'), q\mathcal{O})$  by the same letter  $\hat{s}$ . Now we can expand  $\hat{s}$  in a Taylor series:  $\hat{s} = \sum_{|v|=0}^{\infty} s_v (t/\rho')^v$  where  $s_v \in qI(U)$ .

*Definition of a norm.* When  $\hat{s} \in \Gamma(\hat{U}(\rho'), F)$  we put  $\|\hat{s}\| = \sup_v |s_v(U)|$ .

Strictly speaking the norm  $\|\hat{s}\|$  is taken with respect to the measure chart  $\mathcal{W}$ .

It is not hard to see that for every point  $x \in X_0$  there exists a measure chart  $\mathcal{W}$  such that  $x \in \hat{W}$ . In particular we can cover  $X_0$  by finitely many measure charts  $\mathcal{W}_\iota = (\hat{W}_\iota, \Phi_\iota, \Theta_\iota, \rho_\iota)$ , i.e.  $X_0 \subset \subset \bigcup_1^{\iota^*} \hat{W}_\iota$ . We remark that it

follows that  $X(\rho) = \psi^{-1}(E^n(\rho)) \subset \subset \bigcup_1^{\iota^*} \hat{W}_\iota$  for some  $\rho > 0$  with  $\rho \leq \rho_\iota$  because  $\psi$  is a proper map. The collection  $\mathcal{W} = \{\mathcal{W}_\iota\}_1^{\iota^*}$  is called an atlas around  $X_0$ . From now on  $\mathcal{W}$  is a fixed atlas.

*Measure coverings.* We shall define measure coverings with respect to the given atlas  $\mathcal{W}$  above. If  $U \subset W_\iota$  is open we put  $(U)_\iota(\rho) = \Phi_\iota^{-1}(U \times E^n(\rho))$  when  $\rho \leq \rho_\iota$ . We see that  $(U)_\iota(\rho) \subset \hat{W}_\iota$  and  $(U)_\iota(\rho)$  is Stein if  $U$  is Stein. Let  $\mathfrak{U} = \{U_\iota\}_1^{\iota^*}$  be a Stein covering of  $X_0$  with  $U_\iota \subset \subset W_\iota$  for each  $\iota$ . Let  $\rho > 0$  with  $\rho < \min_\iota \rho_\iota$ . We put  $\hat{U}_\iota(\rho) = (U_\iota)_\iota(\rho)$ . We see that  $\hat{U}_\iota(\rho) \subset \subset \hat{W}_\iota$  and  $\hat{U}_\iota(\rho)$  are Stein. It is now required that  $\hat{\mathfrak{U}}(\rho) =$

$= \{ \hat{U}_i(\rho) \}_1^{i^*}$  is a Stein covering of  $X(\rho)$ . We say then that  $\hat{\mathfrak{U}}(\rho)$  is a measure covering of  $X(\rho)$ .

*Admissible refinements of measure coverings.* Let  $\hat{\mathfrak{U}}(\rho)$  and  $\hat{\mathfrak{U}}^*(\rho)$  be two measure coverings of  $X(\rho)$ . We say that  $\hat{\mathfrak{U}}^*(\rho)$  is an admissible refinement of  $\hat{\mathfrak{U}}(\rho)$  if the following conditions hold:

- 1)  $U_i^* \subset \subset U_i$  for each  $i$ .
- 2) If  $U_{i_0 \dots i_\lambda}^* = U_{i_0}^* \cap \dots \cap U_{i_\lambda}^*$  we put  $(U_{i_0 \dots i_\lambda}^*)_v = \Phi_v^{-1}(U_{i_0 \dots i_\lambda}^* \times E^n(\rho))$  for each  $v \in \{i_0 \dots i_\lambda\}$ . It is now required that  $(U_{i_0 \dots i_\lambda}^*)_v \subset (U_{i_0 \dots i_\lambda}^*)_\mu$  for all  $v, \mu \in \{i_0 \dots i_\lambda\}$ .
- 3)  $\hat{U}_{i_0 \dots i_\lambda}^* = \hat{U}_{i_0}^* \cap \dots \cap \hat{U}_{i_\lambda}^* \subset (U_{i_0 \dots i_\lambda}^*)_\mu$  for each  $\mu \in \{i_0 \dots i_\lambda\}$ .

#### EXISTENCE OF ADMISSIBLE REFINEMENTS OF MEASURE COVERINGS

*Existence Theorem.* For every fixed integer  $s$  we can find, for some  $\rho > 0$ , a sequence  $\mathfrak{U}_s \ll \mathfrak{U}_{s-1} \ll \dots \ll \mathfrak{U}_1 \ll \mathfrak{U}_0$  of finer measure coverings of  $X(\rho)$  each of which is an admissible refinement of the following.

*Proof.* We first construct a measure covering of  $X(\rho)$  for some  $\rho < \min \rho_i$ . Let  $\mathfrak{U}_0 = \{ \mathfrak{U}_i \}_1^{i^*}$  be a Stein covering of  $X_0$  such that  $U_i \subset \subset W_i$  for  $i \in \{1, \dots, i^*\}$ . Choose a fixed  $\rho_0 < \min \rho_i$ . Now the open sets  $\Phi_i^{-1}(U_i \times E^n(\rho_0))$  cover  $X_0$  and hence they also cover  $X(\rho)$  for some sufficiently small  $\rho$ . Hence  $\mathfrak{U}_0$  defines a measure covering of  $X(\rho)$ . It is also clear that  $\mathfrak{U}_0$  defines a measure covering of  $X(\rho')$  for each  $\rho' \leq \rho$ . Let us now construct  $\mathfrak{U}_1$ . We let  $\mathfrak{U}^* = \{ U_i^* \}_1^{i^*}$  be a Stein covering such that  $U_i^* \subset \subset U_i$  always holds. Now we can find  $\rho_1 \leq \rho$  such that  $\{ \hat{U}_i^*(\rho_1) = \Phi_i^{-1}(U_i^* \times E^n(\rho_1)) \}_1^{i^*}$  cover  $X(\rho_1)$ . Hence  $\hat{\mathfrak{U}}^*(\rho_1)$  and  $\hat{\mathfrak{U}}(\rho_1)$  are measure coverings of  $X(\rho_1)$ . But we do not yet know if  $\hat{\mathfrak{U}}^*(\rho_1) \ll \hat{\mathfrak{U}}(\rho_1)$ . We claim that if  $\rho_2 \leq \rho_1$  is sufficiently small then  $\hat{\mathfrak{U}}^*(\rho_2) \ll \hat{\mathfrak{U}}(\rho_2)$ . For suppose this is false. Say that 2) fails for  $\hat{\mathfrak{U}}^*(\rho_2)$  and  $\hat{\mathfrak{U}}(\rho_2)$  when  $0 < \rho_2 \leq \rho_1$ . Hence  $\Phi_v^{-1}(U_{i_0 \dots i_\lambda}^* \times E^n(\rho_2)) - \Phi_\mu^{-1}(U_{i_0 \dots i_\lambda} \times E^n(\rho_2))$  are non empty for suitable indices while  $\rho_2 \rightarrow 0$ . Choose a point  $x_t$  from each of these sets. Because  $x_t \in X(\rho_1)$  which is relatively compact we may assume that  $x_t \rightarrow x_0$ . Obviously we get  $x_0 \in \overline{U_{i_0 \dots i_\lambda}^*} - U_{i_0 \dots i_\lambda}$ , a contradic-

tion because  $\overline{U_{\iota_0 \dots \iota_\lambda}^*} \subset \overline{U_{\iota_0}^*} \cap \dots \cap \overline{U_{\iota_\lambda}^*} \subset U_{\iota_0 \dots \iota_\lambda}$ . In the same way we can prove that condition 3) is satisfied if  $\rho_2$  is sufficiently small and the theorem is clear.

### GENERAL THEORY

Let  $G$  be an analytic manifold. We put  $\hat{G} = G \times E^n(\rho_1)$  where  $\rho_1$  is an  $n$ -tuple of positive numbers. Let  $\pi: \hat{G} \rightarrow E^n(\rho_1)$  and  $\mathfrak{P}: \hat{G} \rightarrow G$  be the projection maps.  $\hat{G}^* \subset \hat{G}$  denotes an open subset and  $G^* = \hat{G}^* \cap G \times \{0\}$ .

The set  $G^*$  can be identified with an open subset of  $G$ . We denote by  $\alpha: G^* \times E^n(\rho_1) \rightarrow \hat{G}^*$  a biholomorphic fiber preserving map, i.e.  $\pi \circ \alpha = \pi^*$  where  $\pi^*: G^* \times E^n(\rho_1) \rightarrow E^n(\rho_1)$  is the natural projection. Let  $\rho \leq \rho_2 = \gamma \rho_1 < \rho_1$  where  $0 < \gamma < 1$  is a fixed number. We put  $\hat{G}(\rho) = G \times E^n(\rho)$ . If  $f$  is a holomorphic function on  $\hat{G}(\rho)$  we write  $f = \sum a_\nu (t/\rho)^\nu$  with  $a_\nu \in I(G)$ . We define the norm  $\|f\|_\rho$  of  $f$  by  $\|f\|_\rho = \sup_\nu \{ \sup |a_\nu(G)| \}$ .

If  $f \in I(\hat{G}(\rho))$  we see that  $f \circ \alpha$  is a well defined function on  $G^* \times E^n(\rho)$  because  $\alpha$  is fiber preserving. We define  $\|f \circ \alpha\|_\rho$  using  $G^*$  instead of  $G$  as above. We have the proposition:

*Proposition 1.* There exists a constant  $K$  such that  $\|f \circ \alpha\|_\rho \leq K \|f\|_\rho$  where  $K = K(\rho_2)$  is independent of  $\rho \leq \rho_2$ .

*Proof.* We write  $f = \sum_{|\nu|=0}^{\infty} a_\nu (t/\rho)^\nu$  with  $a_\nu \in I(G)$ . Now we get  $f \circ \alpha = \sum (a_\nu \circ \mathfrak{P} \circ \alpha) (t/\rho)^\nu$  because  $\alpha$  is fiber preserving. Since  $\mathfrak{P}(\hat{G}^*) \subset G$  we get  $|a_\nu \circ \mathfrak{P}(\hat{G}^*)| \leq |a_\nu(G)| \leq \|f\|_\rho$ . Now  $a_\nu \circ \mathfrak{P} \circ \alpha$  admits a Taylor series:  $a_\nu \circ \mathfrak{P} \circ \alpha = \sum C_{\nu\lambda} (t/\rho)^\lambda$  with  $C_{\nu\lambda} \in I(G^*)$ . Since  $|\sum C_{\nu\lambda} (t/\rho)^\lambda| \leq \|f\|_\rho$  in  $G^* \times E^n(\rho_1)$  and  $\rho \leq \rho_2 = \gamma \rho_1$  Cauchy's inequalities give us  $|C_{\nu\lambda}(G^*)| \leq \|f\|_\rho \gamma^{|\lambda|}$ . Let us put  $b_\mu = \sum_{\nu+\lambda=\mu} C_{\nu\lambda}$ . We get  $|b_\mu(G^*)| \leq \|f\|_\rho \sum \gamma^{|\lambda|} = \|f\|_\rho (1-\gamma)^{-n} = K \|f\|_\rho$ . Now we can write  $f \circ \alpha = \sum_\nu a_\nu \circ \mathfrak{P} \circ \alpha (t/\rho)^\nu = \sum_{\lambda,\nu} C_{\nu\lambda} (t/\rho)^\lambda (t/\rho)^\nu = \sum_\mu b_\mu (t/\rho)^\mu$ . By definition we have  $\|f \circ \alpha\|_\rho = \sup_\mu |b_\mu(G^*)| \leq K \|f\|_\rho$ .

Let us now consider  $\mathbf{h} = (h_{\nu\mu})$  which is a  $q \times q$  matrix with  $h_{\nu\mu} \in I(\hat{G})$ . The  $h_{\nu\mu}$  are also assumed to be bounded on  $\hat{G}$ .



*Proposition 2.* Let  $\mathbf{f} = (f_1 \dots f_q) \in qI(\hat{G}(\rho))$ . Then  $\|\mathbf{h}(\mathbf{f})\|_\rho \leq K \|\mathbf{f}\|_\rho$ . As before  $\rho \leq \rho_2 = \gamma\rho_1 < \rho_1$  and  $K$  only depends on  $\rho_2$ .

*Proof.* We have  $\mathbf{h}(\mathbf{f}) = (g_1 \dots g_q)$  with  $g_\nu = \sum_\mu h_{\nu\mu} f_\mu$ . Let us write  $h_{\nu\mu} = \sum_\lambda a_{\nu\mu\lambda} (t/\rho)^\lambda$ . By assumption  $|h_{\nu\mu}(\hat{G})| \leq M$  for some constant  $M$  and hence we have, by Cauchy's inequalities,  $|a_{\nu\mu\lambda}(G)| \leq M\gamma^{|\lambda|}$ . Let us also write  $f_\mu = \sum_\lambda b_{\mu\lambda} (t/\rho)^\lambda$ . By definition  $\sup_{\mu,\lambda} |b_{\mu\lambda}(G)| = \|\mathbf{f}\|_\rho$ . Now we get  $g_\nu = \sum_\mu \sum_{\lambda_1, \lambda_2} a_{\nu\mu\lambda_1} b_{\mu\lambda_2} (t/\rho)^{\lambda_1 + \lambda_2} = \sum_{\nu,\lambda} C_{\nu\lambda} (t/\rho)^\lambda$  where  $C_{\nu\lambda} = \sum_\mu \sum_{\lambda_1 + \lambda_2 = \lambda} a_{\nu\mu\lambda_1} b_{\mu\lambda_2}$ . We get easily  $|C_{\nu\lambda}(G)| \leq qM \|\mathbf{f}\|_\rho (1-\gamma)^{-n} = K \|\mathbf{f}\|_\rho$ . Hence  $\|\mathbf{h}(\mathbf{f})\|_\rho = \sup_\nu \|g_\nu\|_\rho = \sup_{\nu,\lambda} |C_{\nu\lambda}(G)| \leq K \|\mathbf{f}\|_\rho$ .

We shall now apply these two propositions to our situation. Let  $G^* \subset G \subset W_{\iota_0 \dots \iota_\lambda} \subset X_0$ . Here  $G^*$  and  $G$  are open sets and  $W_{\iota_0 \dots \iota_\lambda}$  comes from the measure atlas  $\mathcal{W}$ . As before  $\rho \leq \rho_2 < \rho_* = \min \rho_{\iota_i}$ . We are given  $\iota$  and  $\iota'$  from  $\{\iota_0, \dots, \iota_\lambda\}$  and the following inclusions are assumed:  $(G^*)_{\iota'}(\rho_1) \subset (G)_\iota(\rho_1)$ ,  $(G^*)_{\iota'}(\rho_1) \subset \subset \hat{W}_{\iota'}$ ,  $(G)_\iota(\rho_1) \subset \subset \hat{W}_\iota$ .

The following theorem is very important.

*Theorem I.* Let  $S \in \Gamma((G)_\iota(\rho), \mathbf{F})$ . Then  $\|S|(G^*)_{\iota'}(\rho)\|_{\iota'} \leq K \|S\|_\iota$ .  $K$  depends only on  $\rho_2$ .

*Proof.* We have the following diagram:

$$\begin{array}{ccc} & \Phi_\iota & \\ & (G)_\iota(\rho_1) \rightarrow G \times E^n(\rho_1) & \\ \text{injection} \uparrow & & \uparrow \alpha \\ & \Phi_{\iota'} & \\ & (G^*)_{\iota'}(\rho_1) \rightarrow G^* \times E^n(\rho_1) & \end{array}$$

$\alpha$  being a fiber preserving holomorphic map. We identify  $S|(G^*)_{\iota'}(\rho)$  with an element of  $qI(G^* \times E^n(\rho))$  using the trivialization of  $F$  in the chart  $\mathcal{W}_{\iota'}$ . Call this element  $S^*$ . Also  $S$  itself is considered as an element of  $qI(G \times E^n(\rho))$  using the trivialization in the chart  $\mathcal{W}_\iota$ . Now we have  $S^* = \mathbf{h}(S \circ \alpha)$  where  $\mathbf{h}$  is a  $q \times q$  matrix. The elements of  $\mathbf{h}$  are holomorphic functions defined on  $\Phi_{\iota'}(\hat{W}_{\iota'}) \supset \supset G^* \times E^n(\rho_1)$ . Hence the elements of  $\mathbf{h}$  are bounded on  $G^* \times E^n(\rho_1)$ . It is now obvious how we can use 1) and 2) to finish the proof.

We shall need one more general result. Let  $G$  be an analytic manifold.  $G$  is assumed to be Stein and  $R^* = \{U_1, \dots, U_{\iota^*}\}$  a Stein covering of  $G$ .



The set  $G^* \subset G$  is open and  $R^{**} = \{V_1, \dots, V_{l^*}\}$  an open covering of  $G^*$  such that  $V_l \subset \subset U_l$  for  $l \in \{1, \dots, l^*\}$ . We have:

*Cartan's Theorem.* There exists a constant  $K$  such that if  $\xi \in Z^l(R^*, q\mathcal{O})$  then  $\xi|_{R^{**}} = \delta\eta$  where  $\eta \in C^{l-1}(R^{**}, q\mathcal{O})$  and  $\|\eta\| \leq K\|\xi\|$  for  $l \geq 1$ .

This is a simple consequence of Theorem B and Banach's open mapping theorem.

Now we apply Cartan's theorem. We keep the notations as above. Let  $\hat{G} = G \times E^n(\rho)$  and put  $\hat{R}^* = \{U_l \times E^n(\rho)\}$ . Now  $\hat{R}^*$  is a Stein covering of  $\hat{G}$ . Let  $\hat{G}^* = G^* \times E^n(\rho)$  and  $\hat{R}^{**} = \{V_l \times E^n(\rho)\}$ . Let  $\hat{\xi} \in Z^l(\hat{R}^*, q\mathcal{O})$  and write  $\hat{\xi} = \sum \xi_{(v)}(t/\rho)^v$  with  $\xi_{(v)} \in Z^l(R^*, q\mathcal{O})$ . We assume  $\|\hat{\xi}\|_\rho = \sup_v \|\xi_{(v)}\| < \infty$ . Now Cartan's theorem gives  $\xi_{(v)}|_{R^{**}} = \delta\eta_v$  with  $\eta_v \in C^{l-1}(R^{**}, q\mathcal{O})$  and  $\|\eta_v\| \leq K\|\xi_{(v)}\| < \infty$ . It follows that  $\hat{\eta} = \sum \eta_v(t/\rho)^v$  is well defined in  $C^{l-1}(\hat{R}^{**}, q\mathcal{O})$  and by definition we have  $\|\hat{\eta}\|_\rho \leq K\|\hat{\xi}\|_\rho$ .

### SMOOTHING

We are given a sequence of admissible refinements of measure coverings in  $X(\rho_1)$ . Here  $\rho_1 < \rho_0 = \min \rho_l$  as usual. Let  $l$  be a fixed integer  $\geq 1$ . We are given  $\mathfrak{B}^* \ll \mathfrak{B}' = \mathfrak{B}_{3l} \ll \mathfrak{B}_{3l-1} \ll \dots \ll \mathfrak{B}_1 \ll \mathfrak{B}_0 \ll \mathfrak{B} \ll \mathfrak{U}^* \ll \mathfrak{U} = \mathfrak{U}_{3l} \ll \dots \ll \mathfrak{U}_0 \ll \mathfrak{U}'$ . Here it is also required that  $(\mathfrak{B}_{v+1}, \mathfrak{U}_{v+1}) \ll (\mathfrak{B}_v, \mathfrak{U}_v)$ ;  $(\mathfrak{B}^*, \mathfrak{U}^*) \ll (\mathfrak{B}', \mathfrak{U})$  and  $(\mathfrak{B}_0, \mathfrak{U}_0) \ll (\mathfrak{B}, \mathfrak{U}')$ . These extra conditions mean: 1)  $\hat{U}_{i_0 \dots i_\kappa}^{(v+1)} \cap \hat{V}_{l_0 \dots l_l}^{(v+1)} \subset (U_{i_0 \dots i_\kappa}^{(v)} \cap V_{l_0 \dots l_l}^{(v)})_i$  for each  $i \in \{i_0, \dots, i_\kappa\}$  and 2)  $(U_{i_0 \dots i_\kappa}^{(v+1)} \cap V_{l_0 \dots l_l}^{(v+1)})_j \subset (U_{i_0 \dots i_\kappa}^{(v)} \cap V_{l_0 \dots l_l}^{(v)})_i$  for all  $i, j \in \{i_0, \dots, i_\kappa, l_0, \dots, l_l\}$ .

Recall that all operations are done with respect to  $\rho_1$ . Let us put  $\hat{R}_{i_0 \dots i_\kappa l_0 \dots l_l}^{(v)} = \hat{U}_{i_0 \dots i_\kappa}^{(v)} \cap \hat{V}_{l_0 \dots l_l}^{(v)}$ . We consider elements  $\xi_{i_0 \dots i_\kappa l_0 \dots l_l} \in \hat{\Gamma}(\hat{R}_{i_0 \dots i_\kappa l_0 \dots l_l}^{(v)}, \mathbb{F})$ .

Now we take a full collection  $\hat{\xi} = \{\hat{\xi}_{i_0 \dots i_\kappa l_0 \dots l_l}\}$  of such elements which is anticommutative in  $\{i_0, \dots, i_\kappa\}$  and  $\{l_0, \dots, l_l\}$ . In this way we get a double complex  $C_v^{k, \kappa}$ . Here  $\delta : C_v^{k, \kappa} \rightarrow C_v^{k+1, \kappa}$  and  $\partial : C_v^{k, \kappa} \rightarrow C_v^{k, \kappa+1}$  are the usual coboundary operators.

NORM IN  $C_v^{k, \kappa}$ : Let  $\hat{\xi} \in C_v^{k, \kappa}$ ; we put

$\|\hat{\xi}\|_\rho = \max_{i, (i_0, \dots, i_k, \iota_0, \dots, \iota_k)} \{ \|\hat{\xi}_{i_0 \dots i_k \iota_0 \dots \iota_k} \mid (R_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v+1)})_i(\rho) \|\}_i$  with  $i \in \{i_0, \dots, i_k\}$ . Here  $\rho \geq \rho_1$  and  $R_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v+1)} = U_{i_0 \dots i_k}^{(v+1)} \cap V_{\iota_0 \dots \iota_k}^{(v+1)}$  and  $\|\cdot\|_i$  is taken with respect to the chart  $\mathscr{W}_i$  as usual.

**SMOOTHING LEMMA:** Let  $\kappa > 0$ . There exists a constant  $K$  such that: If  $\hat{\xi} \in C_v^{k, \kappa}$  with  $\partial \hat{\xi} = 0$  and  $\|\hat{\xi}\|_\rho < \infty$  then we can find  $\hat{\eta} \in C_{v+3}^{k, \kappa-1}$  such that  $\hat{\xi} \mid C_{v+3}^{k, \kappa} = \partial \hat{\eta}$  and  $\|\hat{\eta}\|_\rho \leq K \|\hat{\xi}\|_\rho$ . Here  $\rho \leq \rho_2 = \gamma \rho_1$  with  $0 < \gamma < 1$  and  $K$  depends only on  $\rho_2$ .

*Proof.* Let us fix  $i_0, \dots, i_k$  in the following discussion. Let  $G = U_{i_0 \dots i_k}^{(v+1)}$  and put  $\hat{G} = (G)_i(\rho_1)$  for some  $i \in \{i_0, \dots, i_k\}$  which is also fixed now. Now  $G$  is Stein in  $X_0$  and  $\hat{G}$  is Stein in  $X$ . We put  $R^* = G \cap \mathfrak{B}_{v+1}$  which is a Stein covering of  $G$ . Also  $\hat{R}^* = \{(G \cap V_{\iota}^{(v+1)})_i(\rho_1)\}_{\iota=1, \dots, \iota^*}$  is a Stein covering of  $\hat{G}$ . Let  $\hat{\xi} = \{\hat{\xi}_{i_0 \dots i_k \iota_0 \dots \iota_k}\}$ . Now we look at the elements of  $\{\hat{\xi}_{i_0 \dots i_k \iota_0 \dots \iota_k}\} = \hat{\xi}_{i_0 \dots i_k} \in Z^\kappa(\hat{R}^*, \mathbf{F})$ . Here  $i_0, \dots, i_k$  is fixed as above. We get a cocycle because we have assumed that  $\partial \hat{\xi} = 0$ . More precisely we have considered the restriction of  $\hat{\xi}_{i_0 \dots i_k \iota_0 \dots \iota_k}$  to  $\hat{R}^*$ . We must verify that this restriction is possible.

*Verification:* By definition of  $Z^\kappa(\hat{R}^*, \mathbf{F})$  we have to look at sets of the following type: (these are the sets where the cross-sections are defined)  $(G \cap V_{\iota_0}^{(v+1)})_i \cap \dots \cap (G \cap V_{\iota_k}^{(v+1)})_i = (G \cap V_{\iota_0 \dots \iota_k}^{(v+1)})_i = (R_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v+1)})_i$ . Now by 2) we have  $(R_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v+1)})_i \subset \bigcap_j (R_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v)})_j \subset (U_{i_0}^{(v)})_{i_0} \cap \dots \cap (V_{\iota_k}^{(v)})_{\iota_k} = \hat{R}_{i_0 \dots i_k \iota_0 \dots \iota_k}^{(v)}$ . Q.E.D.

Now we put  $G^* = U_{i_0 \dots i_k}^{(v+2)} \subset \subset G$ . We let  $\hat{R}^{**} = \{(G^* \cap V_{\iota}^{(v+2)})_i\}_{\iota=1, \dots, \iota^*}$ . The system  $\hat{R}^{**}$  is a Stein covering of  $(G^*)_i$ . We are in a good position now. For we are given  $\hat{\xi}_{i_0 \dots i_k} \in Z^\kappa(\hat{R}^*, \mathbf{F})$ . Here  $\hat{R}^*$  is a Stein covering of  $\hat{G}$  and  $\hat{G}$  is a Stein manifold. We are working in the chart  $\mathscr{W}_i$  where the usual identifications are used. Hence we arrive at the following situation:  $G$  is a Stein manifold with a Stein covering  $R^* = \mathfrak{B}_{v+1} \cap G$ . Also  $G^* \subset \subset G$  and  $R^{**} = \mathfrak{B}_{v+2} \cap G^*$  is a Stein covering of  $G^*$  such that  $R^{**} \subset \subset R^*$ . The cocycle  $\hat{\xi}_{i_0 \dots i_k}$  is now considered as an element of  $Z^\kappa(\hat{R}^*, q\mathcal{O})$  which

we simply call  $\hat{\xi}_{i_0 \dots i_k}$  again. Now we apply the result after Cartan's theorem. Hence we can find a constant  $K$  such that for every  $\rho \leq \rho_2$  we get  $\eta \in C^{\kappa-1}(\hat{R}^{**}, q\mathcal{O})$  and  $\|\eta\|_\rho \leq K \|\hat{\xi}_{i_0 \dots i_k}\|_\rho$  with  $\partial\eta = \hat{\xi}_{i_0 \dots i_k}$ . But this means precisely that we can find  $\hat{\eta}_{i_0 \dots i_k} \in C^{\kappa-1}(\hat{R}^{**}(\rho), \mathbb{F})$  such that  $\|\hat{\eta}_{i_0 \dots i_k}\|_{i, \rho} \leq K \|\hat{\xi}_{i_0 \dots i_k}\|_{i, \rho}$  with  $\hat{\xi}_{i_0 \dots i_k} = \partial\hat{\eta}_{i_0 \dots i_k}$ . We have only constructed  $\hat{\eta}_{i_0 \dots i_k}$  using a fixed  $i \in \{i_0, \dots, i_k\}$ . Now we must let  $(i_0, \dots, i_k)$  vary. For each  $(i_0, \dots, i_k)$  we choose some  $i$  which only depends on the unordered  $(k+1)$ -tupel  $(i_0, \dots, i_k)$  and construct an element  $\hat{\eta}_{i_0 \dots i_k}$  as above. Now we can restrict everything to  $C_{v+3}^{k, \kappa-1}$ .

*Verification:* Consider a set where cross-sections over  $C_{v+3}^{k, \kappa-1}$  have to be defined, i.e. a set  $\hat{U}_{i_0 \dots i_k}^{(v+3)} \cap \hat{V}_{i_0 \dots i_k}^{(v+3)}$ . But by 1) follows  $\hat{U}_{i_0 \dots i_k}^{(v+3)} \cap \hat{V}_{i_0 \dots i_k}^{(v+3)} \subset (R_{i_0 \dots i_k, i_0 \dots i_k}^{(v+2)})_i$  for each  $i \in \{i_0, \dots, i_k\}$ . This inclusion shows that we get a well defined element  $\hat{\eta} \in C_{v+3}^{k, \kappa-1}$  by restricting the elements  $\hat{\eta}_{i_0 \dots i_k}$  to  $C_{v+3}^{k, \kappa-1}$ . We find that  $\hat{\xi} \mid C_{v+3}^{k, \kappa} = \partial\hat{\eta}$  now. The norm inequalities are not obvious, but recalling how  $\hat{\eta}$  is constructed here it is seen that we can apply Theorem I to obtain the required estimate.

**SMOOTHING THEOREM.** There exists a constant  $K$  such that: If  $\hat{\xi} \in Z^l(\hat{\mathfrak{B}}(\rho), \mathbb{F})$  with  $\|\hat{\xi}\|_\rho < \infty$  then we can find  $\hat{\xi}^* \in Z^l(\hat{\mathfrak{U}}(\rho), \mathbb{F})$  and  $\hat{\eta} \in C^{l-1}(\hat{\mathfrak{B}}'(\rho), \mathbb{F})$  for which  $\hat{\xi}^* \mid \hat{\mathfrak{B}}'(\rho) = \hat{\xi} \mid \hat{\mathfrak{B}}'(\rho) + \delta\hat{\eta}$  and  $\|\hat{\xi}^*\|_\rho$  and  $\|\hat{\eta}\|_\rho \leq K \|\hat{\xi}\|_\rho$ . Here  $\rho \leq \rho_2 < \rho_1$  and  $K$  only depends on  $\rho_2$ .

*Proof.* Before we can use the double complex  $\{C_v^{k, \kappa}\}$  we must introduce two "ε-maps". To define the  $\varepsilon_1$ -map, let  $Z_v^{k, \kappa} \subset C_v^{k, \kappa}$  consist of all  $\hat{\xi} \in C_v^{k, \kappa}$  such that  $\delta\hat{\xi} = \partial\hat{\xi} = 0$ . Now we shall define the  $\varepsilon_1$ -map:  $\varepsilon_1: Z^l(\hat{\mathfrak{B}}, \mathbb{F}) \rightarrow Z_0^{0, l}$ . A section belonging to an element of  $C_0^{0, l}$  is defined on some set  $\hat{U}_{i_0}^{(0)} \cap \hat{V}_{i_0, \dots, i_l}^{(0)} \subset \hat{V}_{i_0 \dots i_l}$  where sections of elements of  $Z^l(\hat{\mathfrak{B}}, \mathbb{F})$  are defined. Hence we get a natural restriction map  $\varepsilon_1$  which also maps cocycles into cocycles. It is easy to verify that  $\|\varepsilon_1(\hat{\xi})\|_\rho \leq K \|\hat{\xi}\|_\rho$ . Theorem I can be used because  $(U_i^{(1)} \cap V_{i_0 \dots i_l}^{(1)})_i \subset (V_{i_0 \dots i_l}^{(0)})_i$  for every  $i$  and every  $i \in \{i_0, \dots, i_l\}$ . Recall that the norm in  $Z^l(\hat{\mathfrak{B}}, \mathbb{F})$  is defined with respect to

$\hat{\mathfrak{B}}_0$  here. The “ $\varepsilon_2$ -map” : we shall construct a map  $\varepsilon_2: Z_{3l}^{l,0} \rightarrow Z^l(\hat{\mathfrak{U}}, \mathbf{F})$ . Let  $\hat{\xi} = \{ \hat{\xi}_{i_0, \dots, i_l, \iota_0} \} \in Z_{3l}^{l,0}$ . Here  $\hat{\xi}_{i_0, \dots, i_l, \iota_0}$  is defined on  $\hat{R}_{i_0 \dots i_l, \iota_0}^{(3l)}$ . Because  $\hat{\delta}\hat{\xi} = 0$  we see that the elements  $\hat{\xi}_{i_0 \dots i_l, \iota_0}$  are independent of  $\iota_0$ . Now  $\bigcup_{\iota=1}^{\iota^*} \hat{V}_{\iota}^{(3l)}$  covers  $X(\rho_1)$ . If we put  $\varepsilon_2(\hat{\xi})_{i_0 \dots i_l} = \hat{\xi}_{i_0 \dots i_l, \iota_0}$  in  $\hat{U}_{i_0 \dots i_l}^{(3l)} \cap \hat{V}_{\iota_0}^{(3l)}$  then we see that  $\varepsilon_2(\hat{\xi})_{i_0 \dots i_l}$  is a well defined section on  $\hat{U}_{i_0 \dots i_l}^{(3l)}$ . In this way we obtain  $\varepsilon_2(\hat{\xi}) \in Z^l(\hat{\mathfrak{U}}, \mathbf{F})$ . Here  $\varepsilon_2(\hat{\xi})$  is a cocycle because  $\hat{\delta}\hat{\xi} = 0$ . Now we prove that  $\| \varepsilon_2(\hat{\xi}) \|_{\rho} \leq K \| \hat{\xi} \|_{\rho}$ .

*Verification.* A computation of  $\| \varepsilon_2(\hat{\xi}) \|_{\rho}$  involves the following:  $\varepsilon_2(\hat{\xi}) = \{ \xi_{i_0 \dots i_l}^{(2)} \}$ . Look at some  $\xi_{i_0 \dots i_l}^{(2)}$  in the chart  $\mathcal{W}_i$  with  $i \in \{ i_0, \dots, i_l \}$ . We write  $\hat{\xi}_{i_0 \dots i_l}^{(2)} = \sum a_v (t/\rho)^v$  over  $(U_{i_0 \dots i_l}^*)_i$  and compute  $\sup_v | a_v(U_{i_0 \dots i_l}^*) |$ . A computation of  $\| \hat{\xi} \|_{\rho}$  involves the following: Look at  $\hat{\xi}_{i_0 \dots i_l}$  over  $(U_{i_0 \dots i_l}^* \cap V_{\iota}^*)_i$  in a chart  $W_i$ . Here  $\iota$  is fixed. We write  $\hat{\xi}_{i_0 \dots i_l, \iota} = \sum a_v^{(\iota)} (t/\rho)^v$  and compute  $\sup_{\iota^*} | a_v^{(\iota)}(U_{i_0 \dots i_l}^* \cap V_{\iota}^*) |$ . Now  $\bigcup_{\iota} V_{\iota}^*$  covers  $X_0$ . Hence we would have  $\sup_{v, \iota} | a_v^{(\iota)}(U_{i_0 \dots i_l}^* \cap V_{\iota}^*) | = \sup_v | a_v(U_{i_0 \dots i_l}^*) |$  if  $a_v = a_v^{(\iota)}$  in  $U_{i_0 \dots i_l}^* \cap V_{\iota}^*$ . But this is obvious since  $\xi_{i_0 \dots i_l}^{(2)} = \hat{\xi}_{i_0 \dots i_l, \iota}$  in  $(U_{i_0 \dots i_l}^* \cap V_{\iota}^*)_i$ . Hence we have  $\| \varepsilon_2(\hat{\xi}) \|_{\rho} \leq \| \hat{\xi} \|_{\rho}$ .

Now we are ready to start the proof of the smoothing theorem. We let  $K$  denote a constant, which may be different at different occurrences.

We also introduce a double complex  $\{ \tilde{C}_v^{k, \kappa} \}$  using  $(\mathfrak{B}, \mathfrak{B})$ , i.e. it is defined just as the previous double complex was, using  $\mathfrak{B}$ -sets instead of  $\mathfrak{U}$ -sets.

We shall inductively construct the following elements:

$$\hat{\xi}_v = \{ \hat{\xi}_{i_0 \dots i_v, \iota_0 \dots \iota_{l-v}} \} \in Z_{3v}^{v, l-v}$$

$$\tilde{\xi}_v = \{ \tilde{\xi}_{i_0 \dots i_v, \iota_0 \dots \iota_{l-v}} \} \in \tilde{Z}_{3v}^{v, l-v}; \quad v = 0, \dots, l$$

$$\hat{\eta}_v = \{ \hat{\eta}_{i_0 \dots i_{v-1}, \iota_0 \dots \iota_{l-v}} \} \in C_{3v}^{v-1, l-v}$$

$$\tilde{\eta}_v = \{ \tilde{\eta}_{i_0 \dots i_{v-1}, \iota_0 \dots \iota_{l-v}} \} \in \tilde{C}_{3v}^{v-1, v-1}; \quad v = 1, \dots, l$$

$$\tilde{\gamma}_v = \{ \tilde{\gamma}_{i_0 \dots i_{v-1}, i_0 \dots i_{l-v-1}} \} \in \tilde{C}_{3v-3}^{v-1, l-v-1}; \quad v = 1, \dots, (l-1)$$

and  $\tilde{\gamma}_l = \{ \tilde{\gamma}_{i_0 \dots i_{l-1}} \} \in C^{l-1}(\mathfrak{B}_{3l})$ .

The construction:  $\hat{\xi} \in Z^l(\hat{\mathfrak{B}}(\rho), \mathbb{F})$  is given. The whole construction is done using  $\rho$  instead of  $\rho_1$  and we omit  $\rho$  to simplify the notation. We put  $\varepsilon_1(\hat{\xi}) = \hat{\xi}_0 \in Z_0^{0,l}$ . Now we apply the Smoothing Lemma and get  $\hat{\eta}_1$  such that  $\partial \hat{\eta}_1 = \hat{\xi}_0$  with  $\| \hat{\eta}_1 \|_\rho \leq K \| \hat{\xi}_0 \|_\rho \leq K \| \hat{\xi} \|_\rho$ . Put  $\hat{\xi}_1 = \delta \hat{\eta}_1$ . Obviously  $\| \hat{\xi}_1 \|_\rho \leq K \| \hat{\eta}_1 \|_\rho$ . Inductively we find  $\delta \hat{\eta}_v = \hat{\xi}_{v-1}$  and we put  $\hat{\xi}_v = \delta \hat{\eta}_v$  where  $\hat{\eta}_v$  are found from the Smoothing Lemma. Finally we get  $\hat{\xi}_l$  and we have  $\| \hat{\xi}_l \|_\rho \leq K \| \hat{\xi} \|_\rho$ . Now we define  $\tilde{\xi}_v$  and  $\tilde{\eta}_v$  as follows. Put  $\tilde{\xi}_0 = \hat{\xi}_0$  where  $\tilde{\xi}_0 \in \tilde{Z}_0^{0,l}$  is obtained by natural restriction of  $\hat{\xi}_0$ . Put  $\tilde{\eta}_v = (-1)^v \{ \tilde{\xi}_{i_0 \dots i_{v-1}, i_0 \dots i_{l-v}} \}$  which is well defined with respect to  $(\mathfrak{B}_{3v}, \mathfrak{B}_{3v})$  by taking natural restrictions. Put  $\tilde{\xi}_v = \delta \tilde{\eta}_v$  for  $v = 1, \dots, l$ . A computation shows that  $\tilde{\xi}_{v-1} = \partial \tilde{\eta}_v$  when  $v = 1, \dots, l$ . Notice that this is trivial when  $v = 1$ . In the following discussion each  $\hat{\eta}_v$  is restricted to  $(\mathfrak{B}_{3v}, \mathfrak{B}_{3v})$ . We have  $\partial(\tilde{\eta}_1 - \hat{\eta}_1) = 0$ . Hence we find  $\tilde{\eta}_1 - \hat{\eta}_1 = \delta \tilde{\gamma}_1$  by the Smoothing Lemma. Now we define  $\tilde{\gamma}_v$  such that  $\partial \tilde{\gamma}_v = \tilde{\eta}_v - \hat{\eta}_v - \delta \tilde{\gamma}_{v-1}$  inductively. This is possible because  $\partial(\tilde{\eta}_v - \hat{\eta}_v - \delta \tilde{\gamma}_{v-1}) = 0$ , for we have  $\partial(\tilde{\eta}_v - \hat{\eta}_v - \delta \tilde{\gamma}_{v-1}) = \tilde{\xi}_{v-1} - \hat{\xi}_{v-1} - \delta \partial \tilde{\gamma}_{v-1} = \delta \tilde{\eta}_{v-1} - \delta \hat{\eta}_{v-1} - \delta(\tilde{\eta}_{v-1} - \hat{\eta}_{v-1}) = 0$ . We get finally  $\tilde{\gamma}_{l-1} \in \tilde{C}_{3l}^{l-2,0}$  and then  $\delta \tilde{\gamma}_{l-1} \in \tilde{C}_{3l}^{l-1,0}$ . We have  $\partial(\tilde{\eta}_l - \hat{\eta}_l - \delta \tilde{\gamma}_{l-1}) = 0$ . Therefore we can put  $\tilde{\gamma}_l = \varepsilon_2(\tilde{\eta}_l - \hat{\eta}_l - \delta \tilde{\gamma}_{l-1})$ . It follows that  $\tilde{\gamma}_l \in C^{l-1}(\mathfrak{B}_{3l})$  and  $\delta \tilde{\gamma}_l = \varepsilon_2(\tilde{\xi}_l - \hat{\xi}_l)$ . We have  $\varepsilon_2(\tilde{\xi}_l) = -\hat{\xi} | \mathfrak{B}'$  and for  $\varepsilon_2(\tilde{\xi}_l) = -\hat{\xi}^*$  and  $\hat{\eta} = \tilde{\gamma}_l$  the required equation  $\hat{\xi}^* = \hat{\xi} + \delta \hat{\eta}$ . The estimates follow immediately from the construction and the Smoothing Lemma.

APPROXIMATION

We use positive  $n$ -tuples  $\rho, \dots$  with  $\rho \leq \rho_2 < \rho_3 < \rho_4 < \rho_1$  and  $\rho = \gamma'' \rho_1, \rho_2 = \gamma \rho_1, \rho_3 = \gamma' \rho_1, \rho_4 = \gamma''' \rho_1$ . The  $n$ -tuple  $\rho_1$  is defined as in the smoothing theorem.

*Definition:*  $H_*^l = \{ \xi \in H^l(X_0, \underline{F}|X_0) \text{ such that there exists } U = U(0) \text{ in } E^n \text{ with } \hat{\xi} \in H^l(\psi^{-1}(U), \mathbf{F}) \text{ and } \hat{\xi}|X_0 = \xi \}$ . Serre's theorem gives  $\dim_{\mathbf{C}} H_*^l \leq \dim_{\mathbf{C}} H^l(X_0, \underline{F}|X_0) < \infty$ . In the following discussion we are given  $\hat{b}_1, \dots, \hat{b}_r$  in  $Z^l(\hat{\mathcal{U}}'(\rho_4), \mathbf{F})$  such that  $\hat{b}_1|X_0, \dots, \hat{b}_r|X_0$  constitute a base of the complex vector space  $H_*^l$ . For this to be possible,  $\rho_4$  has to be chosen small enough. Here  $\hat{\mathcal{U}}'$  is a Stein covering of  $X(\rho_1)$  and defined as in the smoothing theorem. We also assume that we are given a sequence of measure coverings as there. Further we construct the sequence so that there are still sufficiently many measure coverings in between  $\mathfrak{B}$  and  $\mathcal{U}$ . These are denoted by  $\mathcal{U}_v^*$ . We have  $\mathcal{U} \gg \mathcal{U}_1^* \gg \mathcal{U}_2^* \gg \dots \gg \mathfrak{B}$ . The  $n$ -tuple  $\rho_3$  is also fixed from now on and  $K$  always denotes (possibly different) constants.

*Approximation Lemma:* Let  $\varepsilon > 0$ . Then we can find  $\rho_2$  such that: If  $\rho \leq \rho_2$  and  $\hat{\xi} \in Z^l(\hat{\mathcal{U}}(\rho), \mathbf{F})$  with  $\|\hat{\xi}\|_{\rho} < \infty$  (the norm is taken with respect to  $\hat{\mathcal{U}}_1^*(\rho)$ ), then there exist  $a_1, \dots, a_r \in I(E^n(\rho))$  and  $\hat{\eta} \in C^{l-1}(\hat{\mathfrak{B}}(\rho), \mathbf{F})$  such that  $\tilde{\xi} = \hat{\xi} - \sum_1^r a_i \hat{b}_i - \delta \eta$  on  $\mathfrak{B}(\rho)$ . Here  $\tilde{\xi} \in Z^l(\mathfrak{B}(\rho), \mathbf{F})$  and  $\|\tilde{\xi}\|_{\rho} \leq \varepsilon \|\hat{\xi}\|_{\rho}$  and  $\|a_v\|_{\rho}, \|\hat{\eta}\|_{\rho} \leq K \|\hat{\xi}\|_{\rho}$ .  $K$  is a fixed constant.

*Proof.* We shall first prove some results which are needed later on. Let  $S \in \Gamma(\hat{U}_{\iota_0 \dots \iota_l}(\rho), \mathbf{F})$ . Choose  $\iota \in \{ \iota_0, \dots, \iota_l \}$ . Now  $(U_{\iota_0 \dots \iota_l}^{(1)*})_{\iota} \subset \hat{U}_{\iota_0 \dots \iota_l}$  because  $\mathcal{U}_1^* \ll \mathcal{U}$ . The operations are always defined with respect to  $\rho_1$ . We can now restrict  $S$  to  $(U_{\iota_0 \dots \iota_l}^{(1)*})_{\iota}(\rho)$ . In the chart  $\mathcal{W}_{\iota}$  we can write  $S = \sum a_v (t/\rho)^v$ . Here  $a_v \in qI(U_{\iota_0 \dots \iota_l}^{(1)*})$ . Now the  $a_v$  are extended constantly and we get elements  $\hat{a}_v \in \Gamma((U_{\iota_0 \dots \iota_l}^{(1)*})_{\iota}, \mathbf{F})$ . Let us put  $S_v = \hat{a}_v | \hat{U}_{\iota_0 \dots \iota_l}^{(2)*}$ . We claim that  $\|S_v\|_{\rho_1} \leq K \|S\|_{\rho}$ . For obviously  $\|S\|_{\rho} \geq |a_v (U_{\iota_0 \dots \iota_l}^{(1)*})|$  and

we can use the Theorem I to prove that  $\|S_v\|_{\rho_1} \leq K \|\hat{a}_v\| (U_{\iota_0 \dots \iota_l}^{(1)*}(\rho_1))\|_{\iota} = K \|a_v(U_{\iota_0 \dots \iota_l}^{(1)*})\| \leq K \|S\|_{\rho}$ . Q.E.D.

Let  $S'_v$  be defined using some other  $\iota' \in \{\iota_0, \dots, \iota_l\}$ . Then  $S_v - S'_v \in \Gamma(\hat{U}_{\iota_0 \dots \iota_l}^{(2)*}, \mathbf{F})$ . We claim that  $\|S_v - S'_v\|_{\rho_4} \leq K \gamma''' \|S\|_{\rho}$ .

*Proof.* Define  $\alpha_s = \sum_{|\lambda|=s} a_{\lambda}(t/\rho)^{\lambda}$  and  $\beta_s = \sum_{|\lambda|=0}^{s-1} a_{\lambda}(t/\rho)^{\lambda}$  over  $(U_{\iota_0 \dots \iota_l}^{(1)*})_{\iota}(\rho)$ . We do the same for  $\iota'$  respectively and obtain  $\alpha'_s$  and  $\beta'_s$  over  $(U_{\iota_0 \dots \iota_l}^{(1)*})_{\iota'}(\rho)$ . For the restrictions to  $\hat{U}_{\iota_0 \dots \iota_l}^{(2)*}$  we see that  $\alpha_s - \alpha'_s = -(\beta_s - \beta'_s)$ . Hence we get  $\|\alpha_s - \alpha'_s\|_{\rho_4} \leq K(\gamma''')^s \|\alpha_s - \alpha'_s\|_{\rho_1} = K(\gamma''')^s \|\beta_s - \beta'_s\|_{\rho_1} \leq K(\gamma''')^s \|\beta_s\|_{\rho_1} + K(\gamma''')^s \|\beta'_s\|_{\rho_1} \leq K(\gamma''')^s [\|\beta_s\|_{\rho_1}^* + \|\beta'_s\|_{\rho_1}^*] \leq K(\gamma''')^s (\gamma'')^{1-s} \|S\|_{\rho}$ . Here the norms are defined with respect to  $U_{\iota_0 \dots \iota_l}^{(3)*}$  except  $\|\cdot\|_{\rho_1}^*$  and  $\|S\|_{\rho}$  which are defined with respect to  $U_{\iota_0 \dots \iota_l}^{(1)*}$ . Now we look at the difference  $(S_v - S'_v) t^v/\rho^v$  on  $(U_{\iota_0 \dots \iota_l}^{(3)*})_{\mu}$  with  $|v|=s$ ,  $\mu \in \{\iota_0, \dots, \iota_l\}$ , and the power series development with respect to  $W_{\mu}$ . There is one term of order  $s$  which is equal to the corresponding term of  $\alpha_s - \alpha'_s$ . Therefore its norm is  $\leq K(\gamma''')^s \cdot (\gamma'')^{1-s} \|S\|_{\rho}$ . Moreover we have  $\|S_v(t/\rho)^v - S'_v(t/\rho)^v\|_{\rho_1} \leq (\gamma'')^{-s} \cdot K \|S\|_{\rho}$  where the first norm is defined with respect to  $U_{\iota_0 \dots \iota_l}^{(3)*}$ . For the sum  $\sum$  of terms of higher order than  $s$  in the power series of  $(S_v - S'_v) t^v/\rho^v$  we therefore get:  $\|\sum\|_{\rho_4} \leq (\gamma''')^{s+1} (\gamma'')^{-s} \cdot K \|S\|_{\rho}$ . Hence we get  $\|(S_v - S'_v)\|_{\rho_4} \leq \gamma''' \cdot K \|S\|_{\rho}$ . This proves our statement. We see that  $K$  is independent of  $\rho_4$  and  $S$ . The number  $\gamma'''$  depends on  $\rho_4$  only, so  $\gamma''' \cdot K$  gets very small if we make  $\rho_4$  very small.

Let  $\hat{\xi} \in Z^l(\hat{\mathcal{U}}(\rho), \mathbf{F})$  with  $\hat{\xi} = \{\hat{\xi}_{\iota_0 \dots \iota_l}\}$ . Choose  $\iota = \iota(\iota_0, \dots, \iota_l)$  as a function of the unordered  $(l+1)$ -tuple. We now fix  $\iota_0, \dots, \iota_l$  and write  $S = \hat{\xi}_{\iota_0 \dots \iota_l}$ . We apply to  $S$  the method described above and obtain  $\hat{\xi}_{\iota_0 \dots \iota_l}^{(v)} = S_v$ . We do this now for every  $\iota_0, \dots, \iota_l$  and consider  $\hat{\xi}_{(v)} = \{\hat{\xi}_{\iota_0 \dots \iota_l}^{(v)}\}$  as an element of  $C^l(\hat{\mathcal{U}}_2^*(\rho_4), \mathbf{F})$ . Of course  $\hat{\xi}_{(v)}$  depends on the choice of  $\iota = \iota(\iota_0 \dots \iota_l)$  here. Now we see that  $\|\hat{\xi}_{(v)}\|_{\rho_4} \leq \|\hat{\xi}_{(v)}\|_{\rho_1} \leq K \|\hat{\xi}\|_{\rho}$ . We also wish to estimate  $\delta \hat{\xi}_{(v)}$ . Because  $\hat{\xi} \in Z^l(\hat{\mathcal{U}}(\rho), \mathbf{F})$  we can use the preliminary result on  $\iota$  and  $\iota'$  to obtain  $\|\delta \hat{\xi}_{(v)}\|_{\rho_4} \leq K \gamma''' \|\hat{\xi}\|_{\rho}$ .

We shall also need another result:



*Induction Lemma:* There exists  $\hat{\eta}_v \in C^l(\hat{\mathcal{U}}_4^*(\rho_3), \mathbf{F})$  such that  $\delta\hat{\eta}_v = \delta\hat{\xi}_{(v)}$  on  $\hat{\mathcal{U}}_4^*(\rho_3)$  and  $\|\hat{\eta}_v\|_{\rho_3} \leq K \|\delta\hat{\xi}_{(v)}\|_{\rho_4}$ .

*Proof.* The proof uses the assumption that  $\psi_{(l+1)}(\mathbf{F})$  is coherent. Because the coherence of direct images is proved by downward induction on  $l$ , this assumption can be made. Moreover it is assumed that the main theorem is proved for dimension  $l + 1$  already. Let us now put  $\alpha = \delta\hat{\xi}_{(v)} \in B^{l+1}(\hat{\mathcal{U}}_2^*(\rho_4), \mathbf{F})$  and  $\hat{\eta}_v = \beta \in C^l(\hat{\mathcal{U}}_4^*(\rho_3), \mathbf{F})$ . We have to prove the existence of  $\beta$ . We may assume that  $\rho_4$  is so small that the main theorem is valid for  $\rho \leq \rho_4$  in the case of dimension  $l + 1$ . So there are cocycles  $\omega_1, \dots, \omega_r \in Z^{l+1}(\hat{\mathcal{U}}(\rho_4), \mathbf{F})$  such that  $\alpha = \sum C_\lambda \omega_\lambda + \delta\eta$ , where  $C_\lambda \in I(E^n(\rho_4))$  and  $\eta \in C^l(\hat{\mathcal{U}}_4^*(\rho_4), \mathbf{F})$ . We have to assume that between  $\hat{\mathcal{U}}_4^*$  and  $\mathcal{U}_2^*$  there are very many measure coverings. The cross-sections  $\psi_{(l+1)}(\omega_\lambda)$  give a homomorphism  $r\mathcal{O} \rightarrow \psi_{(l+1)}(\mathbf{F})$  over  $E^n(\rho_4)$ . Because  $\psi_{(l+1)}(\mathbf{F})$  is coherent the kernel  $\mathcal{N}$  is coherent again. Over  $E^n(\rho')$  with  $\rho_3 < \rho' < \rho_4$  we find an epimorphism  $p\mathcal{O} \rightarrow \mathcal{N}$ . Denote by  $n_1, \dots, n_p$  the images of the unit cross-sections in  $p\mathcal{O}$ . Write  $n_\lambda = (e_{\lambda 1}, \dots, e_{\lambda r})$  as an  $r$ -tupel of holomorphic functions. The image of  $n_\lambda$  in  $\Gamma(E^n(\rho'), \psi_{(l+1)}(\mathbf{F}))$  is  $\psi_{(l+1)}(\sum_{\mu=1}^r e_{\lambda\mu} \omega_\mu)$  and zero. We may choose  $\rho_2$  and then  $\rho_3$  and  $\rho'$  very small. Then it follows that  $\hat{n}_\lambda = \sum e_{\lambda\mu} \omega_\mu$  is a coboundary. If  $\rho_3 < \rho'' < \rho'$  there are cochains  $\eta_\lambda \in C^l(\hat{\mathcal{U}}_4^*(\rho''), \mathbf{F})$  such that  $\delta\eta_\lambda = \hat{n}_\lambda$ . Now  $(C_1, \dots, C_r) \in \Gamma(E^n(\rho_4), \mathcal{N})$ . By the methods of sheaf theory we can lift this cross-section to  $p\mathcal{O}$ . Using a "Banach open mapping theorem" we see that the map  $\Gamma(E^n(\rho'), p\mathcal{O}) \rightarrow \Gamma(E^n(\rho'), \mathcal{N})$  is open. This means here that we can find holomorphic functions  $a_\lambda$  over  $E^n(\rho_3)$  such that  $C_\mu = \sum a_\lambda e_{\lambda\mu}$  and  $\|a_\lambda\|_{\rho_3} \leq K \max_\mu \|C_\mu\|_{\rho'} \leq K \max_\mu \|C_\mu\|_{\rho_4}$ . We get  $\sum C_\mu \omega_\mu = \sum a_\lambda e_{\lambda\mu} \omega_\mu = \sum a_\lambda \hat{n}_\lambda = \delta(\sum a_\lambda \eta_\lambda)$ . This leads to  $\alpha \in C^{l+1}(\hat{\mathcal{U}}_4^*(\rho_3)) = \delta(\eta + \sum a_\lambda \eta_\lambda)$ . The estimates required obviously hold. Q.E.D.

Let us now put  $\hat{\xi}_{(v)}^* = \hat{\xi}_{(v)} - \hat{\eta}_v \in Z^l(\hat{\mathcal{U}}_4(\rho_3), \mathbf{F})$ . We can write  $\hat{\xi}_{(v)}^* \mid X_0 = \sum a_{v\lambda} \hat{b}_\lambda \mid X_0 + \delta\gamma_v$  over  $\mathcal{U}_6^*$ . Here  $a_{v\lambda}$  are complex numbers and  $\gamma_v \in C^{l-1}(\mathcal{U}_6^*, \mathbf{F} \mid X_0)$ . Cartan's theorem and the result after that give the estimates  $|a_{v\lambda}| \leq K \|\hat{\xi}_{(v)}^*\|_{\rho_3} \leq K \|\hat{\xi}\|_\rho$  and  $\|\gamma_v\|_{\rho_3} \leq K \|\hat{\xi}_{(v)}^*\|_{\rho_3} \leq K \|\hat{\xi}\|_\rho$ . Here  $\hat{\gamma}_v \in C^{l-1}(\hat{\mathcal{U}}_7^*(\rho_3), \mathbf{F})$  has been obtained by a constant



extension of  $\gamma_v$ . Let us now put  $\hat{\xi}_{(v)}^{(1)} = \hat{\xi}_{(v)}^* - \sum a_{v\lambda} \hat{b}_\lambda - \delta \hat{\gamma}_v$ . Here  $\hat{\xi}_{(v)}^{(1)} \in C^l(\hat{\mathfrak{U}}_7^*(\rho_3), \mathbb{F})$ . Using the previous estimates and the fact that the  $\hat{b}_\lambda$  are finite we find that  $\|\hat{\xi}_{(v)}^{(1)}\|_{\rho_3} \leq K \|\hat{\xi}_{(v)}\|_{\rho_4} \leq K \|\hat{\xi}\|_{\rho}$ .

Now we also have  $\hat{\xi}_{(v)}^{(1)}|_{X_0} = 0$ . It follows that

$$\|\hat{\xi}_{(v)}^{(1)}\|_{\rho} \leq \gamma/\gamma' \|\hat{\xi}_{(v)}^{(1)}\|_{\rho_3} \leq \gamma/\gamma' \cdot K \|\hat{\xi}\|_{\rho}.$$

Finally we put in  $\hat{\mathfrak{U}}_9^*(\rho)$ :

$$\begin{aligned} \hat{\xi}^{(1)} &= \sum \hat{\xi}_{(v)}^{(1)} (t/\rho)^v = \\ &= \sum \hat{\xi}_{(v)} (t/\rho)^v - \sum \hat{\eta}_v (t/\rho)^v - \sum a_{v\lambda} (t/\rho)^v \hat{b}_\lambda - \delta (\sum \hat{\gamma}_v (t/\rho)^v) \\ &= \hat{\xi} - \hat{\eta} - \sum a_\lambda \hat{b}_\lambda - \delta \hat{\gamma}. \end{aligned}$$

Using the fact that the sum of the absolute values of the coefficients in the power series expansion of  $\hat{\xi}_{(v)}^{(1)}$  by  $(t/\rho)$  is smaller than  $\gamma/\gamma' \cdot K \|\hat{\xi}\|_{\rho}$  and that with respect to  $\hat{\eta}_v$  is smaller than  $\gamma''' \cdot K \|\hat{\xi}\|_{\rho}$  we find:  $\|\hat{\xi}^{(1)}\|_{\rho} \leq \gamma/\gamma' \cdot K \|\hat{\xi}\|_{\rho}$  and  $\|\hat{\eta}\|_{\rho} \leq \gamma''' \cdot K \|\hat{\xi}\|_{\rho}$  and  $\|a_\lambda\|_{\rho} \leq K \|\hat{\xi}\|_{\rho}$ . We take the restriction to  $\hat{\mathfrak{B}}(\rho)$  and now  $\tilde{\xi} = \hat{\xi}^{(1)} - \hat{\eta} \in Z^l(\hat{\mathfrak{B}}(\rho), \mathbb{F})$  is the desired element. Of course we have to choose  $\rho_4$  and then  $\rho_2$  small enough, for example let  $\gamma''' < \varepsilon/2 K$  and  $\gamma \leq \varepsilon\gamma'/2 K$ .

### MAIN THEOREM

There exists  $\rho_2$  and a constant  $K$  such that if  $\rho \leq \rho_2$  and  $\hat{\xi} \in Z^l(\hat{\mathfrak{U}}(\rho), \mathbb{F})$  with  $\|\hat{\xi}\|_{\rho} < \infty$  then we can find  $a_1, \dots, a_r \in I(E^n(\rho))$  and  $\hat{\eta} \in C^{l-1}(\hat{\mathfrak{B}}(\rho), \mathbb{F})$  such that  $\hat{\xi} = \sum a_\lambda \hat{b}_\lambda + \delta \hat{\eta}$  on  $\hat{\mathfrak{B}}(\rho)$  with  $\|\hat{\eta}\|_{\rho}$  and  $\|a_v\|_{\rho} \leq K \|\hat{\xi}\|_{\rho}$ .

*Proof.* We have one constant  $K$  from the smoothing theorem. Now we find  $\rho_2$  with an  $\varepsilon$  in the Approximation Lemma such that  $\varepsilon \cdot K < 1/2$ . We shall use this  $\rho_2$  and prove the theorem here. We are given  $\hat{\xi}_0 = \hat{\xi} \in Z^l(\hat{\mathfrak{U}}(\rho), \mathbb{F})$  with  $\|\hat{\xi}\|_{\rho} < \infty$ . The Approximation Lemma gives  $\tilde{\xi}_1 =$

$= \hat{\xi} - \sum a_{1\lambda} \hat{\mathbf{b}}_\lambda - \hat{\delta}\gamma_1$  on  $\hat{\mathfrak{B}}(\rho)$ . Here  $\hat{\gamma}_1 \in C^{l-1}(\hat{\mathfrak{B}}(\rho), \mathbf{F})$  and  $\|\hat{\xi}_1\|_\rho \leq \varepsilon \|\hat{\xi}\|_\rho$ . Now  $\hat{\xi}_1 \in Z^l(\hat{\mathfrak{B}}(\rho), \mathbf{F})$ . The Smoothing Theorem gives  $\hat{\xi}_1 \in Z^l(\hat{\mathfrak{U}}(\rho), \mathbf{F})$  and  $\hat{\eta}_1 \in C^{l-1}(\hat{\mathfrak{B}}(\rho), \mathbf{F})$  such that  $\hat{\xi}_1 = \tilde{\xi}_1 + \hat{\delta}\eta_1$  on  $\hat{\mathfrak{B}}(\rho)$ . Here  $\|\eta_1\|_\rho$  and  $\|\hat{\xi}_1\|_\rho \leq K \|\tilde{\xi}_1\|_\rho < 1/2 \|\hat{\xi}\|_\rho$ . Now we use  $\hat{\xi}_1$  instead of  $\hat{\xi}_0$  as above and get:  $\hat{\xi}_2 = \hat{\xi}_1 + \hat{\delta}\eta_2 - \sum a_{2\lambda} \hat{\mathbf{b}}_\lambda - \hat{\delta}\gamma_2$ . Here  $\|\hat{\xi}_2\|_\rho$  and  $\|\hat{\eta}_2\|_\rho < 1/2 \|\hat{\xi}_1\|_\rho < (1/2)^2 \|\hat{\xi}\|_\rho$  and  $\|a_{2\lambda}\|_\rho$  and  $\|\gamma_2\|_\rho \leq \frac{K}{2} \|\hat{\xi}\|_\rho$ . Inductively we get:  $\hat{\xi}_n = \hat{\xi}_{n-1} - \sum a_{n\lambda} \hat{\mathbf{b}}_\lambda - \hat{\delta}\gamma_n + \hat{\delta}\eta_n$ . Here  $\|\hat{\xi}_n\|_\rho < 2^{-n} \|\hat{\xi}\|_\rho$ ,  $\|\hat{\eta}_n\|_\rho \leq 2^{-n} \|\hat{\xi}\|_\rho$  and  $\|a_{n\lambda}\|_\rho$  and  $\|\gamma_n\|_\rho \leq 2^{-n+1} \cdot K \|\hat{\xi}\|_\rho$  for  $n = 1, 2, 3, \dots$ . A summation is now possible. We get  $0 = \hat{\xi} - \sum_{n,\lambda} a_{n\lambda} \hat{\mathbf{b}}_\lambda - \sum \hat{\delta}\gamma_n + \sum \hat{\delta}\eta_n$ . We put  $a_\lambda = \sum_n a_{n\lambda}$ ,  $\hat{\eta} = \sum (-\hat{\gamma}_n + \hat{\eta}_n)$  and the theorem follows.

For the proof of the coherence the Main Theorem is needed in a weaker and simpler form.

*Main Theorem (\*)*: There exists a positive  $n$ -tuple  $\rho_2 \leq \rho_0$  and cross-sections  $S_1, \dots, S_r \in \Gamma(E^n(\rho_2), \psi_{(l)}(\mathbf{F}))$  such that any  $S = \psi_{(l)}(\hat{\xi}') \in \Gamma(E^n(\rho'), \psi_{(l)}(\mathbf{F}))$  with  $\hat{\xi}' \in H^l(X(\rho'), \mathbf{F})$  can be written over  $E^n(\rho)$  in the form  $S = \sum_1^n a_\lambda S_\lambda$  with  $a_1, \dots, a_r \in I(E^n(\rho))$ . Here  $\rho \leq \rho_2$  and  $\rho < \rho' \leq \rho_0$ .

*Proof.* Define  $S_\lambda = \psi_{(l)}(\hat{\mathbf{b}}_\lambda | X(\rho_2))$ . The cross-section  $S$  can be written in the form  $S = \psi_{(l)}(\hat{\xi}')$  with  $\hat{\xi}' \in Z^l(\hat{\mathfrak{U}}'(\rho'), \mathbf{F})$ . We put  $\hat{\xi} = \hat{\xi}' | \mathfrak{U}(\rho)$ . Then  $\|\hat{\xi}\|_\rho < \infty$  and we have the representation  $\hat{\xi} = \sum a_\lambda \hat{\mathbf{b}}_\lambda + \hat{\delta}\eta$ . For the cohomology classes we get  $\hat{\xi} = \sum a_\lambda \hat{\mathbf{b}}_\lambda$  and for the images  $S | E^n(\rho)$ , this gives  $S | E^n(\rho) = \psi_{(l)}(\hat{\xi}) = \sum a_\lambda S_\lambda$ .

The immediate consequence of this form of the Main Theorem is that the stalk of  $\psi_{(l)}(\mathbf{F})$  at the origin (and hence at every point of course) is finitely generated. However this is not yet the full coherence of  $\psi_{(l)}(\mathbf{F})$ . Nevertheless, the Main Theorem above contains all that is essential, and the rest of the proof is not difficult. We refer to [1, pp. 54-58], or to Knorr [2] for details.

REFERENCES

- [1] GRAUERT, H., *Ein Theorem der analytischen Garbentheorie und die Modulräume komplexer Strukturen*. Publ. math. n° 5 de l'Inst. des Hautes Etudes Scientifiques, Paris, 1960.
- [2] KNORR, K., *Über die Kohärenz von Bildgarben bei eigentlichen Abbildungen in der analytischen Geometrie*. *Ann. Scuola Norm. Pisa*, 1968/69.