

# §5. The Moduli Space of Stable Curves

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

§ 5. THE MODULI SPACE OF STABLE CURVES

Our main result is:

**THEOREM 5.1.** *Fix  $n \geq 5$ , and for any curve  $C$  of genus  $g$  let  $\Phi_n(C) \subset \mathbf{P}^{(2n-1)(g-1)-1}$  be the image of  $C$  embedded by a basis of  $\Gamma(C, \omega_C^{\otimes n})$ . Then if  $C$  is moduli-stable,  $\Phi_n(C)$  is Chow stable.*

In view of the basic results of § 1, and those of [20], this shows:

**COROLLARY 5.2.** *(F. Knudsen)  $\bar{\mathcal{M}}_g$  is a projective variety.*


Recall that  $C$  moduli-stable means

- (1)  $C$  has at worst ordinary double points (by Proposition 3.12, this is necessary for the asymptotic semi-stability of  $C$ ) and is connected,
- (2)  $C$  has no smooth rational components meeting the rest of the curve in fewer than three points:  
this condition is necessary to ensure that  $C$  has only finitely many automorphisms.

We will call  $C$  moduli semi-stable if it satisfies (1) and

- (2')  $C$  has no smooth rational components meeting the rest of the curve in only one point.

Note that if  $C$  is moduli semi-stable, then the set of its smooth rational components meeting the rest of the curve in exactly 2 points form a finite set of chains and if each of these is replaced by a point, we get a moduli stable curve:



We will case these the rational chains of  $C$ .

It would be more satisfactory to have a direct proof of Theorem 5.1 similar to the proof of the stability of smooth curves given in § 4. But curves with double points are not usually linearly stable (cf. the remark following Theorem 4.14) and, in fact, the estimates in Corollary 4.11 do not suffice to prove stability for such curves. We will therefore take an indirect approach.

*Proof of 5.1.* We begin by recalling the useful valuative criterion:

LEMMA 5.3. *Suppose a reductive group  $G$  acts on a  $k$ -vector space  $V$ . Let  $K = k((t))$  and suppose  $x \in V_K$  is  $G$ -stable. Then there is a finite extension  $K' = k'((t')) \supset K$ , and elements  $g \in G_{K'}$ ,  $\lambda \in (K')^*$  such that the point  $\lambda g(x) \in V \otimes_k K'$  lies in  $V \otimes_k k'[[t']]$  and specializes as  $t \rightarrow 0$  to a point  $\overline{\lambda g(x)}$  with closed orbit. Thus  $\overline{\lambda g(x)}$  is either stable or semi-stable with a positive dimensional stabilizer.*

*Proof.* The diagram below is defined over  $k$ :

$$\begin{array}{c} \mathbf{P}(V) \supset \mathbf{P}(V)_{ss} \\ \downarrow \pi \\ X = \text{Proj}(\text{graded ring of invariants on } V) \end{array}$$

The point  $\pi(x) \in X_K$  specializes to a point  $\overline{\pi(x)} \in X_k$ . Let  $\bar{y}$  be a lifting of this point to  $V_{ss}$  with  $O^G(\bar{y})$  closed. In the scheme  $V \times \text{Spec } k[[t]]$  form the closure  $Z$  of  $\overline{\mathbf{G}_m \cdot O^G(x)}$ . The lemma follows if we prove that  $\bar{y} \in Z$ . If  $\bar{y} \notin Z$ , then  $Z$  and  $O^G(\bar{y})$  are closed disjoint  $G$  invariant subsets of  $V \times \text{Spec } k[[t]]$ , hence there exists a homogeneous  $G$ -invariant  $f$  such that  $f(x) = 0$  but  $f(\bar{y}) \neq 0$ . Then for some  $n$ ,  $f^{\otimes n}$  descends to a section of some line bundle on  $X \times \text{Spec } k[[t]]$ . But then  $f(\pi(x)) = 0$  and  $\overline{f(\pi(x))} \neq 0$  are contradictory.

Now suppose that  $C$  is a moduli stable curve of genus  $g$  over  $k$ . Let  $\mathcal{C}/k[[t]]$  be a family of curves with fibre  $C_0$  over  $t = 0$  equal to  $C$  and generic fibre  $C_\eta$  smooth. At the double points of  $C_0$ ,  $\mathcal{C}$  looks formally like  $xy = t^n$ , that is has only  $A_{n-1}$ -type singularities and hence is normal. Embed  $C_\eta$  in  $\mathbf{P}^N$  ( $N = (2n-1)(g-1)-1$ ) by  $\Gamma(C_\eta \omega_{C_\eta}^{\otimes n})$  and let  $\Phi(C_\eta)$  denote its image there. Then Lemma 5.3 says that by replacing  $k[[t]]$  with some finite extension and choosing a suitable basis of  $\Gamma(C_\eta, \omega_{C_\eta}^{\otimes n})$ —this

corresponds to choosing  $g, \lambda$ —we may assume that the closure  $\mathcal{D}$  in  $\mathbf{P}^N \times \text{Spec } k[[t]]$  of  $\Phi(C_\eta)$  satisfies

- i)  $D_\eta = C_\eta$
- ii)  $D_0$  Chow-stable or Chow semi-stable with positive dimensional stabilizer.



I now claim:

(5.4)  $\mathcal{D} = \Phi(\mathcal{C})$ , the image of  $\mathcal{C}$  under a  $k[[t]]$  basis of

$$\Gamma(\mathcal{C}, \omega_{\mathcal{C}}^n / k[[t]])$$

In particular this implies  $D_0 = C_0 = C$  and since  $C$  has finite stabilizer this means  $D_0$ , hence  $C$ , is Chow stable.

The main step in the proof of (5.4) is to show that  $D_0$  is moduli semi-stable as a scheme, and the key difficulty in doing this is to show that  $D_0$  has only ordinary double points. At first glance, this seems rather obvious, since from Proposition 3.12 it follows easily that as a cycle  $D_0$  has no multiplicities and has only ordinary double points. But ordinary double points on a limit cycle arise in two ways:



In the second case the scheme  $D_0$  has an embedded component (the first order normal neighbourhood in the  $z$ -direction) at the double point so in the limit scheme the double point is not ordinary. If case (ii) occurred for  $D_0$ , then since  $D_0$  is Chow semi-stable, it must span  $\mathbf{P}^N$  set-theoretically. But  $\Gamma(D_0, \mathcal{O}_{D_0}(1))$  has a torsion section supported at the double point: so  $D_0$  would have to be embedded by a non-complete linear system  $\Sigma \subset \Gamma(D_0, \mathcal{O}_{D_0}(1))$  of torsion-free sections,  $\dim \Sigma = \dim H^0(D_\eta, \mathcal{O}_{D_\eta}(1))$ . Consequently  $H^1(D_0, \mathcal{O}_{D_0}(1)) \neq (0)$  too. That this cannot happen in the situation of (5.4) follows from:

PROPOSITION 5.5. *Let  $C \subset \mathbf{P}^n$  be a 1-dimensional scheme such that*

- a)  $n + 1 = \deg C + \chi(\mathcal{O}_C)$ ,  $\chi(\mathcal{O}_C) < 0$ ,
- b)  $C$  is Chow semi-stable,
- c)  $\frac{\deg C}{n + 1} < \frac{8}{7}$ .

Then i)  $C$  is embedded by a complete non-special <sup>1)</sup> linear system,

ii)  $C$  is a moduli semi-stable curve with rational chains of length at most one consisting of straight lines.

Moreover if  $v = \frac{\deg C}{\deg \omega_C}$  (where  $\omega_C$  is the Grothendieck dualizing sheaf) and

$C = C_1 \cup C_2$  is a decomposition of  $C$  into two sets of components such that  $\mathcal{W} = C_1 \cap C_2$  and  $w = \# \mathcal{W}$  then

$$\text{iii) } \quad | \deg C_1 - v \deg_{C_1}(\omega_C) | \leq \frac{w}{2}$$

REMARKS. 1) It is clear that  $D_0$  satisfies the hypotheses of the lemma. Indeed a) is satisfied by  $D_\eta$  and is preserved under specialization. The key point of the Proposition to replace this by the stronger condition i)

2) Roughly, iii) says that the degrees of the components of  $C$  are roughly in proportion to their "natural" degrees. We will see later on that this is enough to force  $\mathcal{D} = \mathcal{C}$ .

*Proof.* From b), c) and Proposition 3.1 we know that the cycle of  $C$  has no multiplicity and only ordinary double points. Hence  $C_{\text{red}}$  is a scheme

<sup>1)</sup> Non-special means  $H^1(C, \mathcal{O}_C(1)) = (0)$ .



so by (5.7)

$$\deg C_1 \leq \frac{8}{7} h^0(C_1, \mathcal{O}_{C_1}(1)) \leq \frac{8}{14} \deg C_1 + \frac{8}{7},$$

which implies  $\deg C_1 \leq 2$ , hence  $C_1$  is rational and then  $H^1(C_1, \mathcal{O}_{C_1}(1)) = (0)$  anyway.

b)  $H^1(C_1, \mathcal{O}_{C_1}(1)(-\mathcal{W})) = (0)$ : indeed from (5.7) and Riemann-Roch,

$$\deg C_1 + \frac{1}{2} w \leq \frac{8}{7} (\deg C_1 - g_1 + 1), \text{ whence}$$

$$\deg \mathcal{O}_{C_1}(1)(-\mathcal{W}) = \deg C_1 - w \geq 8(g_1 - 1) + \frac{5}{2} w.$$

The last expression is greater than  $2g_1 - 2$  unless  $w = 0$ , when b) reduces to a), or  $g_1 = 0$  and  $w = 1$  or  $2$ . But in this case  $\mathcal{O}_{C_1}(1)(-\mathcal{W}) = \mathcal{O}_{\mathbb{P}^1}(e)$ , with  $e \geq 1 - 2 = -1$ .

Together a) and b) imply  $H^1(C, \mathcal{O}_C(1)) = 0$ . In fact, if  $C_{\text{red}}$  has components  $C_i$ , then there is an exact sequence

$$0 \rightarrow \bigoplus \mathcal{O}_{C_i}(1)(-\mathcal{W}_i) \rightarrow \mathcal{O}_{C_{\text{red}}}(1) \rightarrow \mathcal{M} \rightarrow 0$$

where  $\mathcal{M}$  has 0-dimensional support, hence  $H^1(C_{\text{red}}, \mathcal{O}_{C_{\text{red}}}(1)) = 0$ , and if  $\mathcal{N}$  is the sheaf of nilpotents in  $\mathcal{O}_C$ , then  $\mathcal{N}$  has 0-dimensional support and the conclusion follows from an examination of the exact sequence

$$0 \rightarrow \mathcal{N} \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_{C_{\text{red}}} \rightarrow 0.$$

Therefore hypothesis (a) can be rewritten  $n + 1 = h^0(\mathcal{O}_C(1))$ . Since  $C$  is not contained in a hyperplane,  $C$  is embedded by a complete linear system. But now if  $\mathcal{N} \neq (0)$ , then set-theoretically  $C$  will still be contained in a hyperplane, contradicting its Chow semi-stability; so  $C = C_{\text{red}}$  and all that we have said about  $C_{\text{red}}$  above is true of  $C$ .

Using the fact that

$$\chi(\mathcal{O}_C) = -\chi(\omega_C) = -(\deg \omega_C + \chi(\mathcal{O}_C))$$

it follows that  $\deg C/n + 1 = 2v/2v - 1$  and we can rewrite (5.7) in terms of  $v$  as

$$\frac{w}{2} + \deg C_1 \leq \left( \frac{2v}{2v-1} \right) (\deg C_1 - g_1 + 1)$$

or equivalently

$$\frac{w}{2} \cong v(2g_1 - 2 + w) - \deg C_1 = v \deg_{C_1}(\omega_C) - \deg C_1.$$


Then since

$$\begin{aligned} 0 &= v(\deg(\omega_C)) - \deg C \\ &= v \deg_{C_1}(\omega_C) + v \deg_{C_2}(\omega_C) - \deg C_1 - \deg C_2, \end{aligned}$$

we obtain iii):  $\frac{w}{2} \cong \left| v \deg_{C_1}(\omega_C) - \deg C_1 \right|.$

Now suppose  $C$  has a smooth rational component  $C_1$  meeting the rest of the curve in  $w$  points  $P_1, \dots, P_w$ . Then  $\omega_C|_{C_1}$  is just the sheaf of differentials on  $C_1$  with poles at  $P_1, \dots, P_w$ , so if  $w \leq 2$ ,  $\deg_{C_1}(\omega_C) \leq 0$ . Using iii) this shows  $\deg C_1 \leq \frac{1}{2}$  if  $w = 1$ , absurd, and  $\deg C_1 \leq 1$  if  $w = 2$ .

Moreover, if, in this last case one of the  $P_1$  lies on a smooth rational curve  $C_2$  meeting the rest of  $C$  in only 1 other point, as in the diagram below



then  $\omega_C|_{C_1} \cong \mathcal{O}_{C_1}$  and  $\omega_C|_{C_2} \cong \mathcal{O}_{C_2}$  so  $\deg_{C_1 \cup C_2}(\omega_C) = 0$ . Using iii) again, we find  $\deg(C_1 \cup C_2) \leq \frac{1}{2} \cdot 2 = 1$ , and as this is absurd, we have proved all parts of the Proposition.

We are now ready to show that  $\mathcal{D} = \mathcal{C}$ . Since  $D_0$  is moduli semi-stable, it follows that  $\mathcal{D}$  is a normal two-dimensional scheme with only type  $A_n$  singularities. Moreover  $\omega_{\mathcal{D}}^{\otimes n}/k[[t]]$  is generated by its sections if  $n \geq 3$  and defines a morphism from  $\mathcal{D}$  to a scheme  $\mathcal{D}'/k[[t]]$ , where  $D'_\eta = D_\eta$ ,  $D'_0 = D_0$  with rational chains blown down to points. Thus  $\mathcal{D}'$  is a family of moduli-stable curves over  $k[[t]]$  with generic fibre  $\mathcal{C}_\eta$ . Since there is only one such (cf. [6]), it follows that  $\mathcal{D}' = \mathcal{C}$ . Thus we have a diagram:



$$\begin{array}{ccccc}
 C_\eta & \xleftarrow{\approx} & D_\eta & \xrightarrow{\Phi_\eta} & \mathbf{P}^N \times \text{Spec } k((t)) \\
 \cap & & \cap & & \cap \\
 \mathcal{C} & \xleftarrow{\quad} & \mathcal{D} & \xrightarrow{\phi} & \mathbf{P}^N \times \text{Spec } k[[t]] \\
 \Phi_\eta^*(\mathcal{O}_{\mathbf{P}^N}(1)) & = & \omega_{D_\eta/k((t))}^{\otimes n} & & 
 \end{array}$$

Let  $L = \mathcal{O}_{\mathcal{D}}(1)$ . It follows that  $L \cong \omega_{\mathcal{D}/k[[t]]}^{\otimes n} (-\sum r_i D_i)$ , where  $D_i$  are the components of  $D_0$ . Multiplying the isomorphism by  $t^{\min(r_i)}$ , we can assume  $r_i \geq 0$ ,  $\min r_i = 0$ . Let  $D_1 = \bigcup_{r_i=0} D_i$ ,  $D_2 = \bigcup_{r_i>0} D_i$ . If  $f$  is a local equation of  $\sum r_i D_i$ , then  $f \not\equiv 0$  in any component of  $D_1$  since  $r_i = 0$  on all these while  $f(x) = 0$ , all  $x \in D_1 \cap D_2$ , so

$$\#(D_1 \cap D_2) \leq \deg_{D_1}(\mathcal{O}_{\mathcal{D}_0}(\sum r_i D_i)).$$

But this last degree equals  $(\deg D_1 - n \deg_{D_1}(\omega_{D_0}))$  which contradicts iii) of Proposition 5.5 unless all  $r_i$  are zero. Hence  $L = \omega_{\mathcal{C}}^{\otimes n}$  which shows  $\mathcal{D} = \mathcal{C}$ .

### LINE BUNDLES ON THE MODULI SPACE

For the remainder of this section we examine  $\text{Pic}(\overline{\mathcal{M}}_g)$ . We fix a genus  $g \geq 2$  and an  $e \geq 3$ . Then for all stable  $C$ ,  $\omega_C^{\otimes e}$  is very ample and in this embedding  $C$  has degree  $d = 2e(g-1)$ , the ambient space has dimension  $v-1$  where  $v = (2e-1)(g-1)$  and  $C$  has Hilbert polynomial  $P(X) = dX - (g-1)$ . Let  $H \subset \text{Hilb}_{\mathbf{P}^{v-1}}^P$  be the locally closed smooth subscheme of  $e$ -canonical stable curves  $C$ , let  $C \subset H \times \mathbf{P}^{v-1}$  be the universal curve and let

$$\text{ch} : H \rightarrow \text{Div} = \text{Div}^{d,d} = \left\{ \begin{array}{l} \text{projective space of bihomogeneous forms} \\ \text{of bidegree } (d, d) \text{ in dual coordinates} \\ u, v \text{ (cf. § 1).} \end{array} \right\}$$

be the Chow map. These are related by the diagram

$$\begin{array}{ccccc}
 & & C & & \\
 & & \downarrow \pi & & \\
 \text{Div} & \xleftarrow{\text{ch}} & H & \xrightarrow{\rho} & \overline{\mathcal{M}}_g = H/PGL(v)
 \end{array}$$

If  $\text{Pic}(H, PGL(v))$  is the Picard group of invertible sheaves on  $H$  with  $PGL(v)$ -action, we have a diagram