

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 25 (1979)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FIFTEEN CHARACTERIZATIONS OF RATIONAL DOUBLE POINTS AND SIMPLE CRITICAL POINTS
Autor: Durfee, Alan H.
Kapitel: 2. Rational singularities
DOI: <https://doi.org/10.5169/seals-50375>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 06.02.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

germs V and W embedded in \mathbf{C}^n at the origin are *isomorphic* if there is a germ of an analytic automorphism of \mathbf{C}^n fixing the origin and taking V to W .

Characterization A1. The analytic set $f^{-1}(0)$ is isomorphic to the zero locus of one of the functions listed in column 1 of Table 1.

2. RATIONAL SINGULARITIES

A *resolution* of a germ of a normal surface singularity V as above is a complex analytic manifold M and an analytic map $\pi: M \rightarrow V$ that is surjective and proper (compact fibers) such that its restriction to $M - \pi^{-1}(\mathbf{v})$ is an analytic isomorphism, and $M - \pi^{-1}(\mathbf{v})$ is dense in M . Resolutions exist, and can be computed with a certain amount of effort. The article [Lipman 2] contains a general discussion of resolutions, and [Laufer 1] and [Hirzebruch, Neumann, and Koh, §9] give a detailed method with examples.

Among all resolutions there is a *minimal resolution* $\pi: M \rightarrow V$ that has the following universal mapping property: Given any other resolution $\pi': M' \rightarrow V$, there is a unique map $\rho: M' \rightarrow M$ with $\pi' = \pi \circ \rho$.

The *geometric genus* p of V is the dimension of the complex vector space $H^1(M, \mathcal{O}_M)$, where M is any resolution of V , and \mathcal{O}_M is the sheaf of holomorphic functions on M [Artin; Wagreich 1, §1.4; Brieskorn 2; Laufer 2]. (V is assumed Stein.) This number is finite, and independent of the choice of resolution. It may alternately be defined as the dimension of the stalk at the origin of the sheaf $R^1 \pi_* \mathcal{O}_M$ on V . The idea behind this definition is that M is a collection of “thickened” curves, and that the genus of a curve X is the dimension of $H^1(X, \mathcal{O}_X)$. For example, $H^1(M, \mathcal{O}_M) = 0$ if M is the total space of a line bundle over a curve of genus zero. On the other hand, $\dim H^1(M, \mathcal{O}_M) = k(k-1)(k-2)/6$ if M is a line bundle of Chern class $-k$ over a curve of genus $(k-1)(k-2)/2$ (the minimal resolution of $f(x, y, z) = x^k + y^k + z^k$). In terms of V alone, p is the dimension of the space of holomorphic two-forms on $V - \mathbf{v}$ divided by square-integrable forms [Laufer 2, Theorem 3.4]. Another formula for p in terms of topological invariants of the resolution M and the nearby fiber F (see §11) is given in [Laufer 6].

The analytic set V has a *rational* singularity if $p = 0$. A rational singularity embeds in codimension 1 if and only if it is a double point (its local ring is of multiplicity two) [Artin, Corollary 6].

Characterization A2. The singularity of $f^{-1}(0)$ is rational.

Characterizations A1 and A2 will both be shown equivalent to Characterization A3.

3. EXCEPTIONAL SETS

Let V be as above, and let $\pi: M \rightarrow V$ be a resolution of V . The *exceptional set* $E = \pi^{-1}(\mathbf{v})$ is compact, one-dimensional, and connected, and hence is a union of irreducible complex curves E_1, \dots, E_s . It is possible to arrange that the E_i are non-singular, the intersection of E_i and E_j is transverse for $i \neq j$, and no three E_i meet at a point. Such a resolution is called *good*. If, in addition, the intersection of E_i and E_j is empty or one point, the resolution is *very good*; this is possible to arrange as well.

Suppose that the resolution is good. Let $E_i \cdot E_j$ equal the number of points of intersection of E_i and E_j if $i \neq j$ (always a non-negative integer), or the first Chern class of the normal bundle to E_i evaluated on the orientation class of E_i if $i = j$ (the self-intersection of E_i). The matrix $\{E_i \cdot E_j\}$ is called the *intersection matrix of the resolution*. It is proved in [Du Val 2] (see also [Mumford; Laufer 1, p. 49]) that this matrix is negative definite. Conversely, given a collection of curves $E = E_1 \cup \dots \cup E_s$ in a two-dimensional manifold M with negative definite intersection matrix $\{E_i \cdot E_j\}$, a theorem of Grauert says that the quotient space M/E has a normal complex structure and that the projection map $M \rightarrow M/E$ is analytic [Laufer 1, p. 60].

Characterization A3. The minimal resolution of $f^{-1}(0)$ is very good, and its exceptional set consists of curves of genus 0 and self-intersection -2 .

The equivalence of Characterizations A2 and A3 is proved in [Du Val 1], and [Artin]. The following facts are needed:

- (i) Let $M \rightarrow V$ be a resolution of a normal singularity V as above. There is a certain unique non-zero divisor $Z = \sum n_i E_i$ on M with $n_i \geq 0$ called the *fundamental cycle*, and it is shown that the singularity of V is rational if and only if the analytic Euler characteristic $\chi(Z)$ of Z is 1 (that is, the arithmetic genus of Z is 0) [Artin, Theorem 3]. It is easy to see that the support of Z is the whole exceptional set of E .
- (ii) Any resolution of a rational singularity V is very good, and the curves in the exceptional set are of genus zero [Brieskorn 2, Lemma 1.3].