

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 28 (1982)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE REPRESENTATION THEORY OF  $SL(2, \mathbb{R})$ , A NON-INFINITESIMAL APPROACH  
**Autor:** Koornwinder, Tom H.  
**Kapitel:** 5.1. Spherical functions  
**DOI:** <https://doi.org/10.5169/seals-52233>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

**Download PDF:** 18.04.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

5. EQUIVALENCE OF IRREDUCIBLE REPRESENTATIONS OF  $SU(1, 1)$   
TO SUBREPRESENTATIONS OF THE PRINCIPAL SERIES

The first two subsections review some generalities about Gelfand pairs and spherical functions. By using the concepts developed there we can next, in §5.4, translate the problem of classifying the irreducible representations of  $SU(1, 1)$  in such a way that the problem can be solved by global methods. For this the generalized Abel transform (§5.3) and the Chebyshev transform pair of Deans (Theorem 5.10) are the main tools. The problem is finally reduced to finding the continuous characters on the convolution algebra  $\mathcal{D}_{\text{even}}(\mathbf{R})$  (Prop. 5.7).

### 5.1. SPHERICAL FUNCTIONS

We remember some of the standard facts about spherical functions (cf. for instance GODEMENT [20], HELGASON [25, Ch. X], FARAUT [12, Ch. 1]). Let  $G$  be a unimodular lcsc. group with compact subgroup  $K$ .  $(G, K)$  is called a *Gelfand pair* if  $C_c(K \backslash G / K)$  is a commutative algebra under convolution. If there is a continuous involutive automorphism  $\alpha$  on  $G$  such that  $\alpha(KxK) = Kx^{-1}K(x \in G)$  then  $(G, K)$  is a Gelfand pair. If  $(G, K)$  is a Gelfand pair and the irreducible representation  $\tau$  of  $G$  is unitary or  $K$ -finite then the representation  $1$  of  $K$  has multiplicity 0 or 1 in  $\tau$ .

Let  $(G, K)$  be a Gelfand pair. A *spherical function* is a function  $\phi \neq 0$  on  $G$  such that

$$\phi(x)\phi(y) = \int_K \phi(xky)dk, \quad x, y \in G.$$

The nonzero continuous algebra homomorphisms from  $C_c(K \backslash G / K)$  (or  $C_c^\times(K \backslash G / K)$  if  $G$  is a Lie group) to  $\mathbf{C}$  are precisely of the form

$$(5.1) \quad f \rightarrow \int_G f(x)\phi(x^{-1})dx,$$

where  $\phi$  is a spherical function. If  $\tau$  is a  $K$ -unitary representation of  $G$  and if  $\mathcal{H}(\tau)$  contains a  $K$ -fixed unit vector  $v$ , unique up to a constant factor, then  $x \mapsto (\tau(x)v, v)$  is a spherical function.