

EXACT SEQUENCES OF WITT GROUPS OF EQUIVARIANT FORMS

Autor(en): **Lewis, D. W.**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-52972>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

EXACT SEQUENCES OF WITT GROUPS OF EQUIVARIANT FORMS

by D. W. LEWIS

We construct two exact octagons i.e. circular eight-term exact sequences of Witt groups of forms invariant under the action of a finite group. When the group is trivial our octagons reduce to the two exact sequences obtained in [3]. See also [4].

We are indebted to Cl. Cibils and M. Kervaire for their suggestions to improve the original version of this paper.

Let F be a skewfield, J an involution on F i.e. an anti-automorphism of period two. We allow the case of J being the identity if F is commutative. Let π be a finite group.

Definition. A form over (F, π, J) is a map $\phi : V \times V \rightarrow F$, V an $F\pi$ -module finite dimensional over F , which is sesquilinear, hermitian symmetric with respect to J , and π -invariant in that $\phi(gx, gy) = \phi(x, y)$ for all $g \in \pi$, all $x, y \in V$. Our forms are assumed to be non-singular i.e. $V \rightarrow V^*$, $x \rightarrow \phi(x, -)$ is bijective for all $x \in V$, where V^* is the F -dual of V . We write $W(F, \pi, J)$ for the Witt group of non-singular forms over (F, π, J) , our definition of Witt group being as in [1]. (Remark—the forms which have Witt class zero are precisely those which are neutral i.e. which contain a submodule equal to its orthogonal complement. Note that we do not insist that this submodule be a direct summand as is required in another definition of Witt group which occurs in the literature. When $\text{char } F$ does not divide $|\pi|$ then there is of course no difference between the two definitions of Witt group but in general they will be different.)

Now let K be a field, $\text{char } K \neq 2$, and let L be a quadratic extension of K so that $L = K(i)$, $i^2 = a$ for some $a \in K$. L admits both the identity map and the map—given by $\bar{i} = -i$ as involutions. We will consider the groups $W(K, \pi, 1)$, $W(L, \pi, 1)$ and $W(L, \pi, -)$. Also we write $W_{-1}(K, \pi, 1)$, $W_{-1}(L, \pi, 1)$ for the Witt groups of non-singular forms ϕ defined as above except that now ϕ is required to be skew-symmetric i.e. $\phi(y, x) = -\phi(x, y)$ for all $x, y \in V$. Also we write $W_{-1}(L, \pi, -)$ for the Witt group of skew-hermitian forms over L , i.e. $\phi(y, x) = -\phi(x, y)$ for all $x, y \in V$. Note that for $\pi = 1$, the groups

$W_{-1}(K, \pi, 1)$, $W_{-1}(L, \pi, 1)$ are trivial since the skew-symmetric forms are even-dimensional and classified by rank alone [2, p. 334]. Note also that $W_{-1}(L, \pi, -)$ is isomorphic to $W(L, \pi, -)$ because if ϕ is hermitian then $i\phi$ is skew-hermitian and vice versa.

Let the trace maps $T_\alpha : L \rightarrow K$, $\alpha = 1, 2$ be defined by

$$T_\alpha(r_1 + r_2i) = r_\alpha, \alpha = 1, 2$$

where each $r_\alpha \in K$. These trace maps induce in an obvious way maps between Witt groups as follows:

$$\begin{aligned} W(L, \pi, -) &\xrightarrow{T_1} W(K, \pi, 1), \\ W(L, \pi, 1) &\xrightarrow{T_2} W(K, \pi, 1), \\ W_{-1}(L, \pi, -) &\xrightarrow{T'_1} W_{-1}(K, \pi, 1), \\ W_{-1}(L, \pi, 1) &\xrightarrow{T'_2} W_{-1}(K, \pi, 1). \end{aligned}$$

We denote the last two maps by T'_1, T'_2 merely to distinguish them from the first two maps.

Also we may use the tensor product in a natural way to define maps

$$\begin{aligned} U_1 : W(K, \pi, 1) &\rightarrow W(L, \pi, 1) \\ U'_1 : W_{-1}(K, \pi, 1) &\rightarrow W_{-1}(L, \pi, 1) \end{aligned}$$

and there are also maps

$$\begin{aligned} U_2 : W(K, \pi, 1) &\rightarrow W_{-1}(L, \pi, -) \\ U'_2 : W_{-1}(K, \pi, 1) &\rightarrow W(L, \pi, -) \end{aligned}$$

given by tensor product together with multiplication by the element $i \in L$. E.g. given a form $\phi : V \times V \rightarrow K$ over $(K, \pi, 1)$, $U_2(\phi)$ is the map


$$V \otimes_K L \times V \otimes_K L \rightarrow L$$

given by

$$(U_2(\phi))(x \otimes \lambda, y \otimes \mu) = \bar{\lambda}i\phi(x, y)\mu$$

for all $x, y \in V$, all $\lambda, \mu \in L$. It is easily checked that all these maps are well-defined.

THEOREM 1. *There is an exact octagon of Witt groups*



Proof. We first show exactness of the portion

$$W(L, \pi, -) \xrightarrow{T_1} W(K, \pi, 1) \xrightarrow{U_1} W(L, \pi, 1),$$

i.e. we show that image of T_1 is the kernel of U_1 .

Let $\phi : V \times V \rightarrow L$ represent an element of $W(L, \pi, -)$. To see that $U_1 T_1 \phi$ is neutral as a form over $(L, \pi, 1)$ we consider the subspace W of $V \otimes_K L$ as defined by

$$W = \{iv \otimes 1 + v \otimes i : v \in V\}.$$

Clearly W is an $L\pi$ -submodule and $2 \dim_K W = \dim_K(V \otimes_K L)$. We will show that $W = W^\perp$, orthogonal complement with respect to $U_1 T_1 \phi$. Now if $v, v' \in V$ then

$$(U_1 T_1 \phi)(iv \otimes 1 + v \otimes i, iv' \otimes 1 + v' \otimes i)$$

is easily verified to be zero using the sesquilinearity of ϕ and the definitions of T_1, U_1 . Thus $W \subset W^\perp$. It follows that in fact $W = W^\perp$ since they have the same dimension.

Next let $\psi : V \times V \rightarrow K$ represent an element of $W(K, \pi, 1)$. We may assume ψ is anisotropic by [1]. Now if $U_1 \psi$ is zero in $W(L, \pi, 1)$ then $V \otimes_K L$ contains a self-orthogonal L -submodule W . This enables us to define an L -space structure on V as follows:

Observe that

$$2 \dim_L W = \dim_L V \otimes_K L, \dim_L W = \dim_L V \otimes i,$$

and that $W \cap (V \otimes i) = 0$ since ψ is anisotropic. Thus $V \otimes_K L \cong (V \otimes i) \oplus W$. It now follows that, given $v \in V$, there exists a unique element $v' \in V$ such that $v \otimes 1 + v' \otimes i \in W$. Then define the operator $J: V \rightarrow V$ by $J(v') = v$ for each $v \in V$. It is easily verified that J is skew-adjoint, $J^2 = a$ and that J commutes with the π -action. Thus J can be used to give V an $L\pi$ -module structure, $i \in L$ operating as J on V .

Now define a form $\phi: V \times V \rightarrow L$ by

$$\phi(x, y) = \psi(x, y) + i^{-1} \psi(x, Jy)$$

for all $x, y \in V$. Then ϕ is a non-singular form over $(L, \pi, -)$ and $T_1\phi = \psi$.

This proves exactness at $W(K, \pi, 1)$. At the three points in the sequence

$$W(L, \pi, 1) \xrightarrow{T_2} W(K, \pi, 1) \xrightarrow{U_2} W_{-1}(L, \pi, -),$$

$$W_{-1}(L, \pi, -) \xrightarrow{T'_1} W_{-1}(K, \pi, 1) \xrightarrow{U'_1} W_{-1}(L, \pi, 1),$$

$$W_{-1}(L, \pi, 1) \xrightarrow{T'_2} W_{-1}(K, \pi, 1) \xrightarrow{U'_2} W(L, \pi, -)$$

exactness is proven by the same arguments.

Now consider the piece

$$W_{-1}(K, \pi, 1) \xrightarrow{U'_1} W_{-1}(L, \pi, 1) \xrightarrow{T'_2} W_{-1}(K, \pi, 1).$$

If $\phi: V \times V \rightarrow K$ represents an element of $W_{-1}(K, \pi, 1)$ then we see that $T'_2 U'_2 \phi$ is neutral by looking at

$$W \subset V \otimes_K L, W = V \otimes 1$$

and checking that $W = W^\perp$.

$$T'_2 U'_2 \phi(v_1 \otimes 1, v' \otimes 1) = T'_2 \phi(v, v') = 0$$

for all $v, v' \in V$ so that $W \subset W^\perp$. Hence $W = W^\perp$ since

$$2 \dim_K W = \dim_K V \otimes_K L.$$

Conversely if ψ , representing an element of $W_{-1}(L, \pi, 1)$, is such that $T'_2 \psi$ is neutral then $\psi: V \times V \rightarrow L$, V an $L\pi$ -module, and there exists a $K\pi$ -module W of V with $W = W^\perp$, orthogonal complement with respect to $T'_2 \psi$. Also

$2 \dim_K W = \dim_K V$. Defining $\phi : W \times W \rightarrow V$ by $\phi(x, y) = \psi(x, y)$ for $x, y \in W$ then $W \otimes_K L \cong V$ as $L\pi$ -modules via the isomorphism

$$w \otimes \lambda \rightarrow \lambda w, \lambda \in L, w \in W.$$

Moreover $U'_1(\phi) = \psi$ completing the proof of exactness at $W_{-1}(L, \pi, 1)$.

For the three remaining points of the sequence, which each have U followed by T , the above arguments go through virtually unchanged.

This completes the proof.

Now suppose we have a quaternion division algebra D over K , $D = \left(\frac{a, b}{K} \right)$ generated by i, j with $i^2 = a, j^2 = b, ij = -ji$ etc. We have involutions $-$ and $\hat{}$ on D given by $\bar{i} = -i, \bar{j} = -j$ and $\hat{i} = i, \hat{j} = j$ respectively. Let L be the maximal subfield $K(i)$ of D . There are trace maps $T_i : D \rightarrow L, i = 1, 2$ given by $T_i(z_1 + z_2 j) = z_1$ where $z_1, z_2 \in L$, and these induce natural maps of Witt groups

$$W(D, \pi, -) \xrightarrow{T_1} W(L, \pi, -),$$

$$W(D, \pi, \hat{}) \xrightarrow{T_2} W(L, \pi, 1),$$

$$W(D, \pi, \hat{}) \xrightarrow{T'_1} W(L, \pi, -),$$

$$W(D, \pi, -) \xrightarrow{T'_2} W_{-1}(L, \pi, 1).$$

Also we have maps

$$W(L, \pi, -) \xrightarrow{U_1} W(D, \pi, \hat{}),$$

$$W(L, \pi, 1) \xrightarrow{U_2} W(D, \pi, \hat{}),$$

$$W(L, \pi, -) \xrightarrow{U'_1} W(D, \pi, -),$$

$$W_{-1}(L, \pi, 1) \xrightarrow{U'_2} W(D, \pi, -),$$

U_1, U'_1 given by the tensor product, U_2, U'_2 by the tensor product together with multiplication by the element $k = ij$ of D . E.g. given a form $\phi : V \times V \rightarrow L$ over $(L, \pi, 1)$, $U_2(\phi)$ is the form $V \otimes_L D \times V \otimes_L D \rightarrow D$ defined by

$$U_2(\phi)(x \otimes \lambda, y \otimes \mu) = \hat{\lambda} \phi(x, y) k \mu \quad \text{for } \lambda, \mu \in D, x, y \in V.$$

(Beware that the position of k matters as D is not commutative!).

THEOREM 2. *There is an exact octagon of Witt groups*

$$\begin{array}{ccc}
 W(D, \pi, -) & \xrightarrow{T_1} & W(L, \pi, -) \\
 \nearrow U'_2 & & \searrow U_1 \\
 W_{-1}(L, \pi, 1) & & W(D, \pi, \wedge) \\
 \uparrow T'_2 & & \downarrow T_2 \\
 W(D, \pi, -) & & W(L, \pi, 1) \\
 \nwarrow U'_1 & & \swarrow U_2 \\
 W(L, \pi, -) & \xleftarrow{T'_1} & W(D, \pi, \wedge)
 \end{array}$$

Proof. We need only modify the proof of theorem 1 slightly. Specifically j will play the role that i did in theorem 1. For example at the start of the proof we must put

$$W = \{jv \otimes 1 + v \otimes j : v \in V\}$$

and later on the operator J is defined in a similar fashion to that of theorem 1 except that we get $J^2 = b$ leading to a $D\pi$ -module structure. The lack of commutativity of D causes no problem, although care must be taken in dealing with the maps U_2, U'_2 . (See the comment above.) We leave the reader to check that with these modifications the proof goes through completely.

Comment 1. When $\pi = 1$ the Witt groups $W_{-1}(K, \pi, 1)$ and $W_{-1}(L, \pi, 1)$ are trivial as we remarked earlier in this paper. Our sequences now reduce to those of [3].

Comment 2. Note that $W_{-1}(L, \pi, -) \cong W(L, \pi, -)$ for the reason stated earlier.

Also $W(D, \pi, \wedge) \cong W_{-1}(D, \pi, -)$ since forms hermitian with respect to \wedge are equivalent to those skew-hermitian with respect to $-$ and vice versa. (The correspondence $\phi \leftrightarrow i\phi$ gives this since $\hat{x} = i^{-1}\bar{x}i$ for all $x \in D$.) A consequence of the above is that the two octagons each display an interesting symmetry

feature. In the “antipodal” position to $W(F, \pi, J)$ in the octagon we always have $W_{-1}(F, \pi, J)$.

Comment 3. Our proof is different from that of [3] and it may well be possible that this new method of proof can also be used to generalize the sequences of [3] to the case when K is a commutative ring and L is some kind of Galois extension with Galois group cyclic of order two.

REFERENCES

- [1] CIBILS, Claude. Groupe de Witt d'une algèbre avec involution. *L'Enseignement Mathématique* 29 (1983), 27-43.
- [2] JACOBSON, N. *Basic Algebra*. W. H. Freeman, San Francisco, 1974.
- [3] LEWIS, D. W. New improved exact sequences of Witt groups. *J. of Algebra* 74 (1982), 206-210.
- [4] ——— A note on hermitian and quadratic forms. *Bull. London Math. Soc.* 11 (1979), 265-267.

(Reçu le 16 juillet 1982)

D. W. Lewis

Department of Mathematics
University College
Belfield
Dublin 4
Ireland

Vide-leer-empty