

3. Digression on hyperbolic geometry

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

proof requires Tits' theorem is due to Gromov: a finitely generated group has polynomial growth if and only if it is almost nilpotent [G].

The analogue of Tits' theorem for division rings does not hold as such [L1], but conjectural statements have been formulated [L2]. Another generalisation of the theorem is proposed as a research problem in remark 1.4.2 of [BL].

3. DIGRESSION ON HYPERBOLIC GEOMETRY

Let n be an integer, $n \geq 1$. The hyperbolic space H^{n+1} of dimension $n + 1$ is the open unit ball of the euclidean space \mathbf{R}^{n+1} . Hyperbolic lines (called lines below) in H^{n+1} are traces on H^{n+1} of circles and euclidean lines in \mathbf{R}^{n+1} which are orthogonal to \mathbf{S}^n . Two distinct points $P, Q \in H^{n+1}$ are on a unique line which determines two points $P_\infty, Q_\infty \in \mathbf{S}^n$, say with P, Q, Q_∞, P_∞ arranged in cyclic order on the euclidean circle defining this line. The (hyperbolic) distance between P and Q is given by a cross-ratio of euclidean distances; more precisely, it is defined to be

$$d(P, Q) = \text{Log}(P, Q, Q_\infty, P_\infty) = \log \left(\frac{|P - Q_\infty|}{|P - P_\infty|} : \frac{|Q - Q_\infty|}{|Q - P_\infty|} \right).$$

The *proper Mæbius group* $GM(n)_0$ is the group of orientation preserving isometries of \mathbf{R}^{n+1} for this distance. Any $g \in GM(n)_0$ extends to a homeomorphism of the closed ball $H^{n+1} \cup \mathbf{S}^n$. One may check that $GM(1)_0$ is isomorphic to $PGL(2, \mathbf{R})$ and $GM(2)_0$ to $PGL(2, \mathbf{C})$.

There is an equivalent description with H^{n+1} the half space $\mathbf{R}^n \times \mathbf{R}_+^*$. The set of "points at infinity" is then $\mathbf{R}^n \cup \{\infty\}$ rather than \mathbf{S}^n .

For all this, see e.g. [A] or [Si].

An isometry $g \in GM(n)_0$ is said to be

elliptic if there is some point in H^{n+1} fixed by g ,

parabolic if there is in \mathbf{S}^n exactly one point fixed by g ,

hyperbolic if there is a line in H^{n+1} invariant by g on which g has no fixed point.

(Following Thurston [Th], we call "hyperbolic" elements which are "loxodromic" in classical literature, such as in [Gr].)

PROPOSITION. *Elliptic, parabolic and hyperbolic elements define a partition of the proper Mæbius group in three disjoint classes.*

Proof. Let us first check that the three classes do not overlap in $GM(n)_0$. If g is hyperbolic, it has two fixed points in \mathbf{S}^n and thus cannot be parabolic; if g was

also elliptic, the foot of the perpendicular from the fixed point of g onto the invariant line of g would be fixed by g , and this cannot be. If g was at the same time elliptic with fixed point $a \in H^{n+1}$ and parabolic with fixed point $b \in S^n$, the line from a towards b would have two points at infinity b and b' both fixed by g , and this cannot be.

That any $g \in GM(n)_0$ belongs to one of the three classes follows for example from Brouwer's fixed point theorem. (See also 4.9.3 in [Th].) \square

Observe that an hyperbolic isometry $g \in GM(n)_0$ has a unique invariant line δ . Suppose indeed that there are two of them, say δ and δ' . If $\delta \cap \delta' \neq \emptyset$, the intersection point (which is unique) is fixed by g , and this cannot be. If $\delta \cap \delta' = \emptyset$ and if δ, δ' have no common point at infinity, there is a unique line perpendicular to both δ and δ' ; but this line intersects δ in a point fixed by g , and this cannot be. Assume finally that $\delta \cap \delta' = \emptyset$ and that δ and δ' have a common point at infinity; choose some number $\rho > 0$ and consider the set C_ρ of points in H^{n+1} at a distance of ρ from δ' ; the intersection $C_\rho \cap \delta$ is a point fixed by g , and again this cannot be. One may consequently also define an isometry $g \in GM(n)_0$ to be

elliptic if $d(a, g(a)) = 0$ for some $a \in H^{n+1}$,

parabolic if $\inf d(a, g(a)) = 0$, with the infimum over $a \in H^{n+1}$ not attained,

hyperbolic if $\inf d(a, g(a)) > 0$ (and the infimum is then attained exactly on the invariant line of g).

We shall need below the following *dynamical description*. An hyperbolic isometry $g \in GM(n)_0$ has on S^n one attracting point P_a and one repulsing point P_r . This means that, for any neighborhood U of P_a in S^n and for any compact subset K of $S^n - \{P_r\}$, one has $g^k(K) \subset U$ for k large enough. (And similarly with g^{-1} instead of g when exchanging P_a and P_r .) Consider now a parabolic isometry $g \in GM(n)_0$ with fixed point $P \in S^n$. Let U be a neighborhood of P in S^n and let K be compact in $S^n - \{P\}$; then $g^k(K) \subset U$ for any $k \in \mathbf{Z}$ with $|k|$ large enough. (This is obvious when g is a translation in $\mathbf{R}^n \times \mathbf{R}_+^*$ by some vector in \mathbf{R}^n , and any parabolic isometry of H^{n+1} is conjugate to such a translation.)

4. FREE SUBGROUPS OF $GL(2, \mathbf{R})$ AND OF $GL(2, \mathbf{C})$

We show in this section that a subgroup of the proper Mœbius group $G = PGL(2, \mathbf{R})$ which is not almost solvable contains free groups; the same fact for $GL(2, \mathbf{R})$ follows straightforwardly. We discuss also the case of $GL(2, \mathbf{C})$.