

§5. Euclidean Spaces

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **30 (1984)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$f_w(SO_{n+1}^m)$ is not contained in a proper algebraic subset (in this case, A) of SO_{n+1} . This completes the proof of Theorem 1 (a) for S^n .

Next, consider Theorem 1 (c) for S^n . First observe that this can be proved for SO_3 by the technique above, if A is taken to consist simply of the identity. This is because the action of SO_3 on S^2 is locally commutative, so all that is needed is a perfect set of free generators, which in turn requires only that each R_w be nowhere dense. Theorem 1 of [5] again applies, because A is an algebraic set: membership in A is equivalent to the simultaneous vanishing of $(n+1)^2$ polynomials which, by using a sum of squares, is equivalent to the vanishing of a single polynomial. For higher dimensions, we appeal to the technique used by Borel to get locally commutative free subgroups of SO_{n+1} . In [5, p. 162] he showed that, if $n \geq 2$, SO_3 may be represented as a subgroup H of SO_{n+1} where H 's action on S^n is locally commutative. Hence the perfect free generating set in SO_3 yields a perfect subset of H which is the desired free generating set in SO_{n+1} .

§ 5. EUCLIDEAN SPACES

For the Euclidean case of Theorem 1, it suffices to consider \mathbf{R}^3 , since any isometry of \mathbf{R}^3 can be extended to one in higher dimensions by simply fixing the additional coordinates; this introduces no new fixed points. Now, \mathbf{R}^3 can be handled in a way entirely similar to S^n . Any orientation-preserving isometry of \mathbf{R}^3 is a screw-motion, i.e. a rotation $\rho \in SO_3$ followed by a translation τ . Such isometries may be represented as elements of $SL_4(\mathbf{R})$ as follows: if $\sigma = \tau\rho$ where ρ corresponds to $(a_{ij}) \in SO_3$ and τ is a translation by (v_1, v_2, v_3) , then identify σ with the matrix

$$\begin{pmatrix} & & & v_1 \\ & & & v_2 \\ & a_{ij} & & v_3 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Since composition of isometries corresponds to matrix multiplication, this shows that $G(\mathbf{R}^3)$ may be viewed as a connected (6-dimensional) analytic submanifold of \mathbf{R}^{12} . Now, the proof can proceed exactly as for spheres, once it is shown that the existence of a fixed point is equivalent to the

vanishing of a polynomial. But a screw-motion σ has a fixed point if and only if the translation vector is perpendicular to the axis of the rotation. Since the axis of a rotation $(a_{ij}) \in SO_3$ is parallel to $(a_{32} - a_{23}, a_{13} - a_{31}, a_{21} - a_{12})$, σ has a fixed point if and only if $v_1(a_{32} - a_{23}) + v_2(a_{13} - a_{31}) + v_3(a_{21} - a_{12}) = 0$. This completes the proof of Theorem 1 (a) for \mathbf{R}^n .

§ 6. HYPERBOLIC SPACES

Here we meet a case where the existence of a free, fixed-point free group of isometries having rank 2 does not imply the existence of such a group having uncountable rank. The hyperbolic plane is such a space.

If H^2 is identified with the upper half-plane of \mathbf{C} , then $G(H^2)$ corresponds to linear fractional transformations $z \mapsto \frac{az + b}{cz + d}$, where a, b, c, d are real and $ad - bc \neq 0$. Since it may be assumed that $ad - bc = 1$, this group is isomorphic to $PSL_2(\mathbf{R})$. A nonidentity element of $PSL_2(\mathbf{R})$ is called elliptic, parabolic, or hyperbolic according as the absolute value of its trace is less than, equal to, or greater than two; the nonidentity elements of $G(H^2)$ with a fixed point in H^2 correspond to the elliptic elements of $PSL_2(\mathbf{R})$. See [18] for more details about this interpretation of $PSL_2(\mathbf{R})$. The following theorem clarifies the situation regarding fixed-point free subgroups of $G(H^2)$.

THEOREM 3. (Siegel) *If F is a free subgroup of $PSL_2(\mathbf{R})$ then F is discrete if and only if F has no elliptic elements.*

Theorem 3 is a rephrasing of the result of [34] (see also [15]). An elementary proof appears in [41]. The forward direction is an immediate consequence of the fact that the nondiscrete cyclic subgroups of $PSL_2(\mathbf{R})$ are precisely the ones generated by an elliptic element of infinite order. This fact also yields the reverse direction in the case when F is cyclic. Siegel gave an algebraic proof of the reverse direction for noncyclic free groups. This can also be obtained by first using techniques of Lie algebras to show that a nondiscrete, nonsolvable subgroup of $PSL_2(\mathbf{R})$ is dense in $PSL_2(\mathbf{R})$, and observing that the elliptics form an open set; this approach is due, independently, to A. Borel and D. Sullivan.

The forward (easy) direction of Theorem 3 yields a proof of the positive part of Theorem 1 (b) for H^2 (and hence for H^n , $n \geq 2$), since it implies that a discrete free group of rank two has no elliptic elements. Therefore