

Zeitschrift: L'Enseignement Mathématique
Band: 34 (1988)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ABOUT THE PROOFS OF CALABI'S CONJECTURES ON COMPACT
KÄHLER MANIFOLDS
Kapitel: 5. A PRIORI ESTIMATES: THE ORIGINAL WAY
Autor: Delanoë, Ph. / Hirschowitz, A.
DOI: <https://doi.org/10.5169/seals-56591>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 07.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. A PRIORI ESTIMATES: THE ORIGINAL WAY

According to proposition 4.3 we must prove now that, given any sequence of positive real numbers $(K_i), i \in \mathbf{N}$, there exists a sequence (C_i) such that

$$\forall i \in \mathbf{N}, \quad \| D^i P_\lambda(\varphi) \| \leq K_i$$

implies

$$\| \varphi \| \leq C_0, \quad \forall i \in \mathbf{N}, \quad \| D^i \nabla \bar{\nabla} \varphi \| \leq C_{i+2}.$$

These are *a priori* estimates of order zero, two, three, and so on ... In case $\lambda > 0$, the C^0 estimate is straightforward [2]. In case $\lambda = 0$, it becomes very tricky; proofs simpler than Yau's original one [24] (p. 352-359), based on the idea of uniformly estimating the $L^p(dX_g)$ norms of φ , may be found in [16] (dimension 2), [3] [21] and [4] (p. 148-149).

Estimates of order two and three are carried out by means of tensor calculus and of the Maximum Principle (for elliptic equations) [20] applied to *suitable* test functions. Though it is not everywhere clear in [21] [24], it is worth noting that the computations can be written intrinsically, i.e. without any reference to a *particular* system of coordinates (e.g. [2]), or even *coordinate free* (see section 6 below).

Further regularity is then recovered by Schauder theory e.g. [5] (lemma 1). In the sequel, we show how further estimates can be carried out instead, *just going ahead with coordinate free tensor calculus*. This occurs actually for any fully nonlinear second order elliptic equation on a compact Riemannian manifold, via a straightforward imitation of the device below.

Remark 5.1. It follows from the C^2 *a priori* estimates that the metrics g' are *a priori* uniformly equivalent to the original metric g (see e.g. [3], p. 75).

6. COORDINATE FREE TENSOR CALCULUS

Even coordinate free tensor calculus needs indices. Usually these indices refer to a *local* frame. Another way is to view these indices *globally* as labelling copies of the holomorphic and antiholomorphic tangent and cotangent bundles. From this point of view, a tensor written with indices is a section of the tensor product of a family of bundles indexed by an *unordered* set of indices (disregarding those indices subject to the summation convention).