

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 35 (1989)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE CANTOR SET AND A GEOMETRIC CONSTRUCTION
Autor: Pavone, Marco
Kapitel: Introduction
DOI: <https://doi.org/10.5169/seals-57361>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE CANTOR SET AND A GEOMETRIC CONSTRUCTION

by Marco PAVONE

INTRODUCTION

The Cantor ternary set consists of all those real numbers x in $[0, 1]$ which have a ternary expansion $x = \sum_{n=1}^{\infty} a_n/3^n$ for which a_n is never 1. Equivalently, C can be obtained in a purely geometrical fashion by first removing from $[0, 1]$ the middle third $(1/3, 2/3)$, then removing the middle thirds $(1/9, 2/9)$ and $(7/9, 8/9)$ of the remaining intervals, and so on (C will be exactly the complement of the countable union of the removed intervals). If $x = \sum_{n=1}^{\infty} a_n/3^n$ is in C , the geometric interpretation of its ternary expansion is that x is the unique point in $[0, 1]$ which is reached by first staying to the left or to the right of $(1/3, 2/3)$ if $a_1 = 0$ or $a_1 = 2$ respectively, then staying to the left or to the right of the next removed interval if $a_2 = 0$ or $a_2 = 2$ respectively, and so on. It follows from the construction that C is a nowhere dense closed subset of $[0, 1]$.

A well known property of C is that any real number in $[0, 2]$ can be written as the sum of two numbers in C . The purpose of this note is to give an elementary proof of $C + C = [0, 2]$ which only uses the geometric definition of C . A refinement of the proof shows in fact that for any k in $[0, 2]$ there exists either a finite or an uncountable number of pairs x, y from C such that $x + y = k$. We also discuss the analogy between this decomposition result and certain properties of continued fractions.

THE GEOMETRIC CONSTRUCTION

We set, as usual, $C \times C = \{(x, y) \in \mathbf{R}^2 : x, y \in C\}$. Then $C + C = [0, 2]$ can be geometrically restated as

- (*) for any k in $[0, 2]$ the line $x + y = k$ intersects $C \times C$ in at least one point.