

# 1. The groups $G_s$ and their representations

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **35 (1989)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.07.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

computed; they turn out to be isomorphic to  $\pi_s(U)$  and  $\pi_s(O)$  respectively. Moreover a product is defined in the direct sum of the  $E_s^U(E_s^O)$  turning it into a graded ring  $E_*^U(E_*^O)$ . The claim of Theorem A is proved in Section 3; we show that the maps  $\phi: E_s^U \rightarrow \pi_s(U)$ ,  $\psi: E_s^O \rightarrow \pi_s(O)$  given by the  $f_s$  of 0.1 are isomorphisms. Using the product structure in  $\pi_*(U)$  and  $\pi_*(O)$  known from  $K$ -theory the proof reduces to simple verifications in low dimensions. The symplectic case is dealt with in Section 4. In Section 5 we make a remark concerning the "linearization phenomenon" for the homotopy groups of  $U$ ,  $O$  and  $Sp$ .

## 1. THE GROUPS $G_s$ AND THEIR REPRESENTATIONS

1.1. We will denote throughout by  $G_s$  the group given by the presentation

$$G_s = \langle \varepsilon, a_1, \dots, a_s \mid \varepsilon^2 = 1, a_j^2 = \varepsilon, a_j a_k = \varepsilon a_k a_j, j, k = 1, 2, \dots, s, j \neq k \rangle.$$

Clearly any set  $A_1, \dots, A_s$  of HR-matrices yields a (unitary or orthogonal) representation of  $G_s$  of degree  $n$  by  $\varepsilon \mapsto -E$ ,  $a_j \mapsto A_j$ ,  $j = 1, 2, \dots, s$ . Conversely a representation of  $G_s$  with  $\varepsilon \mapsto -E$ , in short an  $\varepsilon$ -representation, yields a set of  $s$  HR-matrices. For the elementary properties of  $G_s$  and its representations we refer to [E]. We just recall that the order of  $G_s$  is  $2^{s+1}$ , that  $\varepsilon$  is central, and that the irreducible unitary  $\varepsilon$ -representations of  $G_s$  are of degree  $2^{s/2}$  if  $s$  is even (one equivalence class), of degree  $2^{(s-1)/2}$  if  $s$  is odd (two equivalence classes). These degrees are the minimal values  $n_0$  in case  $U$ . As for the case  $O$ , one has to recall that a representation is equivalent to an orthogonal one if and only if it is equivalent to a real (and orthogonal) one. Thus, unless an irreducible unitary  $\varepsilon$ -representation is already real, one has to add its conjugate-complex representation, and the discussion of the various cases depending on  $s$  yields the minimal values  $n_0$  (case  $O$ ) mentioned in the introduction; in other words, the degrees of the irreducible orthogonal  $\varepsilon$ -representations of  $G_s$ .

1.2. A very simple and useful scheme for studying the groups  $G_s$  and their  $\varepsilon$ -representations (and many other things) has been devised by T. Y. Lam and T. Smith [LS]. They have expressed the  $G_s$  as products of very small and well-known groups. Namely  $C = G_1$ , the cyclic group of order 4;  $Q = G_2$ , the quaternionic group of order 8;  $K$ , the Klein 4-group; and  $D$ , the dihedral group of order 8. Although  $K$  and  $D$  do not belong to the family  $G_s$ , they are of a similar nature and contain a distinguished central element  $\varepsilon$  of order 2 (distinguished arbitrarily in  $K$ ). "Product" here means the central product obtained from the direct product by identifying the

two  $\varepsilon$ 's. The expression for the  $G_s$  then is as follows, displaying a fundamental periodicity modulo 8:

(2)	$s$	0	1	2	3	4	5	6	7	8	9	...
	$G_s$	$\mathbf{Z}/2$	$C$	$Q$	$QK$	$QD$	$D^2C$	$D^3$	$D^3K$	$D^4$	$D^4C$	...

and  $G_{s+8} = D^4G_s$ .

The tensor product of  $\varepsilon$ -representations of two of the groups  $G_s, K, D$  is an  $\varepsilon$ -representation of their product above, and all  $\varepsilon$ -representations of the  $G_s$  can be obtained in that explicit way from those of  $C, Q, K, D$ , which are well-known. This yields, in particular, the characters  $\chi$  and the Schur indices  $I$  of the irreducible unitary  $\varepsilon$ -representation (the Schur index  $I = 1$  if the representation is equivalent to a real one; if it is not,  $I = -1$  if it is equivalent to the conjugate-complex one,  $I = 0$  otherwise). Both  $\chi$  and  $I$  behave multiplicatively with respect to the central product.

1.3. The Schur indices of the irreducible  $\varepsilon$ -representations are: 0 for  $C = G_1$ ,  $-1$  for  $Q = G_2$ , and 1 for  $K$  and  $D$  (two equivalence classes for  $K$ , one for  $D$ ). This yields the Schur indices  $I_s$  of the irreducible  $\varepsilon$ -representations of the  $G_s$ , as listed in (2) below; we further list the numbers  $v_s^U$  of inequivalent unitary, and  $v_s^O$  of inequivalent orthogonal irreducible  $\varepsilon$ -representations, and the respective degrees  $d_s^U, d_s^O$ . Note that  $I_s$  is periodic with period 8, and  $d_s^O$  is periodic with period 8 in the sense that  $d_{s+8}^O = 16d_s^O$ . Finally we include in the same table the Grothendieck groups  $D_s^U$  and  $D_s^O$  of (equivalence classes of) irreducible  $\varepsilon$ -representations of  $G_s$ , with respect to the direct sum of representations.

(3)	$s$	0	1	2	3	4	5	6	7	8	9	...
	$I_s$	1	0	-1	-1	-1	0	1	1	1	0	...
	$v_s^U$	1	2	1	2	1	2	1	2	1	2	
	$v_s^O$	1	1	1	2	1	1	1	2	1	1	
	$d_s^U$	1	1	2	2	4	4	8	8	16	16	
	$d_s^O$	1	2	4	4	8	8	8	8	16	32	
	$D_s^U$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	
	$D_s^O$	$\mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z} \oplus \mathbf{Z}$	$\mathbf{Z}$	$\mathbf{Z}$	

The values of  $d_s^O$  follow immediately from the  $I_s$  and the  $d_s^U$ . The values  $n_0$  for the case  $O$ , as given in the Introduction, are the  $d_s^O$ .

## 2. THE REDUCED $\varepsilon$ -REPRESENTATION RING

2.1. For all  $s \geq 0$  the group  $G_s$  is the subgroup of  $G_{s+1}$  obtained by omitting the generator  $a_{s+1}$ ; let  $h_s: G_s \rightarrow G_{s+1}$  be the embedding homomorphism. Via  $h_s$  we can restrict an  $\varepsilon$ -representation of  $G_{s+1}$  to  $G_s$ , which in terms of HR-matrices means omitting  $A_{s+1}$ .

Let  $h_s^*: D_{s+1}^U \rightarrow D_s^U$  be the corresponding homomorphism of Grothendieck groups, and  $E_s^U = D_s^U / h_s^* D_{s+1}^U$  the "reduced" groups; similarly  $E_s^O = D_s^O / h_s^* D_{s+1}^O$ . They can easily be computed by means of the characters of  $\varepsilon$ -representations, as follows.

For  $Q$  and  $D$  the character of an irreducible unitary  $\varepsilon$ -representation is 0 except on 1 and  $\varepsilon$ . For  $C$  and  $K$  it is  $\neq 0$  on all 4 elements; on the essential generator ( $\neq \varepsilon$ ) of  $C$  it is  $+i$  or  $-i$  for the two inequivalent representations, and  $+1$  or  $-1$  in the case of  $K$ . For  $G_s$ ,  $s$  even, we infer from the table (2) that the character is 0 except on 1,  $\varepsilon$ . For  $G_s$ ,  $s$  odd, the character is 0 except on 1,  $\varepsilon$  and two further elements  $z, \varepsilon z$ ; on these the two inequivalent  $\varepsilon$ -representations differ just by the sign of the character.

If  $s$  is even,  $d_{s+1}^U = d_s^U = 2^{s/2}$ ; thus the restriction of an irreducible  $\varepsilon$ -representation must be irreducible, whence  $h_s^* D_{s+1}^U = D_s^U$ ,  $E_s^U = 0$ . If  $s$  is odd,  $d_{s+1}^U = 2d_s^U = 2^{(s+1)/2}$ ; thus the restriction is the sum of two irreducible  $\varepsilon$ -representations, and since the character is 0 (except on 1,  $\varepsilon$ ) these two must be inequivalent. Therefore  $h_s^* D_{s+1}^U$  is the "diagonal" of  $D_s^U = \mathbf{Z} \oplus \mathbf{Z}$ , and  $E_s^U = \mathbf{Z}$ ; its generator  $\rho_s$  is represented by either of the two inequivalent irreducible  $\varepsilon$ -representations of  $G_s$ ,  $-\rho_s$  by the other one.

In the orthogonal case the  $E_s^O$  are computed similarly from (3). Since  $d_1^O = 2$  and  $d_0^O = 1$ , the restriction from  $D_1^O$  to  $D_0^O$  yields twice the generator, and  $E_0^O = \mathbf{Z}/2$ ; the same argument holds for  $s \equiv 0 \pmod 8$ ,  $d_{s+1}^O = 2d_s^O$ . Since  $d_2^O = 4$  and  $d_1^O = 2$ , we get  $E_1^O = \mathbf{Z}/2$ . From  $d_3^O = d_2^O = 4$  we get  $E_2^O = 0$ . As for  $s = 3$ , the character argument shows that  $h_3^* D_4^O =$  diagonal of  $D_3^O (= \mathbf{Z} \oplus \mathbf{Z})$ , and  $E_3^O = \mathbf{Z}$ . For  $s = 4, 5, 6$  the dimensions  $d_{s+1}^O = d_s^O$  show that  $E_4^O = E_5^O = E_6^O = 0$ . For  $s = 7$ , the character argument yields  $h_7^* D_8^O =$  diagonal of  $D_7^O (= \mathbf{Z} \oplus \mathbf{Z})$ , and  $E_7^O = \mathbf{Z}$ . Finally one has, for all  $s$ ,  $E_{s+8}^O \cong E_s^O$ .