

Zeitschrift: L'Enseignement Mathématique
Band: 36 (1990)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE INVERSIVE DIFFERENTIAL GEOMETRY OF PLANE CURVES
Kapitel: §7. The canonical map $g:G$
Autor: Cairns, G. / Sharpe, R. W.
DOI: <https://doi.org/10.5169/seals-57907>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 04.10.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The stabilizer of the expression $y = x^4/24 + O(x^6)$ has order 2 and is generated by

$$z \mapsto -\bar{z}.$$

It follows that the non-degeneracy of a vertex is an invariant of inversive geometry.

§7. THE CANONICAL MAP $g: \gamma \rightarrow G$

The considerations of the last section allow us to define a canonical map $g_\gamma: \gamma \rightarrow G$ for vertex free curves γ by mapping a point $p \in \gamma$ to $g_\gamma(p) \in G$, which is the unique group element such that $g_\gamma(p)^{-1}$ sends p to the origin and $g_\gamma(p)^{-1}(\gamma)$ has oriented contact of order 4 with the standard curve $y = x^3/6$ at the origin. We note that if $\gamma' = h(\gamma)$ for some $h \in G$, then obviously $g_{\gamma'}(h(p)) = h(g_\gamma(p))$. Of course altering the initial choice of the origin and the axes used there to describe the model will alter g_γ , but only by right multiplication by some fixed element of G . If $\sigma: (\alpha, \beta) \rightarrow \mathbf{C}$ is a parametrization of the curve by Euclidean arc-length s , and $\sigma'(s) = e^{i\theta(s)}$, then the curvature of the curve at $\sigma(s)$ is $\theta'(s) = \kappa(s)$, and we have the following explicit formula for g .

$$g(s) = \begin{pmatrix} 1 & \sigma \\ 0 & 1 \end{pmatrix} \begin{pmatrix} e^{i\theta/2} & 0 \\ 0 & e^{-i\theta/2} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ (\kappa'' - 2i\kappa\kappa')/4\kappa' & 1 \end{pmatrix} \begin{pmatrix} |\kappa'|^{-1/4} & 0 \\ 0 & |\kappa'|^{1/4} \end{pmatrix}$$

The first two factors are Euclidean motions whose inverse puts γ into oriented first order contact with the oriented x -axis. The rest improve the order of contact to 4 as in §6. It is convenient to regard g as a function of the inverse arc-length v . Now $g(v)$ is a curve on the Lie group G , with tangent vector dg/dv at $g(v)$. Left translation by $g(v)^{-1}$ moves this tangent vector to the origin to yield

$$(7.1) \quad c(v) = g(v)^{-1} \frac{dg}{dv}$$

which is a vector in the Lie algebra $sl_2(\mathbf{C})$ of 2 by 2 complex matrices of trace zero. As v varies $c(v)$ inscribes a curve on this Lie algebra. Indeed it is well known (e.g. [13], p. 71) that this curve determines the original curve $g(v)$ up to left translation by an arbitrary constant element of G . Here is an explicit formula for the curve $c(v)$. It is easy but rather tedious to verify it.

$$c(v) = \begin{pmatrix} 0 & 1 \\ T & 0 \end{pmatrix}, \quad \text{where } T = \frac{1}{2} \operatorname{sgn}(\kappa') (Q - i)$$

and Q is as in §6. It follows that the inversive curvature Q determines the curve up to an orientation preserving inversive automorphism.

§8. RELATION WITH CARTAN'S MOVING FRAMES

Let us sketch a more usual way of obtaining a Frenet lift. The connection with the Schwartzian described here can be found, for example, in Cartan's book [4] and very succinctly in [7]. The canonical line bundle

$$p: \xi \rightarrow \mathbf{P}^1(\mathbf{C})$$

has a pedestrian description (away from the zero-section) as:

$$\begin{array}{ccc} (z_1, z_2) \in \xi - \{\text{zero section}\} = \mathbf{C}^2 - \{0\} & & \\ \downarrow & p \downarrow & \\ z = \frac{z_1}{z_2} \leftrightarrow [z_1, z_2] \in \mathbf{P}^1(\mathbf{C}) & & \end{array}$$

Let $\sigma: (\alpha, \beta) \rightarrow \mathbf{R}^2 \subset \mathbf{P}^1(\mathbf{C})$ be a curve; we choose an arbitrary lift $\hat{\sigma} = (z_1(t), z_2(t))$ and set $f_1 = \lambda \hat{\sigma}$, $f_2 = \dot{f}_1 = \lambda(z_1, z_2) + \lambda(\dot{z}_1, \dot{z}_2)$, where $\dot{} = \frac{d}{dt}$. Thus (f_1, f_2) is a frame in \mathbf{C}^2 . We try to choose λ so that this frame has area 1. The condition on λ is:

$$\begin{aligned} 1 &= \text{Area}(f_1, f_2) = \text{Area}(\lambda(z_1(t), z_2(t)), \lambda(\dot{z}_1, \dot{z}_2)) \\ &= \lambda^2(z_1 \dot{z}_2 - z_2 \dot{z}_1), \quad \text{or } 1 = -(\lambda z_2)^2 \dot{z} \end{aligned}$$

Thus $\lambda = \frac{i}{z_2 \sqrt{\dot{z}}}$ will do, and we have

$$\begin{aligned} f_1 &= \frac{i}{\sqrt{\dot{z}}} (z, 1), \\ \text{and } f_2 &= \dot{f}_1 = -\frac{1}{2} i \ddot{z} \dot{z}^{-3/2} (z, 1) + i \dot{z}^{-1/2} (z, 0). \end{aligned}$$

Finally a calculation shows that $\dot{f}_2 = S f_1$, where $S = \frac{3}{4} \ddot{z}^2 \dot{z}^{-2} - \frac{1}{2} \ddot{z} \dot{z}^{-1}$.

Of course S is the *Schwartzian derivative* which this calculation interprets as a "curvature" of σ . Now the Schwartzian S depends on the particular parametrization which is used for the curve. For our purposes we wish to use