

**Zeitschrift:** L'Enseignement Mathématique  
**Band:** 38 (1992)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE CATEGORY OF NILMANIFOLDS

**Kurzfassung**

**Autor:** Oprea, John

**DOI:** <https://doi.org/10.5169/seals-59480>

**Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

**Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

**Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 07.10.2024

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## THE CATEGORY OF NILMANIFOLDS

by John OPREA

ABSTRACT. The techniques of rational homotopy theory are used to compute the category of a nilmanifold:  $\text{cat}(M) = \dim M = \text{rank}(\pi_1 M)$ . This information is of interest to dynamicists since the theorem of Lusternik-Schnirelmann then shows that the number of critical points of a smooth function of  $M$  is bounded below by  $\text{rank}(\pi_1 M) + 1$ .

### INTRODUCTION

As a first step to understanding the structure of certain dynamical systems on nilmanifolds, one might hope to have computable lower bounds on the number of critical points of smooth functions. Of course, one is then led to the Lusternik-Schnirelmann definition of category and their well-known result that category  $(+ 1)$  is such a bound. Unfortunately, category is rarely computable, so those who require numerical bounds often employ the fact that category majorizes cuplength. Hence cuplength (which, generally, is a more computable homotopy invariant than category) is the numerical invariant frequently sought for in order to provide a lower bound for the number of critical points of smooth functions on a manifold.

Indeed, some time ago, for the reasons above, Chris McCord asked me if I knew of a formula for the cuplength of a nilmanifold. I did not then, and after many computations I do not now! Thus, I pose:

QUESTION. What is the cuplength (with  $\mathbf{Q}$ -coefficients say) of a nilmanifold?

Suprisingly, however, the need for such knowledge by dynamicists is obviated by the following.