

13. Connections with Ergodic Theory

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **38 (1992)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

For other results connecting discrepancy and the boundedness of the partial quotients, see the papers of Niederreiter [218] and Dupain and Sós [94, 95]. Also see Beck and Chen [25] and Richert [258].

We can also consider the so-called L^2 discrepancy, T_n , defined as follows: let

$$R_n(t) = \frac{S_n([0, t), \omega)}{n} - t$$

and put

$$T_n(\omega) = \left(\int_0^1 R_n^2(t) dt \right)^{1/2}.$$

It is possible to generalize the definitions of D_n and T_n to the multi-dimensional case, though we omit the details. By appealing to numbers with bounded partial quotients, Davenport [73] constructed sequences in two dimensions with low L^2 discrepancy. Also see Proinov [250, 251, 252].

Another measure connected with sequences is called *dispersion*. Let $\omega = (x_1, x_2, \dots)$ and define the dispersion

$$d_n(\omega) = \sup_{x \in [0, 1]} \min_{1 \leq k \leq n} |x - x_k|,$$

essentially half the distance between the most widely separated points of the sequence x_1, x_2, \dots, x_n . (Compare with the function $\delta_\theta(n)$ in Section 11.)

Niederreiter [221] considered the dispersion of the sequence $\{n\theta\}$. He showed that if θ has bounded partial quotients, then $d_n(\omega) = O(1/n)$. He also gave a more detailed estimate, showing that $d_n(\omega)$ is approximately $K(\theta)/4n$. Also see Drobot [93] and Larcher [311].

13. CONNECTIONS WITH ERGODIC THEORY

Let θ be irrational, $\omega = (\theta, 2\theta, \dots)$ and $S_n(I, \omega)$ be defined as in the previous section. Veech [293] developed connections between S_n and ergodic theory. We mention one result that is number-theoretic in nature. Let $x_n = S_n(I, \omega) \bmod 2$, and define

$$\mu_\theta(I) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{1 \leq k \leq n} x_k,$$

if the limit exists. Then Veech showed that $\mu_\theta(I)$ exists for all $I \subseteq [0, 1)$ if and only if the partial quotients of θ are bounded.

For other connections with ergodic theory, see the papers of Stewart [286]; del Junco [154]; Dani [70, 72]; and Baggett and Merrill [14, 15].

14. PSEUDO-RANDOM NUMBER GENERATION

Lehmer [183] introduced the *linear congruential method* for pseudo-random number generation. Let X_0, m, a, c be given, and define

$$X_{k+1} = aX_k + c \pmod{m},$$

for $k \geq 0$. For this to be a good source of “random” numbers, we want the sequence X_k to be uniformly distributed, as well as the sequence of pairs (X_k, X_{k+1}) , triples, etc.

A test for randomness called the *serial test* on pairs (X_k, X_{k+1}) amounts to the two-dimensional version of the discrepancy mentioned above in Section 12. This turns out to be essentially the function $\rho(\mathbf{g}, m)$ defined in Section 10. Thus linear congruential generators that pass the pairwise serial test arise from rationals a/m having small partial quotients in their continued fraction expansion. See the papers of Dieter [87, 88]; Niederreiter [219, 220, 222]; Knuth [170, Section 3.3.3]; and Borosh and Niederreiter [42].

15. FORMAL LANGUAGE THEORY

Let $w = w_0w_1w_2 \cdots$ be an infinite word over a finite alphabet. We say that the finite word $x = x_0x_1 \cdots x_n$ is a *subword* of w if there exists $m \geq 0$ such that $w_{m+i} = x_i$, for $0 \leq i \leq n$. We say that w is *k-th power free* if x^k is never a subword of w , for all nonempty words x . Here is a classical example: let $s(n)$ denote the number of 1's in the binary expansion of n . Then the infinite word of Thue-Morse

$$t = t_0t_1t_2 \cdots = 0110100110010110 \cdots,$$

defined by $t_n = s(n) \pmod{2}$, is cube-free.

Another way to define infinite words is as the fixed point of a homomorphism on a finite alphabet. For example, the Thue-Morse word t is a fixed point of φ , where $\varphi(0) = 01$ and $\varphi(1) = 10$.