

4. S^1 -FIBRATIONS

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

indeed by [GN₁, Proposition 5.7], for odd p , Γ is cyclic of order $2p^2$. The proof there also shows that $2[\gamma_{1,1}]$ is of order p^2 and that $p[\gamma_{0,p}]$ is of order 2 in Γ , so $[\gamma_{2,2+p^2}]$ generates Γ .

(D) THE PROJECTIVE PLANE

We saw that when X is aspherical and $\chi(X) \neq 0$ then $\Gamma = 0$ and so our first order invariants vanish. In the presence of non-trivial higher homotopy these invariants need not vanish, despite $\chi(X) \neq 0$, as demonstrated by the example of the real projective plane $X = P^2$.

Write $G \equiv \pi_1(P^2) \cong \mathbf{Z}/2$; denote the generator of G by t . Give P^2 the customary cell structure consisting of one cell in each of dimensions 0, 1, and 2. The universal cover \tilde{P}^2 is naturally identified with S^2 and the corresponding cellular chain complex is:

$$C_2(S^2) \xrightarrow{1+t^{-1}} C_1(S^2) \xrightarrow{t^{-1}-1} C_0(S^2).$$

Every element of Γ can be represented by a basepoint preserving homotopy $F: P^2 \times I \rightarrow P^2$ with $F_0 = F_1 = \text{id}_{P^2}$. We have $\tilde{F}_0 = \tilde{F}_1 = \text{id}_{S^2}$ because the basepoint is preserved. It is easy to verify that the corresponding chain homotopy $\tilde{D}_*: C_*(S^2) \rightarrow C_*(S^2)$ is then zero on $C_0(S^2)$ and takes \tilde{e}_1 to $\tilde{e}_2 m(1-t^{-1})$ where $m \in \mathbf{Z}$. By elementary obstruction theory, there exists $F \equiv F^{(m)}$ realizing any $m \in \mathbf{Z}$. In this case $\text{trace}(\tilde{\partial} \otimes \tilde{D}) = (1+t^{-1}) \otimes m(1-t^{-1})$ which is homologous to the canonical form $mt^{-1} \otimes tt^{-1} - mt^{-1} \otimes tt^{-2}$. Since $\chi(P^2) = 1 \neq 0$, the Gottlieb group $\eta_{\#}(\Gamma) \equiv \mathcal{G}(P^2) = 0$ and so the derivation $\tilde{X}_1(P^2)$ is a homomorphism and need not be distinguished from its cohomology class $\tilde{\chi}_1(P^2) \in H^1(\Gamma, HH_1(\mathbf{Z}(\mathbf{Z}/2))) \cong \text{Hom}(\Gamma, HH_1(\mathbf{Z}(\mathbf{Z}/2)))$. It follows that

$$\tilde{\chi}_1(P^2) ([F^{(m)}]) = (m, -m) \in \mathbf{Z}/2 \oplus \mathbf{Z}/2 \cong HH_1(\mathbf{Z}(\mathbf{Z}/2)).$$

In particular, when m is odd $\tilde{\chi}_1(P^2) ([F^{(m)}]) \neq 0$. On the other hand, this shows $\chi_1(P^2) = 0$.

4. S^1 -FIBRATIONS

In this section we investigate the first order Euler characteristic of the total space of an orientable Serre fibration with S^1 -fiber.

Let $S^1 \rightarrow X \xrightarrow{\pi} B$ be an orientable Serre fibration where B is a (not necessarily finite) connected CW complex and X has the homotopy type of a finite complex. By classical obstruction theory, fiber homotopy

equivalence classes of orientable S^1 -fibrations over a CW complex B are classified by the integral cohomology group $H^2(B; \mathbf{Z})$. Given an element $e \in H^2(B; \mathbf{Z}) \cong [B, \mathbf{C}P^\infty]$ one obtains a principal $U(1)$ -bundle over B by pulling back, via a continuous map $B \rightarrow \mathbf{C}P^\infty$ representing e , the $U(1)$ -bundle associated to the canonical complex line bundle over the infinite dimensional complex projective space $\mathbf{C}P^\infty$. Thus we can assume, without loss of generality, that $S^1 \rightarrow X \xrightarrow{\pi} B$ is a principal $U(1)$ -bundle. In particular, there is a free $U(1)$ -action on X which we will write as $\Phi: X \times S^1 \rightarrow X$. Let $\tau \in \Gamma \equiv \pi_1(\mathcal{C}(X), 1)$ be the element represented by Φ ($\Phi = \Phi^\tau$ in the notation of §1). For any coefficient ring R , let $\{r\} \in H_1(X; R)$ denote the image of τ under the composite:

$$\Gamma \xrightarrow{\eta} \pi_1(X) \rightarrow H_1(X) \rightarrow H_1(X; R) .$$

Also, let e_R be the image of the element $e \in H^2(B; \mathbf{Z})$ which classifies $S^1 \rightarrow X \xrightarrow{\pi} B$ under the homomorphism $H^2(B; \mathbf{Z}) \rightarrow H^2(B; R)$.

LEMMA 4.1. *If \mathbf{F} is a field, then $\{\tau\} \in H_1(X; \mathbf{F})$ is non-zero if and only if $e_{\mathbf{F}} = 0$.*

Proof. Consider the Gysin homology sequence for the fibration $S^1 \rightarrow X \xrightarrow{\pi} B$:

$$\cdots \rightarrow H_2(B; \mathbf{F}) \xrightarrow{e_{\mathbf{F}} \cap} H_0(B; \mathbf{F}) \xrightarrow{\theta_0} H_1(X; \mathbf{F}) \xrightarrow{\pi_*} H_1(B; \mathbf{F}) \rightarrow 0 .$$

Since $H_2(B; \mathbf{F}) \xrightarrow{e_{\mathbf{F}} \cap} H_0(B; \mathbf{F}) \cong \mathbf{F}$ is just evaluation of the cohomology class $e_{\mathbf{F}}$ on homology, θ_0 is non-zero if and only if $e_{\mathbf{F}} = 0$. Let $v \in X$ be a basepoint and let $\{\pi(v)\} \in H_0(B; \mathbf{F})$ be the generator determined by the inclusion of $\pi(v)$ into B . The fact that $\theta_0(\{v\}) = \{\tau\}$ follows from the naturality of the Gysin sequence homology sequence, by mapping the Gysin sequence of the trivial fibration $S^1 \rightarrow S^1 \rightarrow \pi(v)$, via the homomorphism induced by inclusion, into the Gysin sequence for $S^1 \rightarrow X \xrightarrow{\pi} B$. \square

THEOREM 4.2. *Let \mathbf{F} be a field. If $e_{\mathbf{F}} \neq 0$ then $\chi_1(X; \mathbf{F})(\tau) = 0$. If $e_{\mathbf{F}} = 0$ then $H_*(B; \mathbf{F})$ is finite dimensional over F and $\chi_1(X; \mathbf{F})(\tau) = -\chi(B; \mathbf{F})\{\tau\}$ where $\chi(B; \mathbf{F}) = \sum_{i \geq 0} (-1)^i \dim_{\mathbf{F}} H_i(B; \mathbf{F})$.*

Proof. In this proof, all homology and cohomology groups will have coefficients in the field \mathbf{F} . Since B is the orbit space of the $U(1)$ -action on X given by Φ , there is a commutative square:

$$\begin{array}{ccc} X \times S^1 & \xrightarrow{\Phi} & X \\ \pi \times \text{id} \downarrow & & \pi \downarrow \\ B \times S^1 & \xrightarrow{p} & B \end{array}$$

where $p: B \times S^1 \rightarrow B$ is projection. This square induces a commutative ladder mapping the Gysin homology sequence of $S^1 \rightarrow X \times S^1 \xrightarrow{\pi \times \text{id}} B \times S^1$ to the Gysin homology sequence of $S^1 \rightarrow X \xrightarrow{\pi} B$:

$$\begin{array}{ccccccc} H_i(B \times S^1) & \xrightarrow{\theta'} & H_{i+1}(X \times S^1) & \xrightarrow{(\pi \times \text{id})^*} & H_{i+1}(B \times S^1) & \rightarrow & H_{i-1}(B \times S^1) \\ p_* \downarrow & & \Phi_* \downarrow & & p_* \downarrow & & p_* \downarrow \\ H_i(B) & \xrightarrow{\theta} & H_{i+1}(X) & \xrightarrow{\pi_*} & H_{i+1}(B) & \xrightarrow{e_{\mathbf{F}} \cap} & H_{i-1}(B) \end{array}$$

For each integer $0 \leq i \leq \dim X$ choose a basis $\{b_1^i, \dots, b_{\beta_i}^i\}$ for $H_i(X)$ such that for some integer $m_i \leq \beta_i$ $\{b_{m_i+1}^i, \dots, b_{\beta_i}^i\}$ is a basis for the kernel of $\pi_*: H_i(X) \rightarrow H_i(B)$. The corresponding dual basis for $H^i(X)$ will be denoted by $\{\bar{b}_1^i, \dots, \bar{b}_{\beta_i}^i\}$. Since we are using coefficients in a field, we make the identifications $H_*(B \times S^1) \cong H_*(B) \otimes H_*(S^1)$ and $H_*(X \times S^1) \cong H_*(X) \otimes H_*(S^1)$ via the natural isomorphism given by the homology exterior product. Let $u \in H_1(S^1)$ be the generator determined by the standard orientation of S^1 . Using Definition B_1 ,

$$\chi_1(X; \mathbf{F})(\tau) = \sum_{k \geq 0} (-1)^{k+1} \sum_{j=1}^{\beta_k} \bar{b}_j^k \cap \Phi_*(b_j^k \otimes u).$$

Consider $b_j^i \otimes u \in H_{i+1}(X \times S^1)$ where $m_i + 1 \leq j \leq \beta_i$. Since b_j^i lies in $\ker \pi_*$, the exactness of the Gysin sequence implies that $b_j^i \otimes u = \theta'(c \otimes u)$ for some $c \in H_i(B)$. Consequently,

$$\Phi_*(b_j^i \otimes u) = \Phi_*(\theta'(c \otimes u)) = \theta(p_*(c \otimes u)) = 0$$

because $p_*(c \otimes u) = 0$. It follows that

$$(4.3) \quad \chi_1(X; \mathbf{F})(\tau) = \sum_{k \geq 0} (-1)^{k+1} \sum_{j=1}^{m_k} \bar{b}_j^k \cap \Phi_*(b_j^k \otimes u).$$

For each k , the set $\{\pi_*(b_1^k), \dots, \pi_*(b_{m_k}^k)\}$ is a basis for the image of $\pi_*: H_k(X) \rightarrow H_k(B)$. Extend this set (in any manner) to basis for $H_k(B)$ and let $\{\overline{\pi_*(b_1^k)}, \dots, \overline{\pi_*(b_{m_k}^k)}\}$ denote the corresponding portion of the dual basis for $H^k(B)$. Then $\bar{b}_j^k = \pi^*(\overline{\pi_*(b_j^k)})$, $0 \leq j \leq m_k$. Consider the commutative diagram:

$$\begin{array}{ccc} H^k(B \times S^1) & \xrightarrow{(\pi \times \text{id})^*} & H^k(X \times S^1) \\ p^* \uparrow & & \Phi^* \uparrow \\ H^k(B) & \xrightarrow{\pi^*} & H^k(X) \end{array}$$

Then, for $0 \leq j \leq m_k$,

$$\begin{aligned}
\bar{b}_j^k \cap \Phi_*(b_j^k \otimes u) &= \Phi_*(\Phi^*(\bar{b}_j^k) \cap (b_j^k \otimes u)) \\
&= \Phi_*(\Phi^*(\pi^*(\overline{\pi_*(b_j^k)}))) \cap (b_j^k \otimes u) \\
&= \Phi_*((\pi \times \text{id})^*(p^*(\overline{\pi_*(b_j^k)}))) \cap (b_j^k \otimes u) \\
&\quad \text{using the above diagram} \\
&= \Phi_*((\bar{b}_j^k \otimes 1) \cap (b_j^k \otimes u)) \\
&= \Phi_*((\bar{b}_j^k \cap b_j^k) \otimes u) = \Phi_*({\nu} \otimes u) = \{\tau\}
\end{aligned}$$

where $\{\nu\}$ is the natural generator of $H_0(X)$ determined by the inclusion of the basepoint ν into X . From the proof of Lemma 4.1, $\Phi_*({\nu} \otimes u) = \{\tau\}$. Substituting the above computation into Formula 4.3 yields $\chi_1(X; \mathbf{F})(\tau) = (\sum_{k \geq 0} (-1)^{k+1} m_k) \{\tau\}$. If $e_{\mathbf{F}} \neq 0$ then Lemma 4.1 implies that $\{\tau\} = 0$ and so $\chi_1(X; \mathbf{F})(\tau) = 0$. Thus the conclusion of the theorem is valid in this case. If $e_{\mathbf{F}} = 0$ then from the portion

$$H_k(X) \xrightarrow{\pi_*} H_k(B) \xrightarrow{e_{\mathbf{F}} \cap} H_{k-2}(B)$$

of the Gysin homology sequence we deduce that π_* is onto and consequently $m_k = \dim_{\mathbf{F}} H_k(B, \mathbf{F})$. Thus $\dim_{\mathbf{F}} H_*(B, \mathbf{F})$ is finite and $\sum_{k \geq 0} (-1)^{k+1} m_k = -\chi(B; \mathbf{F})$. \square

Theorem 4.2 can be used to recalculate $\chi_1(X; \mathbf{F})$ in Examples 3.8 and 3.9.

Next, we consider integer coefficients. Suppose that $S^1 \rightarrow X \xrightarrow{\pi} B$ is a smooth orientable $U(1)$ -bundle over a smooth, closed, oriented manifold B . Let λ be the one dimensional subbundle of the tangent bundle of X consisting of vectors which are tangent to the circle fibers and let ν be a complementary bundle to λ . Then $\nu \cong \pi^*(T_B)$ where T_B is the tangent bundle of B . Let $[B] \in H_n(B; \mathbf{Z})$ be the fundamental class of B where $n = \dim B$. The Euler class, $\text{Eul}(\nu) \in H^n(X; \mathbf{Z})$, is given by

$$\text{Eul}(\nu) = \text{Eul}(\pi^*(T_B)) = \pi^*(\text{Eul}(T_B)) = \chi(B) \pi^*([B]^*)$$

where $[B]^* \in H^n(B; \mathbf{Z})$ is the generator determined by the condition $[B]^*([B]) = 1$; see [MS, Corollary 11.12]. The Gysin homology sequence for $S^1 \rightarrow X \xrightarrow{\pi} B$ determines a fundamental class for X ; $[X] \in H_{n+1}(X)$ is the image of $[B]$ under the homomorphism $\theta_n: H_n(B; \mathbf{Z}) \rightarrow H_{n+1}(X; \mathbf{Z})$. For any closed oriented m -dimensional manifold M , let $\text{PD}_M: H^i(M) \rightarrow H_{m-i}(M)$ be the Poincaré duality isomorphism explicitly given by $\text{PD}_M(x) = (-1)^{i(m-i)} x \cap [M]$ where $x \in H^i(M)$ and $[M] \in H_m(M)$ is the

fundamental class $((-1)^{i(m-i)})$ appears because of our use of Dold's sign conventions). An immediate consequence of Theorem 3.1 of [GN₂] is the following computation of $\chi_1(X)$ (with integer coefficients):

THEOREM 4.4. $\chi_1(X)(\tau) = -\text{PD}_X(\text{Eul}(v))$. \square

THEOREM 4.5. *Under the above hypotheses, $\chi_1(X)(\tau) = -\chi(B)\{\tau\}$.*

Proof. There is a Poincaré duality isomorphism between the Gysin homology sequence and the Gysin cohomology sequence, a portion of which is shown below:

$$\begin{array}{ccccc} H_0(B; \mathbf{Z}) & \xrightarrow{\theta_0} & H_1(X; \mathbf{Z}) & \xrightarrow{\pi_*} & H_1(B; \mathbf{Z}) \\ \text{PD}_B \uparrow & & \text{PD}_X \uparrow & & \text{PD}_B \uparrow \\ H^n(B; \mathbf{Z}) & \xrightarrow{\pi^*} & H^n(X; \mathbf{Z}) & \rightarrow & H^{n-1}(B; \mathbf{Z}) \end{array}$$

Let $v \in X$ be a basepoint, and let $\{\pi(v)\} \in H_0(B; \mathbf{Z})$ be the generator determined by the inclusion of $\pi(v)$ into B . From the above diagram, $\text{PD}_X(\pi^*([B]^*)) = \theta_0(\{\pi(v)\})$. Also, from the proof of Lemma 4.1, $\theta_0(\{\pi(v)\}) = \{\tau\}$. Thus $\text{PD}_X(\text{Eul}(v)) = \chi(B)\{\tau\}$. Regarding the free $U(1)$ -action on X as a flow, we can now invoke Theorem 4.4 to conclude that $\chi(B)\{\tau\} = -\chi_1(X)(\tau)$. \square

Example 4.6. Let Σ_g be a closed oriented surface of genus $g > 1$ and let L_n be a complex line bundle over Σ_g with Chern number n . Let $M_{n,g}$ be the total space of the $U(1)$ -bundle associated to L_n . Then $M_{n,g}$ is a closed oriented aspherical 3-manifold which fibers over Σ_g . The center of $\pi_1(M_{n,g})$ is the infinite cyclic group generated by τ (represented by a circle fiber); the image, $\{\tau\}$, of τ in $H_1(M_{n,g}) \cong \mathbf{Z}^{2g} \oplus \mathbf{Z}/n$ generates the \mathbf{Z}/n summand. By Theorem 4.5, $\chi_1(M_{n,g}): \mathbf{Z} \rightarrow H_1(M_{n,g})$ is given by $\chi_1(M_{n,g})(\tau) = (2g - 2)\{\tau\}$.

Let T^n , where $n > 1$, be the n -torus (i.e. the n -fold product of copies of $U(1)$). Let X be a closed oriented smooth manifold and let $\rho: T^n \times X \rightarrow X$ be a smooth free action of T^n . This action defines a homomorphism $\bar{\rho}: T^n \rightarrow \text{Diff}(X)$ where $\text{Diff}(X)$ is the diffeomorphism group of X . Let $\Gamma_\rho \subset \Gamma$ be the image of the composite:

$$\pi_1(T^n, 1) \xrightarrow{\bar{\rho}\#} \pi_1(\text{Diff}(X), \text{id}) \rightarrow \pi_1(\mathcal{L}(X), \text{id}) = \Gamma.$$

PROPOSITION 4.7. *The restriction of $\chi_1(X): \Gamma \rightarrow H_1(X)$ to Γ_ρ is the zero homomorphism.*

Proof. Since $n > 1$, if $T \subset T^n$ is a circle subgroup then $\chi(X/T) = 0$. Applying Theorem 4.5 to the bundle $T \rightarrow X \rightarrow X/T$ yields the conclusion. \square

COROLLARY 4.8. *If $n > 1$ then $\chi_1(T^n): \mathbf{Z}^n \rightarrow \mathbf{Z}^n$ is zero.* \square

5. A HIGHER ANALOG OF GOTTLIEB'S THEOREM

Let G be a group of type \mathcal{F} . Gottlieb's theorem (see Propositions 1.3 and 2.4) asserts that if $\chi(G) \neq 0$ then $Z(G)$, the center of G , is trivial. We prove an analogous theorem for $\chi_1(G; \mathbf{Q})$: if $\chi_1(G; \mathbf{Q}) \neq 0$ then the center of G is infinite cyclic provided G satisfies an extra hypothesis (explained below) related to the Bass Conjecture; see Proposition 5.2 and Theorem 5.4.

Throughout this section R will be a commutative ground ring. Let S be any associative R -algebra with unit. The Hochschild homology group $HH_0(S)$ is the R -module $S/[S, S]$ where $[S, S]$ is the R -submodule of S generated by $\{ab - ba \mid a, b \in S\}$; see §2. Recall that $K_0(S)$ is the abelian group F/A where F is the free abelian group generated by the set of all isomorphism classes $[M]$ of finitely generated projective right S -modules $M \subset \bigoplus_{i=1}^{\infty} S$ and A is the subgroup of F generated by relations of the form $[M_1 \oplus M_2] - [M_1] - [M_2]$. Since a finitely generated projective module is the image of a finitely generated free module under an idempotent homomorphism, each element of $K_0(S)$ can be represented by an idempotent matrix over S . The *Hattori-Stallings* trace $T_0: K_0(S) \rightarrow HH_0(S)$ is defined as follows. Let $A: M \rightarrow M$ be an idempotent endomorphism of a free, finitely generated right S -module M representing $x \in K_0(S)$. If $[A]$ is the matrix of A with respect to a given basis for M then $T_0(x)$ is defined to be $T_0([A]) \in HH_0(S)$.

Consider the groupring, RG , of a group G over R . Then $HH_0(RG)$ is naturally isomorphic to the free R -module generated by G_1 , the set of conjugacy classes of G (see §2 for an explanation in the case $R = \mathbf{Z}$). Recall that for $g \in G$ we write $C(g) \in G_1$ for the conjugacy class of g , $HH_0(RG)_{C(g)}$ for the summand of $HH_0(RG)$ corresponding to $C(g)$ and $x_{C(g)}$ for the $C(g)$ -component of $x \in HH_0(RG)$. Also write $HH_0(RG) = HH_0(RG)_{C(1)} \oplus HH_0(RG)'$ where $1 \in G$ is the identity element of G , and $HH_0(RG)'$ is the direct sum of the remaining summands. The augmentation homomorphism $\varepsilon: RG \rightarrow R$ induces a homomorphism $\varepsilon_*: HH_0(RG) \rightarrow HH_0(R) = R$.