

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GAUSS-BONNET FORMULA FOR LOCALLY SYMMETRIC SPACES OF NONCOMPACT TYPE
Autor: Leuzinger, Enrico
DOI: <https://doi.org/10.5169/seals-87876>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 30.01.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ON THE GAUSS-BONNET FORMULA
FOR LOCALLY SYMMETRIC SPACES OF NONCOMPACT TYPE

by Enrico LEUZINGER

ABSTRACT. Let X be a Riemannian symmetric space of noncompact type and rank ≥ 2 and let Γ be a non-uniform, irreducible lattice in the group of isometries of X . A Gauss-Bonnet formula for the locally symmetric quotient $V = \Gamma \backslash X$ was first proved by G. Harder. We present a new simple proof which is based on an exhaustion of V by Riemannian polyhedra with uniformly bounded second fundamental forms.

INTRODUCTION

The generalized Gauss-Bonnet theorem of C.B. Allendoerfer, A. Weil and S.S. Chern asserts that the Euler characteristic of a *closed* Riemannian manifold (M, g) is given by

$$\chi(M) = \int_M \omega_g$$

where the Gauss-Bonnet-Chern form $\omega_g = \Psi_g dv_g$ is (locally) computable from the metric g (see [AW], [C]).

In several articles J. Cheeger and M. Gromov investigated the Gauss-Bonnet theorem for *open* complete Riemannian manifolds with bounded sectional curvature and finite volume. They in particular showed that such manifolds M^n admit an exhaustion by compact manifolds with smooth boundary, M_i^n , such that $\text{Vol}(\partial M_i^n) \rightarrow 0$ ($i \rightarrow \infty$) and for which the second fundamental forms $\text{II}(\partial M_i^n)$ are uniformly bounded (see [CG1], [CG2], [CG3] and also [G] 4.5.C). By a formula of Chern one has $\chi(M_i^n) = \int_{M_i^n} \omega_g + \int_{\partial M_i^n} \eta_i$ where η_i is a certain form on the boundary ∂M_i^n (see [C]). The above two properties imply that $\lim_{i \rightarrow \infty} \int_{\partial M_i^n} \eta_i = 0$ and hence $\chi(M_i^n) = \int_{M^n} \omega_g$ for sufficiently large i . As a consequence the Gauss-Bonnet theorem holds whenever $\chi(M_i^n) = \chi(M^n)$ for all sufficiently large i .

We now consider a Riemannian symmetric space X of noncompact type and rank ≥ 2 and a non-uniform, torsion-free lattice Γ in the group of isometries of X . The quotient $V = \Gamma \backslash X$ is a locally symmetric space with bounded non-positive sectional curvature and finite volume. Locally symmetric spaces thus provide important examples for the above class considered by Cheeger and Gromov. If Γ is irreducible a remarkable theorem of G. A. Margulis asserts that Γ is *arithmetic* (see [Z], Ch. 6). For quotients of such lattices the Gauss-Bonnet formula was first proved by G. Harder (see [H]). Following M. S. Raghunathan [R1] he explicitly constructed a smooth exhaustion function h on V which has no critical points outside a compact set. A certain defect of the function h , however, is the quite complicated geometry of its sublevel sets (their second fundamental forms, for instance, are not uniformly bounded). As a consequence the proof given in [H] involves rather long and technical estimates.

The purpose of the present note is two-fold. On the one hand to give a new, more geometric proof of the Gauss-Bonnet theorem for locally symmetric spaces, which avoids the technically complicated estimates of [H]. And, on the other hand, to provide an explicit (and independent) illustration of general results in [CG3].

Our approach is based on an exhaustion $V = \bigcup_{s \geq 0} V(s)$ of locally symmetric spaces *not* by *smooth* submanifolds but by *polyhedra*, i.e. compact submanifolds with corners (see [L2]). The corners which appear here are naturally related to the geometry of V at infinity (and therefore should not be smoothed). Moreover, for each $s \geq 0$ the polyhedron $V(s)$ is a strong deformation retract of V (see [L3]). The essential new feature of this exhaustion is that the boundaries of $\partial V(s)$ consist of subpolyhedra of $V(s)$ which are projections of pieces of horospheres in X . As a consequence their second fundamental forms are uniformly bounded. This property together with the generalized Gauss-Bonnet formula for Riemannian polyhedra of Allendoerfer-Weil and Chern leads to a considerably simplified new proof of the Gauss-Bonnet theorem for locally symmetric spaces (see Theorem 4.1).

NOTATION. Explicit constants are irrelevant for our purpose. If f and g are positive real valued functions on a set S we thus simply write $f \prec g$ if there is a constant $c > 0$ such that $f(s) \leq cg(s)$ for all $s \in S$.

1. THE FORMULA OF ALLENDOERFER AND WEIL

A C^∞ (resp. C^ω) manifold with corners is a topological Hausdorff space locally modeled upon a product of lines and half-lines and such that coordinate changes are of class C^∞ (resp. C^ω). For precise definitions and basic information about this concept we refer to [DH]. A *Riemannian polyhedron* is a compact manifold with corners equipped with a Riemannian metric.

Let \mathcal{P}^n be an n -dimensional Riemannian polyhedron with boundary consisting of a finite family of lower dimensional subpolyhedra

$$\mathcal{P}_E^{n-k} \quad (0 \leq k \leq n-1)$$

and with Riemannian metric induced from \mathcal{P}^n . The *outer angle* $O(p)$ at a point p of \mathcal{P}_E^{n-k} is defined as the set of all unit tangent vectors $v \in T_p\mathcal{P}^n$ such that $\langle v, w \rangle_p \leq 0$ for all w in the tangent cone of \mathcal{P}^n at p . Note that $O(p)$ is a spherical cell bounded by “great spheres” in the $(k-1)$ -dimensional unit sphere of the normal space of $\mathcal{P}_E^{n-k} \subset \mathcal{P}^n$ at p . In [AW] Allendoerfer and Weil define a certain real valued function $\Psi_{E,k}$ on the outer angles of \mathcal{P}_E^{n-k} . The explicit form of this function will not be needed in this paper. We shall only use the fact that $\Psi_{E,k}$ is locally computable from the components of the metric and the curvature tensor of \mathcal{P}^n and from the components of the second fundamental forms $\Pi_Z(p), Z \in O(p)$, of \mathcal{P}_E^{n-k} in \mathcal{P}^n . Let Ψdv denote the Gauss-Bonnet-Chern form on \mathcal{P}^n and dv_E (resp. $d\omega_{k-1}$) the volume element of \mathcal{P}_E^k (resp. of the standard unit sphere S^{k-1}). The *inner Euler characteristic* χ' of \mathcal{P}^n is by definition the Euler characteristic of the open complex consisting of all inner cells in an arbitrary simplicial subdivision of \mathcal{P}^n .

We can now state the generalized Gauss-Bonnet formula of Allendoerfer-Weil for Riemannian polyhedra (see [AW]).

PROPOSITION 1.1. *Let \mathcal{P}^n be a Riemannian polyhedron with boundary consisting of a finite family of subpolyhedra \mathcal{P}_E^{n-k} . Then the inner Euler characteristic of \mathcal{P}^n is given by*

$$(-1)^n \chi'(\mathcal{P}^n) = \int_{\mathcal{P}^n} \Psi dv + \sum_{k=1}^n \sum_E \int_{\mathcal{P}_E^{n-k}} \left(\int_{O(p)} \Psi_{E,k} d\omega_{k-1} \right) dv_E(p).$$

2. AN EXHAUSTION OF LOCALLY SYMMETRIC SPACES

Let X be a Riemannian symmetric space of noncompact type and rank ≥ 2 and let Γ be a non-uniform, torsion-free lattice in the group of isometries of X . In this section we briefly describe the basic features of an exhaustion of the locally symmetric space $V = \Gamma \backslash X$ by Riemannian polyhedra, which was previously constructed in [L2].

The idea is to work with a *fundamental set* $\Omega \subset X$ for the discrete (arithmetic) group Γ . Such “coarse” fundamental domains are provided by *reduction theory*; they are finite unions of translates of so-called Siegel sets. We begin with reviewing some facts about linear algebraic groups and set up the notation. Roughly speaking, the lattice Γ determines a “ \mathbb{Q} -structure” on the real Lie group of isometries of X such that Γ is given by integer matrices. The symmetric space X in turn inherits canonical parametrizations adopted to this structure (generalized horocyclic coordinates). Siegel sets are then defined with respect to such parametrizations.

2.1. REDUCTION THEORY AND GEOMETRY AT INFINITY

We denote by G the identity component of the group of isometries of X ; it is a connected, semisimple Lie group with trivial center. We shall always assume in the following that the non-uniform lattice Γ is *irreducible* (see [R2] 5.20). Then, by the arithmeticity theorem of Margulis, there is a connected semisimple linear algebraic group \mathbf{G} defined over \mathbb{Q} , \mathbb{Q} -embedded in a general linear group $\mathbf{GL}(N, \mathbb{C})$, and a Lie group isomorphism $p : G \rightarrow \mathbf{G}(\mathbb{R})^0$ such that $p(\Gamma)$ is *arithmetic*, i.e. $p(\Gamma) \subset \mathbf{G}(\mathbb{Q}) \subset \mathbf{GL}(N, \mathbb{C})$ is commensurable with the group $\mathbf{G}(\mathbb{Z}) = \mathbf{G} \cap \mathbf{GL}(N, \mathbb{Z})$ (see [Z] 3.1.6 and 6.1.10). The symmetric space X can be recovered as the manifold of maximal compact subgroups of the identity component of the group $\mathbf{G}(\mathbb{R}) = \mathbf{G} \cap \mathbf{GL}(N, \mathbb{R})$ of \mathbb{R} -rational points of \mathbf{G} . For simplicity we will always identify G with $\mathbf{G}(\mathbb{R})^0$ and Γ with $p(\Gamma)$.

Let \mathbf{S} (resp. \mathbf{T}) be a maximal \mathbb{Q} -split (resp. \mathbb{R} -split) algebraic torus of \mathbf{G} , i.e. a subgroup of \mathbf{G} which is isomorphic over \mathbb{Q} (resp. \mathbb{R}) to the direct product of q (resp. $r \geq q$) copies of \mathbb{C}^* . All such tori are conjugate under $\mathbf{G}(\mathbb{Q}) = \mathbf{G} \cap \mathbf{GL}(N, \mathbb{Q})$ (resp. $\mathbf{G}(\mathbb{R})$) and their common dimension q (resp. r) is called the \mathbb{Q} -rank (resp. \mathbb{R} -rank) of \mathbf{G} . The identity component of $\mathbf{S}(\mathbb{R})$ (resp. $\mathbf{T}(\mathbb{R})$) will be denoted by A (resp. A_0), the corresponding Lie algebras by \mathfrak{a} (resp. \mathfrak{a}_0). The \mathbb{R} -rank of \mathbf{G} coincides with the rank of the symmetric space X , i.e. the maximal dimension of totally geodesic flat subspaces. The choice of a maximal compact subgroup K of G

is equivalent to the choice of a base point x_0 of X . We can choose K with Lie algebra \mathfrak{k} so that under the corresponding Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ of the Lie algebra \mathfrak{g} of G we have $\mathfrak{a} \subseteq \mathfrak{a}_0 \subset \mathfrak{p} \cong T_{x_0}X$. Here \mathfrak{a}_0 is maximal abelian in \mathfrak{p} , i.e. the tangent space at x_0 of the (maximal \mathbb{R} -) flat $A_0 \cdot x_0$ in X . The pair of Lie algebras $(\mathfrak{g}, \mathfrak{a}_0)$ gives rise to the root system ${}_{\mathbb{R}}\Phi$ of the symmetric space. Similarly there is a system of \mathbb{Q} -roots ${}_{\mathbb{Q}}\Phi$ associated to the pair $(\mathfrak{g}, \mathfrak{a})$ (see [B3] §21). It is always possible to choose orderings of ${}_{\mathbb{Q}}\Phi$ and ${}_{\mathbb{R}}\Phi$ such that the restrictions of simple \mathbb{R} -roots of ${}_{\mathbb{R}}\Phi$ to \mathfrak{a} are either *simple* \mathbb{Q} -roots of ${}_{\mathbb{Q}}\Phi$, i.e. the elements of a basis $\Delta = {}_{\mathbb{Q}}\Delta$ of ${}_{\mathbb{Q}}\Phi$, or zero (see [BT] 6.8). The basis ${}_{\mathbb{R}}\Delta$ defines a closed \mathbb{R} -Weyl chamber $\overline{\mathfrak{a}_0^+}$ in \mathfrak{a}_0 and Δ then determines a closed \mathbb{Q} -Weyl chamber $\overline{\mathfrak{a}^+} := \{H \in \mathfrak{a} \mid \alpha(H) \geq 0, \text{ for all } \alpha \in \Delta\}$ in \mathfrak{a} . We set $\overline{A^+} = \exp \overline{\mathfrak{a}^+}$ (resp. $\overline{A_0^+} = \exp \overline{\mathfrak{a}_0^+}$). A \mathbb{Q} -Weyl chamber in X is a translate of the basic chamber $\overline{A^+} \cdot x_0 \subseteq \overline{A_0^+} \cdot x_0$. The elements of Δ are differentials of characters (defined over \mathbb{Q}) of the maximal \mathbb{Q} -split torus \mathbf{S} . It is convenient to identify the elements of Δ also with such characters. When restricted to A their values are denoted by $\alpha(a)$ ($a \in A, \alpha \in \Delta$). Notice that $\overline{A^+} = \{a \in A \mid \alpha(a) \geq 1 \text{ for all } \alpha \in \Delta\}$.

A closed subgroup \mathbf{P} of \mathbf{G} defined over \mathbb{Q} is a *parabolic \mathbb{Q} -subgroup* if \mathbf{G}/\mathbf{P} is a projective variety (see [B3] §11). A *parabolic \mathbb{Q} -subgroup* P of $G = \mathbf{G}(\mathbb{R})^0$ is by definition the intersection of G with a parabolic \mathbb{Q} -subgroup of \mathbf{G} (see [BS]). The conjugacy classes under $\mathbf{G}(\mathbb{Q})$ of parabolic \mathbb{Q} -subgroups are in one-to-one correspondence with the subsets Θ of the (chosen) set Δ of simple \mathbb{Q} -roots; they are represented by the *standard parabolic \mathbb{Q} -subgroups* \mathbf{P}_Θ of \mathbf{G} (see [B3] §21.11). The corresponding standard parabolic \mathbb{Q} -subgroups of G are denoted by P_Θ . The minimal parabolic subgroup $P = P_\emptyset$ has a decomposition $P = UMA$, where U is unipotent and M is reductive; A centralizes M and normalizes U (see [B1]). This yields a (generalized) Iwasawa decomposition for G , i.e. $G = P \cdot K = UMAK$, which implies that P acts transitively on the symmetric space X . The intersection of the maximal compact subgroup K of G with M is maximal compact in M and the quotient $Z = M/(K \cap M)$ is (in general) the Riemannian product of a symmetric space of noncompact type by a (flat) Euclidean space. Let $\tau : M \rightarrow Z$ be the natural projection. Then the “horocyclic coordinate map”

$$\mu : Y = U \times Z \times A \rightarrow X \quad ; \quad (u, \tau(m), a) \mapsto uma \cdot x_0$$

is an isomorphism of analytic manifolds (see [BS] or [B2]).

A *generalized Siegel set* $\mathcal{S} = \mathcal{S}_{\omega, \tau}$ in X (relative to the \mathbb{Q} -Weyl chamber $\overline{A^+} \cdot x_0$) is a subset of X of the form $\mathcal{S}_{\omega, \tau} = \omega A_\tau \cdot x_0$ where ω is relatively compact in UM and, for $\tau > 0$, $A_\tau = \{a \in A \mid \alpha(a) \geq \tau, \alpha \in \Delta\}$. If we define $a_0 \in A$ by $\alpha(a_0) = \tau$ for all $\alpha \in \Delta$, then $A_\tau = A_1 a_0 = \overline{A^+} a_0$ and $\mathcal{C} = A_\tau \cdot x_0 \subset \mathcal{S}$ is a (translate of a) \mathbb{Q} -Weyl chamber in X . Siegel sets provide the building blocks for (approximate) fundamental domains for arithmetic groups. A subset $\Omega \subset X$ is called a *fundamental set* for an arithmetic group Γ if the following two conditions hold

- (i) $X = \Gamma \cdot \Omega$;
- (ii) for every $q \in \mathbf{G}(\mathbb{Q})$ the set $\{\gamma \in \Gamma \mid q\Omega \cap \gamma\Omega \neq \emptyset\}$ is finite.

The existence of fundamental sets is guaranteed by reduction theory for arithmetic groups (see [B1] §13 and §15).

PROPOSITION 2.1 (Borel, Harish-Chandra). *Let \mathbf{G} be a semisimple algebraic group defined over \mathbb{Q} with associated Riemannian symmetric space $X = G/K$. Let \mathbf{P} be a minimal parabolic \mathbb{Q} -subgroup of \mathbf{G} and let Γ be an arithmetic subgroup of $\mathbf{G}(\mathbb{Q})$. Then there exists a generalized Siegel set $\mathcal{S} = \mathcal{S}_{\omega, \tau}$ (with respect to $\overline{A^+} \cdot x_0$) such that, for a (fixed) set $\{q_i \mid 1 \leq i \leq m\}$ of representatives of the finite set of double cosets $\Gamma \backslash \mathbf{G}(\mathbb{Q}) / \mathbf{P}(\mathbb{Q})$, the union $\Omega = \bigcup_{i=1}^m q_i \cdot \mathcal{S}$ is a fundamental set (of finite volume) for Γ in X .*

It will be useful in the sequel to dispose of geometric interpretations of the above algebraic concepts and assertions.

First recall that the symmetric space X , as a Riemannian manifold of nonpositive curvature, has an *ideal boundary at infinity* $\partial_\infty X$. The latter is defined as the set of equivalence classes of asymptotic geodesic rays (see [BGS]). In the same way one also defines the ideal boundary at infinity $\partial_\infty V$ of $V = \Gamma \backslash X$. If Γ is an arithmetic lattice in a group \mathbf{G} of \mathbb{Q} -rank $q = 1$, the boundary $\partial_\infty V$ of the associated locally symmetric space consists of m points (corresponding to the cusps), where m is as in Proposition 2.1. For \mathbb{Q} -rank $q \geq 2$ it turns out that $\partial_\infty V$ is isomorphic to a finite simplicial complex $\Gamma \backslash |\mathcal{T}|$, a geometric realization of the Tits building of \mathbf{G} modulo Γ (see [JM] and [L1]). We recall the construction of the latter.

Let \mathcal{P} be the set of all parabolic \mathbb{Q} -subgroups of \mathbf{G} . The conjugacy classes of elements of \mathcal{P} are in one-to-one correspondence with the subsets Θ of the set Δ of simple \mathbb{Q} -roots. Every conjugacy class has a standard representative denoted by \mathbf{P}_Θ . One can show that the sets of double cosets $\Gamma \backslash \mathbf{G}(\mathbb{Q}) / \mathbf{P}_\Theta(\mathbb{Q})$ are *finite* for all Θ (see [B1], §15.6). Let $\Delta = [e_1, \dots, e_q] \subset \mathbb{R}^q$ denote a

standard geometric $q-1$ simplex ($q = \mathbb{Q}$ -rank of \mathbf{G}). If $\Delta = \{\alpha_1, \dots, \alpha_q\}$ and $\Delta - \Theta = \{\alpha_{i_1}, \dots, \alpha_{i_s}\}$ with $1 \leq i_1 < \dots < i_s \leq q$, we define the boundary simplex $\Delta(\Theta)$ of Δ as $\Delta(\Theta) := [e_{i_1}, \dots, e_{i_s}]$. Let \mathbf{P} be a *minimal* parabolic \mathbb{Q} -subgroup of \mathbf{G} and let the set $\Gamma \backslash \mathbf{G}(\mathbb{Q}) / \mathbf{P}(\mathbb{Q})$ be represented by $\{q_1, \dots, q_m\}$ (see Proposition 2.1). We take m copies $\Delta^j = [e_1^j, \dots, e_q^j]$ of Δ with faces $\Delta^j(\Theta)$ corresponding to Θ . The corresponding homeomorphisms $\Delta \simeq \Delta^j$ are denoted by φ_j . The simplicial complex $\Gamma \backslash |\mathcal{T}|$, which provides a geometric realization of the quotient of the Tits building of \mathbf{G} modulo Γ , is constructed from the simplices $\Delta^1, \dots, \Delta^m$ through the following *incidence relations*:

Two simplices Δ^j and Δ^l are pasted together along the faces $\Delta^j(\Theta)$ and $\Delta^l(\Theta)$ by the homeomorphism $\varphi_j \circ \varphi_l^{-1} |_{\Delta^l(\Theta)}$ if and only if

$$\Gamma q_j \mathbf{P}_\Theta(\mathbb{Q}) = \Gamma q_l \mathbf{P}_\Theta(\mathbb{Q}).$$

We remark that the points of $\Gamma \backslash |\mathcal{T}|$ are in one-to-one correspondence with equivalence classes of geodesic rays in the locally symmetric space $V = \Gamma \backslash X$ (see [Hat], [L1] and [JM]).

2.2. AN EXHAUSTION BY POLYHEDRA

We index the “edges” of the Weyl chamber $\overline{\mathfrak{a}^+}$ (or equivalently of $\overline{A^+} \cdot x_0$) by *simple* \mathbb{Q} -roots. More precisely, the edges of $\overline{A^+} \cdot x_0$ are given by geodesic rays $c_\alpha(t) = \exp(tH_\alpha) \cdot x_0$ where $H_\alpha \in \overline{\mathfrak{a}^+}$, $\|H_\alpha\| = 1$ and $\beta(H_\alpha) = 0$ for $\beta \neq \alpha$ ($\alpha, \beta \in \Delta$). We further write $c_{k\alpha}$ for the edges $q_k a_0 c_\alpha$ of the chambers $q_k \mathcal{C}$ in the fundamental set Ω (see Section 2.1 for the notation). If a geodesic ray c represents a point $z \in \partial_\infty X$ we write $z = c(\infty)$. The group G act naturally on $\partial_\infty X$ through $g \cdot c(\infty) = (g \cdot c)(\infty)$. For every $\alpha \in \Delta$ the isotropy group of $c_\alpha(\infty)$ under that action coincides with the (maximal) parabolic subgroups $P_{\Delta - \{\alpha\}}$ introduced above (see [L2] Lemma 1.2).

To a geodesic ray $c: [0, \infty) \rightarrow X$ (parametrized by arc-length) which represents a point z in the ideal boundary $\partial_\infty X$ of X is associated a *Busemann function on X at z* given by

$$h_z: X \rightarrow \mathbb{R} \quad ; \quad h_z(x) = \lim_{t \rightarrow \infty} [d(x, c(t)) - t].$$

The level sets of a Busemann function are *horospheres*, which foliate the symmetric space. We denote the Busemann functions which correspond to the rays $c_{k\alpha}$ by $h_{k\alpha}$. Note that $h_{k\alpha}(c_{k\alpha}(t))$ tends to $-\infty$ if the arc-length t of the geodesic $c_{k\alpha}$ tends to $+\infty$.

In contrast to an exact fundamental domain there are not only points on the boundary of a fundamental set Ω but possibly also interior points which are

identified under the action of Γ . However, there is only a finite set of isometries $\gamma \in \Gamma$ with $\gamma\Omega \cap \Omega \neq \emptyset$. Furthermore it suffices to look at the (finite) set \mathcal{D} of those γ for which this intersection is not relatively compact in X (all other intersections are contained in some compact subset of Ω). It turns out that every $\gamma \in \mathcal{D}$ has the crucial property that there are indices i, j such that $q_j^{-1}\gamma q_i$ is parabolic i.e. fixes at least one point in the ideal boundary $\partial_\infty X$ (see [L2] Proposition 2.2). Then for every $\gamma \in \mathcal{D}$ there are indices i, j, α such the family of horospheres of the form $h_{i\alpha}^{-1}(s), s \in \mathbb{R}$, is mapped isometrically to the family $h_{j\alpha}^{-1}(s), s \in \mathbb{R}$ (see [L2] Lemma 3.2). These identifications correspond to the incidence relations described above in the construction of the simplicial complex $\Gamma \backslash |\mathcal{T}|$. (To see this one has to use the fact that the Siegel set at infinity $\partial_\infty(q_j\mathcal{S})$ is canonically isomorphic to $\Delta^j = [e_1^j, \dots, e_q^j]$.) The main technical step is then to renormalize the Busemann functions as $\tilde{h}_{i\alpha} = h_{i\alpha} - s_{ij}$ (for certain constants s_{ij}) in such a way that each $\gamma \in \mathcal{D}$ maps a horosphere of some given level, say $\{\tilde{h}_{i\alpha} = s\}$, to another one, $\{\tilde{h}_{j\alpha} = s\}$, of the *same* level s (see [L2] Lemma 3.4). This fact finally allows us to truncate the constituents $q_i\mathcal{S}$ of the fundamental set Ω by removing the open horoballs $\mathcal{B}_{i\alpha}(s) := \{\tilde{h}_{i\alpha} < -\tau_\alpha s\}$ (for certain constants τ_α and for $s > 0$ sufficiently large). The above construction guarantees that the truncated fundamental set $\Omega(s) := \bigcup_{i=1}^m q_i\mathcal{S}(s)$ of Ω is relatively compact in X and invariant under the (restricted) action of Γ . Moreover for s sufficiently large the Γ -invariant “core” $X(s) := \Gamma \cdot \Omega(s)$ can be written as the complement in X of a union of (countably many) open horoballs: $X(s) = X - \Gamma \cdot \bigcup_{i=1}^m \bigcup_{\alpha \in \Delta} \mathcal{B}_{i\alpha}(s)$ (see [L3] Theorem 3.6). These horoballs are disjoint if and only if Γ is an arithmetic subgroup of a \mathbb{Q} -rank 1 group. The projection $\pi : X \rightarrow V$ maps $X(s)$ to a compact submanifold with corners $V(s)$ of V whose fundamental group is isomorphic to Γ . The “centers” of the projected horoballs in $\partial_\infty V$ are in bijection with the vertices of $\Gamma \backslash |\mathcal{T}|$. The exhaustion function h is eventually defined in such a way that its level sets coincide with the boundaries $\partial V(s)$. We summarize the result in the following proposition (see [L2] Theorem 4.2).

PROPOSITION 2.2. *Let X be a Riemannian symmetric space of noncompact type and \mathbb{R} -rank ≥ 2 and let Γ be an irreducible, torsion-free, non-uniform lattice in the group of isometries of X . On the locally symmetric space $V = \Gamma \backslash X$ there exists a piecewise real analytic exhaustion function $h : V \rightarrow [0, \infty)$ such that, for each $s \geq 0$, the sublevel set $V(s) := \{h \leq s\}$ is a Riemannian polyhedron in V . Moreover the level sets $\{h = s\} = \partial V(s)$ consist of projections of pieces of horospheres in X .*

Each polyhedron $V(s)$ is homotopically equivalent to V . More precisely we have

PROPOSITION 2.3. *For every sufficiently large s the locally symmetric space V is homeomorphic to the interior of the polyhedron $V(s)$ in V , and $V(s)$ is a strong deformation retract of V .*

For the proof see [L3], Theorems 5.2 and 5.5.

3. ESTIMATES FOR THE BOUNDARY SUBPOLYHEDRA

We wish to apply Proposition 1.1 to the polyhedra $V(s)$ in the above exhaustion and then take the limit for $s \rightarrow \infty$. To that end we need estimates for the second fundamental forms and the volumes of the (lower dimensional) boundary polyhedra.

For each Siegel set $\mathcal{S}_i := q_i \mathcal{S}$ which is part of the fundamental set Ω we have its truncated part

$$\mathcal{S}_i(s) := \mathcal{S}_i - \bigcup_{\alpha \in \Delta} (\mathcal{B}_{i\alpha}(s) \cap \mathcal{S}_i).$$

The top dimensional boundary faces of $\mathcal{S}_i(s)$ in \mathcal{S}_i (resp. of $\Omega(s)$ in Ω) are subsets of horospheres :

$$\mathcal{H}_{i\alpha}(s) := \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\} \cap \mathcal{S}_i(s), \quad \alpha \in \Delta.$$

The ‘‘horospherical’’ pieces $\mathcal{H}_{i\alpha}(s)$ together with their Γ -translates form the boundary of the manifold with corners $X(s)$ in X . For any nonempty subset Θ of Δ we set

$$\mathcal{H}_{i\Theta}(s) := \bigcap_{\alpha \in \Theta} \mathcal{H}_{i\alpha}(s) \subset \mathcal{S}_i(s).$$

The various boundary subpolyhedra of $V(s)$ are then unions of projections of the pieces $\mathcal{H}_{i\Theta}(s)$ under the canonical projection $\pi : X \rightarrow V$. More precisely, as explained in Section 2, for any subset $\Theta \subset \Delta$, we have the equivalence relation on the set $I = \{1, \dots, m\}$

$$j \sim_\Theta l \text{ if and only if } \Gamma q_j P_\Theta = \Gamma q_l P_\Theta$$

(the q_i are as in Proposition 2.1). This relation \sim_Θ induces a partition, $I(\Theta)$, of the set I whose components will be denoted by E . Let $n = \dim X = \dim V$, let k be the cardinality of Θ and let $E \in I(\Theta)$. Then $V_E^{n-k}(s) := \pi(\bigcup_{i \in E} \mathcal{H}_{i\Theta}(s))$ is a $(n - k)$ -dimensional boundary polyhedron of $V(s)$; and moreover, any boundary polyhedron arises in this way (see [L3] §4).

REMARK. The minimal possible dimension which occurs is $n - q$ where q is the \mathbb{Q} -rank of \mathbf{G} . It is also interesting to note (though not needed below) that the outer angles are isomorphic to \mathbb{Q} -Weyl chambers and their walls at infinity.

We shall use the following well-known fact about Jacobi fields in symmetric spaces (see [K] Theorem 2.2.9). A Jacobi field along a geodesic ray is called *stable* if its length is bounded.

LEMMA 3.1. *Let $r : [0, \infty) \rightarrow X$ be a unit-speed geodesic ray in the symmetric space X (of noncompact type). Set $p = r(0)$. Then the unique stable Jacobi field $J_u(s)$ along $r(s)$ with $J_u(0) = u \in T_p X$ can be written as*

$$J_u(s) = \sum_j e^{-\lambda_j s} a_j E_j(s)$$

where $\{E_j(s)\}$ is an orthonormal frame of parallel fields along r , the λ_j are non-negative (uniform) constants and $u = \sum_j a_j E_j(0)$.

LEMMA 3.2. *Let $s \geq 0$. The second fundamental forms of every boundary polyhedron $V_E^{n-k}(s)$ with respect to outer angles in $V(s)$ are uniformly bounded by a constant independent of E, k and s .*

Proof. Since the claim is local we can work in the universal covering space X . As we noted above the preimage of $V_E^{n-k}(s)$ in X under the projection π is the union of a finite number of horospherical sets

$$\mathcal{H}_{i\Theta}(s) = \bigcap_{\alpha \in \Theta} \mathcal{H}_{i\alpha}(s) \subset \bigcap_{\alpha \in \Theta} \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\},$$

where Θ is a subset of Δ with k elements. The (inner) unit normal field of the horosphere $\{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\}$ is given by $Z_{i\alpha} := -\text{grad } \tilde{h}_{i\alpha}$ (see e.g. [HI] Proposition 3.1). Using $d\pi$ any element in the outer angle $O(\pi(p))$ of $V_E^{n-k}(s)$ at a point $\pi(p) \in V_E^{n-k}(s)$ can then be identified with a positive linear combination (of norm 1) of the radial fields $Z_{i\alpha}(p)$, $\alpha \in \Theta$. It therefore suffices to show that for any pair (i, α) the second fundamental form of $V_E^{n-k}(s)$ relative to $d\pi Z_{i\alpha}$ is uniformly bounded. We fix i and α and write Z for $Z_{i\alpha}$. For $p \in X$ let $\langle \cdot, \cdot \rangle_p$ denote the Riemannian metric of X at p . Let $u, v \in T_p X$ be such that $d\pi(u), d\pi(v) \in T_{\pi(p)} V_E^{n-k}(s)$. Using the above identifications the second fundamental form of $V_E^{n-k}(s) \subset V(s)$ with respect to Z can be written as

$$\Pi_Z(u, v)(p) = \langle D_u Z, v \rangle_p.$$

According to [HI], Proposition 3.1, we have $D_u Z(p) = J'_u(0)$ where J_u is the stable Jacobi field along the (unique) geodesic ray, say r , in X which joins p to $c_{i\alpha}(\infty) \in \partial_\infty X$ and with initial value $J_u(0) = u$. By Lemma 3.1 there are orthonormal parallel fields $E_j(s)$ along r and constants $\lambda_j \geq 0$ such that $J_u(s) = \sum_j e^{-\lambda_j s} a_j E_j(s)$ with $u = \sum_j a_j E_j(0)$. Consequently we get $J'_u(0) = -\sum_j \lambda_j a_j E_j(0)$ and finally, for $v = \sum_j b_j E_j(0)$, $|\Pi_Z(u, v)(p)| = |-\sum_j \lambda_j a_j b_j| \prec \|u\| \|v\|$. \square

We next estimate the volumes of the boundary polyhedra. Recall from Section 2.1 the parametrization of X by horocyclic coordinates

$$\mu: Y = U \times Z \times A \longmapsto X; (u, \tau(m), a) \longmapsto uma \cdot x_0.$$

Let dx^2 be the G -invariant Riemannian metric on X induced by the Cartan-Killing form of the Lie algebra \mathfrak{g} of G and let dz^2 be the invariant metric on Z . Further let da^2 (resp. du^2) be the left-invariant metric on A (resp. U). Finally set $dy^2 := \mu^* dx^2$.

LEMMA 3.3. *Let dv_Y, dv_U, dv_Z and dv_A denote the volume elements of the metrics dy^2, du^2, dz^2 and da^2 . Then at the point $(u, z, a) \in Y$ we have*

$$2^e dv_Y = \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_A$$

where $e = \frac{1}{2} \dim U$ and ρ is the sum of all positive roots (counted with multiplicity); it can be written in the form $\rho = \sum_{\alpha \in \Delta} c_\alpha \alpha, c_\alpha > 0$.

For the proof see [B2] Corollary 4.4.

LEMMA 3.4. *For the $(n - k)$ -dimensional volume of each boundary polyhedron $V_E^{n-k}(s)$ of $V(s)$ one has the estimate*

$$\text{Vol}(V_E^{n-k}(s)) \prec s^{q-k} e^{-cs},$$

where $q = \dim A$ is the \mathbb{Q} -rank of \mathbf{G} and $c > 0$ is a constant (independent of E, k and s).

Proof. We again consider the preimage of $V_E^{n-k}(s)$ in X under the map π . We need to estimate the volume of each horospherical piece

$$\mathcal{H}_{i\Theta}(s) = \bigcap_{\alpha \in \Theta} \{\tau_\alpha^{-1} \tilde{h}_{i\alpha} = -s\} \cap \mathcal{S}_i(s), \quad i \in E.$$

It clearly suffices to carry out the estimates for $i = 1$; note that $q_1 = e$. For the horocyclic coordinate map $\mu: Y \rightarrow X$ and the canonical projection

$\pi_A : Y \rightarrow A$ we set $A_\Theta(s) := \pi_A \circ \mu^{-1}(\mathcal{H}_{1\Theta}(s)) \subset A$. The set $A_\Theta(s)$ is contained in an "affine" subspace of A of the form $a_1 a_*(s) A^{q-k}$ where $a_1 a_*(s) \in A$ and A^{q-k} is a $q-k$ -dimensional subgroup of A (see Sections 3 and 4 of [L2]). We denote the restriction of dv_A to A^{q-k} by $dv_{A^{q-k}}$; for $k = q$ we have $A^0 = e$ and we set $dv_{A^0} \equiv 1$. By Lemma 3.3 we have (for k equal to the number of elements of Θ)

$$\text{Vol}(V_E^{n-k}(s)) \prec \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}}.$$

Since the horospherical piece $\mathcal{H}_{1\Theta}(s)$ is part of a Siegel set $\mathcal{S}_{\omega, \tau}$ with ω relatively compact (and hence of finite volume) in UM we get

$$\begin{aligned} \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}} &\prec \\ &\prec \int_{\omega} dv_U \wedge dv_Z \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}} \prec \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}}. \end{aligned}$$

Also by definition of a Siegel set we have $\alpha(a) \geq \tau \succ 1$ for all $\alpha \in \Delta$. Moreover, the computations in the proof of Lemma 4.1 (and Lemma 3.5) in [L2] show that for all $\alpha \in \Theta$ one has $\alpha(a_1 a_*(s)) \succ e^{\mu_\alpha s}$ with $\mu_\alpha > 0$. Hence, as $\Theta \subset \Delta$ is not empty and since $\rho = \sum_{\alpha \in \Delta} c_\alpha \alpha$ ($c_\alpha > 0$), there is a uniform constant $c > 0$ such that $\rho(a)^{-1} \prec e^{-cs}$ for all $a \in A_\Theta(s)$. As noted above the set $A_\Theta(s)$ is contained in a $(q-k)$ -dimensional affine cone in A . It is similar (in the sense of Euclidean geometry) to $A_\Theta(0)$ with similarity factor s (see the proof of Lemma 4.1 in [L2]). Hence we eventually get $\int_{A_\Theta(s)} dv_{A^{q-k}} \prec s^{q-k}$ and the Lemma follows. \square

4. A NEW PROOF OF THE GAUSS-BONNET FORMULA

In this section we present a new simplified proof of the Gauss-Bonnet theorem for higher rank locally symmetric spaces.

THEOREM 4.1. *Let X be a Riemannian symmetric space of noncompact type and \mathbb{R} -rank ≥ 2 and let Γ be an irreducible, torsion-free (non-uniform) lattice in the group of isometries of X . Then for the locally symmetric space $V = \Gamma \backslash X$ the Gauss-Bonnet formula holds:*

$$\chi(V) = \int_V \Psi dv.$$

Proof. By Proposition 2.2 there is an exhaustion $V = \bigcup_{s \geq 0} V(s)$ of V by Riemannian polyhedra $V(s)$. Each polyhedron $V(s)$ in this exhaustion is equipped with the Riemannian metric induced by the one of V . Proposition 1.1 applied to $V(s)$ yields

$$\left| (-1)^n \chi'(V(s)) - \int_{V(s)} \Psi dv \right| \prec \sum_{k=1}^q \sum_E \int_{V_E^{n-k}(s)} \int_{O(p)} \|\Psi_{E,k}\| d\omega_{k-1} dv_E(p)$$

where $q = \dim A$ is the \mathbb{Q} -rank of \mathbf{G} (see Section 2.1) and where the index E runs through a finite set. As we remarked in Section 1 the function $\Psi_{E,k}$ is locally computable from the components of the metric and the curvature tensor of $V(s)$ and from the components of the second fundamental form of $V_E^{n-k}(s)$ in $V(s)$. The fact that V is locally symmetric together with Lemma 3.2 thus implies that $\|\Psi_{E,k}\| \prec 1$ for all E, k . Using Lemma 3.4 we conclude that

$$\left| (-1)^n \chi'(V(s)) - \int_{V(s)} \Psi dv \right| \prec \sum_{k,E} \text{Vol}(V_E^{n-k}(s)) \prec e^{-cs} \sum_{k=1}^q s^{q-k}.$$

By Proposition 2.3 we have $\chi'(V(s)) = \chi(V)$. The polyhedra $V(s)$ exhaust V and $\chi(V)$ is an integer; hence $(-1)^n \chi(V) = \int_{V(s)} \Psi dv$ for sufficiently large s . Finally, for n odd $\Psi \equiv 0$ by definition (see [AW]) and the claimed formula follows. \square

REFERENCES

- [AW] ALLENDOERFER, C. B. and A. WEIL. The Gauss-Bonnet theorem for Riemannian polyhedra. *Trans. Amer. Math. Soc.* 53 (1943), 101–129.
- [BGS] BALLMANN, W., M. GROMOV and V. SCHROEDER. *Manifolds of Nonpositive Curvature*. Boston, 1985.
- [B1] BOREL, A. *Introduction aux groupes arithmétiques*. Paris, 1969.
- [B2] ——— Stable and real cohomology of arithmetic groups. *Ann. scient. Éc. Norm. Sup.* 4^e série 7 (1974), 235–272.
- [B3] ——— *Linear Algebraic Groups*. Second edition, New York, 1991.
- [BS] BOREL, A. and J.-P. SERRE. Corners and arithmetic groups. *Comment. Math. Helv.* 48 (1973), 436–491.
- [BT] BOREL, A. and J. TITS. Groupes réductifs. *I.H.E.S. Publ. Math.* 27 (1965), 55–150.
- [C] CHERN, S. S. A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds. *Ann. of Math.* 45 (1944), 747–752.
- [CG1] CHEEGER, J. and M. GROMOV. On the characteristic numbers of complete manifolds of bounded curvature and finite volume. *Differential Geometry and Complex Analysis*. H. E. Rauch Memorial Volume (I. Chavel, H. M. Farkas, eds.), Berlin, 1985, 115–154.

- [CG2] CHEEGER, J. and M. GROMOV. Bounds on the von Neumann dimension of L_2 -cohomology and the Gauss-Bonnet theorem for open manifolds. *J. Diff. Geom.* 21 (1985), 1–39.
- [CG3] CHEEGER, J. and M. GROMOV. Chopping Riemannian manifolds. A symposium in honour of M. do Carmo (B. Lawson, K. Tenenblat, eds.). *Pitman Monographs and Surveys in Pure and Applied Mathematics* 52, New York, 1991, 85–94.
- [DH] DOUADY, A. et L. HÉRAULT. Arrondissement des variétés à coins. Appendix of [BS], 484–489.
- [G] GROMOV, M. Volume and bounded cohomology. *Publ. Math. IHES* 56 (1982), 5–100.
- [H] HARDER, G. A Gauss-Bonnet formula for discrete arithmetically defined groups. *Ann. scient. Éc. Norm. Sup. 4^e série* 4 (1971), 409–455.
- [Hat] HATTORI, T. Collapsing of quotient spaces of $SO(n)\backslash SL(n, \mathbb{R})$ at infinity. Preprint, Tokyo, 1991.
- [HI] HEINTZE, C. and H.-C. IM HOF. Geometry of horospheres. *J. Diff. Geom.* 12 (1977), 481–491.
- [JM] JI, L. and R. MACPHERSON. Geometry of compactifications of locally symmetric spaces. Preprint, Cambridge MA, 1993.
- [K] KLINGENBERG, W. *Riemannian Geometry*. Berlin, 1982.
- [L1] LEUZINGER, E. Geodesic rays in locally symmetric spaces. *Diff. Geom. & its Appl.* (to appear).
- [L2] — An exhaustion of locally symmetric spaces by compact submanifolds with corners. *Inventiones math.* 121 (1995), 389–410.
- [L3] — Polyhedral retracts and compactifications of locally symmetric spaces. Preprint, Zürich, 1994.
- [R1] RAGHUNATHAN, M.S. A note on quotients of real algebraic groups by arithmetic subgroups. *Invent. math.* 4 (1968), 318–335.
- [R2] — *Discrete subgroups of Lie groups*. New York, 1972.
- [Z] ZIMMER, R.J. *Ergodic Theory and Semisimple Groups*. Boston, 1984.

(Reçu le 21 avril 1995)

Enrico Leuzinger

Institut für Mathematik
Universität Zürich
Winterthurer-Str. 190
CH-8057 Zürich
Switzerland

E-mail address : leuzinge@math.unizh.ch