

# 1. A BASIC CONSTRUCTION

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## 4-MANIFOLDS, GROUP INVARIANTS, AND $l_2$ -BETTI NUMBERS

by Beno ECKMANN

It has been known for some time that closed 4-manifolds provide, via the fundamental group and the Euler characteristic, interesting invariants for finitely presented groups. In this short survey we describe these and more refined invariants (using also the signature of the manifold), and explain some of their significance. The invariants are not easily calculated in general, but quite good information is obtained using  $l_2$ -Betti numbers.

The topic has been developed by several authors, more or less independently. We mention Hausmann-Weinberger [H-W], Kotschick [K], Lück [L], and myself [E1], [E2]. The paper [K] contains a wealth of information on the invariants and further important references; the application of  $l_2$ -Betti numbers appears in [E2] and in [L].

### 1. A BASIC CONSTRUCTION

1.1. We will always denote by  $M$  a connected orientable closed 4-manifold (compact without boundary) admitting a cell decomposition. The fundamental group  $G = \pi_1(M)$  is finitely presented. Indeed, homotopy classes of loops can be represented by edge-polygons and null-homotopies of these by using 2-cells. Conversely, any finitely presented group  $G$  is the fundamental group of a closed 4-manifold. If

$$G = \langle g_1, \dots, g_m \mid r_1, \dots, r_n \rangle$$

is a presentation of  $G$ , there is a standard procedure for constructing such a manifold: One first puts  $M' = S^1 \times S^3 + \dots + S^1 \times S^3$ , connected sum, one copy for each generator  $g_i$  of  $G$ . Then  $\pi_1(M')$  is a free group on generators  $g_1, \dots, g_m$ . A relator, say  $r_1$ , is a word in the  $g_i$  and can be represented by a loop  $S^1$  in  $M'$ .

A tubular neighbourhood  $S^1 \times B^3$  of  $S^1$ , where  $B^k$  is the  $k$ -dimensional ball, has boundary  $S^1 \times S^2$ . Replacing the interior by  $B^2 \times S^2$  with the same

boundary yields a new 4-manifold where the element corresponding to  $r_1$  has been killed; and similarly for the other  $r_i$ . Let  $M_0$  be the 4-manifold thus obtained, fulfilling  $\pi_1(M_0) = G$ . The idea of that construction can already be found in the old book [S-T]. Much later the procedure, in a more general context, has been called "elementary surgery".

1.2. We recall that the (good old) Euler characteristic  $\chi(X)$  of a finite cell complex  $X$  is the alternating sum

$$\chi(X) = \sum (-1)^i \alpha_i,$$

where  $\alpha_i$  is the number of  $i$ -cells. It is easily computed for  $M_0$  above: For  $M'$  it is  $2 - 2m$  since it is  $= 0$  for  $S^1 \times S^3$  and since it decreases by 2 in a connected sum. Under the surgery process above it increases by 2 [use the fact that for the union of two complexes  $X$  and  $Y$  with intersection  $Z$  the characteristic is  $\chi(X) + \chi(Y) - \chi(Z)$ ; and that  $\chi(B^2 \times S^2) = 2$ ]. Whence

$$\chi(M_0) = 2 - 2m + 2n = 2 - 2(m - n).$$

The difference  $m - n$  is called the deficiency of the presentation of  $G$ .

1.3. On the other hand the characteristic can be expressed by the Betti numbers of the cell complex  $X$  as  $\sum (-1)^i \beta_i(X)$  where  $\beta_i(X) = \dim_{\mathbf{R}} H_i(X; \mathbf{R})$  (and is therefore a topological invariant). Moreover the  $\beta_i$  of a manifold fulfill Poincaré duality, i.e. they are equal in complementary dimensions. Thus  $\chi(M) = 2 - 2\beta_1(M) + \beta_2(M)$ . We recall that homology in dimension 1 depends on the fundamental group  $G$  only;  $\beta_1$  is the  $\mathbf{Q}$ -rank of  $G$  Abelianised and we write  $\beta_1(G)$  for  $\beta_1(M)$ . Comparing with  $\chi(M_0)$  above we see that the deficiency of the presentation is  $\leq \beta_1(G)$ . Thus there is a maximum for the deficiency of all presentations of  $G$ , called the deficiency  $\text{def}(G)$  of  $G$ . [For this simple side result there are, of course, much easier arguments.]

## 2. THE HAUSMANN-WEINBERGER INVARIANT

2.1. As seen above, the Euler characteristic of a 4-manifold  $M$  with given finitely presented fundamental group  $G$  is bounded below by  $2 - 2\beta_1(G)$ . The minimum of  $\chi(M)$  for all such  $M$  has been considered by Hausmann-Weinberger [H-W] and denoted by  $q(G)$ . Using  $M_0$  above we have the inequalities

$$2 - 2\beta_1(G) \leq q(G) \leq 2 - 2 \text{def}(G).$$