

0. Introduction

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

AMENABILITY AND GROWTH OF ONE-RELATOR GROUPS

by Tullio G. CECCHERINI-SILBERSTEIN and Rostislav I. GRIGORCHUK

ABSTRACT. An algorithm showing whether a group given by a one-relator presentation is amenable or not is constructed. Sufficient conditions for a one-relator group of exponential growth to have uniformly exponential growth are also given.

0. INTRODUCTION

A one-relator group is a group G which admits a presentation

$$(*) \quad G = \langle a_1, a_2, \dots, a_m : R(a_1, a_2, \dots, a_m) = 1 \rangle$$

with one defining relation.

The paper by G. Baumslag [B 1] is a comprehensive survey of results about one-relator groups. In particular this paper stresses the role of algorithmic problems in the theory of one-relator groups.

Recently the interest in functional-analytical and asymptotical properties of one-relator groups has increased. For instance, the entropy of one-relator groups was discussed in [GrLP], random walks and Markov operators on one-relator groups were investigated in [CV], [BCCH], [BC], and the K-functor of reduced C^* -algebras of one-relator groups was computed in [BBV]. Also the growth functions of the groups $\Gamma_n = \langle t, a : tat^{-1} = a^n \rangle$, $n \neq 0, \pm 1$, and of some other one-relator groups were calculated in [CEG] and [EJ].

Recall that a discrete group G is amenable if there exists a finitely additive measure $\mu: \mathcal{P}(G) = \{0, 1\}^G \longrightarrow [0, 1]$ which is G -(left)-invariant ($\mu(gE) = \mu(E)$ for all $g \in G$ and $E \subset G$) and such that, in addition, $\mu(G) = 1$. For our purpose it will be enough to know that a group containing a free subgroup of rank two is not amenable, and that, on the contrary, any solvable group is amenable ([G]).

As easily follows from the paper of Karrass and Solitar [KS], all amenable one-relator groups are in the following list:

- (**)
1. $\langle a : a^n = 1 \rangle \cong \mathbf{Z}_n$, cyclic groups of finite order $n = 1, 2, \dots$;
 2. $\langle a, b : b = 1 \rangle \cong \mathbf{Z}$, the infinite cyclic group;
 3. $\langle a, b : bab^{-1} = a^n \rangle$, $n \neq 0$.
 This class splits into two subclasses:
 - 3_a. $n = +1$: $\langle a, b : bab^{-1} = a \rangle \cong \mathbf{Z}^2$;
 - $n = -1$: $\langle a, b : bab^{-1} = a^{-1} \rangle$:
 this group contains a subgroup $\cong \mathbf{Z}^2$ of index two,
 but it is not $\cong \mathbf{Z}^2$;
 - 3_b. $n \neq 0, \pm 1$: $\langle a, b : bab^{-1} = a^n \rangle$:
 these groups are 2 step-solvable and of exponential
 growth (pairwise non-isomorphic).

Also Tits' alternative does hold for one-relator groups: any one-relator group either contains a free subgroup of rank two or is solvable (and from the above list).

But in the Karrass-Solitar paper no algorithm is given answering the question whether, given a one-relator presentation, the corresponding group is solvable or not. In Section 1 we present a simple algorithm and, as a consequence, we re-obtain the above list of all amenable one-relator groups.

In the second part of the paper we investigate the uniformly exponential growth for one-relator groups of exponential growth.

Recall that if G is a group with a finite generating system A ,

$$|g|_A = \min \{n : g = a_1 a_2 \cdots a_n, a_i \in A\}$$

is the *length* of an element $g \in G$ with respect to A and $\gamma_A^G(n) = |\{g \in G : |g|_A \leq n\}|$ is the *growth function* of G with respect to the generating system A . The limit

$$\lambda_A(G) = \lim_{n \rightarrow \infty} \sqrt[n]{\gamma_A^G(n)}$$

exists and $\lambda_A(G) \geq 1$. The group G is said to have *exponential growth* (respectively *sub-exponential growth*) if $\lambda_A(G) > 1$ (resp. $\lambda_A(G) = 1$) for some (and therefore for any other) finite system of generators A .

Denoting now by

$$\lambda_*(G) = \inf_A \lambda_A(G)$$

the *minimal growth rate* of G , where the infimum is taken over all finite generating systems, the group G has *uniform exponential growth* if $\lambda_*(G) > 1$. This last concept is due to Avez [A] where the number

$$h(G) = \log(\lambda_*(G))$$

is called the *entropy* of the group G and it is discussed in [GrLP], [SW] and in the survey paper [GH].

The simplest example of a group with uniformly exponential growth is the free group \mathbf{F}_m of finite rank $m \geq 2$ for which the minimal growth rate is $\lambda_*(\mathbf{F}_m) = 2m - 1$, see for instance [GH].

It is not known whether a group of exponential growth has necessarily uniformly exponential growth or not. We formulate the following:

0.1. CONJECTURE. *All one-relator groups of exponential growth have uniformly exponential growth.*

Conjecture 0.1 is true for one-relator groups of rank $m \geq 3$ and for one-relator groups with torsion, therefore we focus our attention on two-generated one-relator groups and give sufficient conditions for such groups to have uniformly exponential growth. We present a new method for estimating the minimal growth rate of a finitely generated group using growth functions of the corresponding graded Lie algebra and apply it to one-relator groups.

1. AN ALGORITHM FOR CHECKING AMENABILITY

Let G be a one-relator group with presentation $(*)$; the number m of the generators of G in the presentation is called the rank of the presentation. Until Section 4 we shall assume that R is cyclically reduced and non trivial.

The next observation is well known. We shall include the proof stressing the algorithmic aspect of the statement.

1.1. LEMMA. *Let $G = \langle a, b, \dots : R(a, b, \dots) \rangle$ be a one-relator group with at least two generators. Then G has a presentation $\langle t, \dots : R'(t, \dots) \rangle$ with $\sigma_t(R') = 0$, where $\sigma_t(R')$ denotes the sum of the exponents of t in the word R' . This second presentation can in fact be produced, starting from the original one, in an algorithmical way.*