

2. ATKIN-LEHNER ACTION ON THETA FUNCTIONS

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

EXAMPLE 4. Similarly, $K = \mathbf{Q}\left(e\left(\frac{1}{8}\right), \sqrt{\pm q}\right)$ may be used to obtain strongly modular lattices Λ in $G_{8r}(2^+q^+)$ from rank r unimodular hermitian lattices over \mathcal{O}_K , setting now

$$x \cdot y = \frac{1}{2} \operatorname{Tr}_{K/\mathbf{Q}}(h(x, y)/(2 - \sqrt{2})).$$

Again, we always have $\min \Lambda \geq 4$. E.g., \mathcal{O}_K itself gives the tensor product of D_4 and the binary lattice with Gram matrix $\begin{pmatrix} 2 & 1 \\ 1 & (q+1)/2 \end{pmatrix}$.

2. ATKIN-LEHNER ACTION ON THETA FUNCTIONS

The subject treated in this section is not new, but appears to be difficult to cite from the literature (in the form we need it). For convenience, I give a rather detailed account, starting from a classical formula (due to Jacobi and others). Let Λ be an even lattice. The theta function of a coset $\bar{v} = v + \Lambda$ in Λ^* (and, in particular, Θ_Λ for $v = 0$) is that function defined on the upper half-plane by

$$\Theta_{\bar{v}}(z) = \sum_{x \in \bar{v}} e\left(\frac{1}{2}(x \cdot x)z\right).$$

Now let $n = 2k$ (k integral), and recall that $SL_2(\mathbf{R})$ acts on functions f on the upper half-plane by

$$(f |_k S)(z) = (cz + d)^{-k} f\left(\frac{az + b}{cz + d}\right), \quad S = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Let S be in $SL_2(\mathbf{Z})$, with $c > 0$. For $u, v \in \Lambda^*$ define

$$\phi_S(u, v) = \sum_x e((ax \cdot x + 2x \cdot v + dv \cdot v)/2c)$$

where x runs through a system of representatives of those elements of $\Lambda^*/c\Lambda$ which reduce to \bar{u} in $D = \Lambda^*/\Lambda$. Each summand clearly depends only on the class $x + c\Lambda$, and the whole sum depends only on \bar{u} and \bar{v} . The latter statement is trivial for \bar{u} , while for \bar{v} it is proved (using $1 = ad - bc$) by

$$\begin{aligned} \phi_S(u, v) &= \sum_x e(a(x + dv) \cdot (x + dv)/2c) e(-b(2x \cdot v + dv \cdot v)/2) \\ (2.1) \quad &= \phi_S(u + dv, 0) e(-b(2u \cdot v + dv \cdot v)/2). \end{aligned}$$

So we may write $\phi_S(u, v) = \phi_S(\bar{u}, \bar{v})$. Then the formula we need is (see [Mi], p. 189)

$$(2.2) \quad \Theta_{\bar{u}}|_k S = (\det \Lambda)^{-\frac{1}{2}} (ic)^{-k} \sum_{\bar{v} \in D} \phi_S(\bar{u}, \bar{v}) \Theta_{\bar{v}}.$$

Let Λ be of level ℓ . Then it is a well-known consequence of (2.2) that Θ_{Λ} is a modular form for the subgroup $\Gamma_0(\ell)$ of $SL_2(\mathbf{Z})$ defined by $c \equiv 0 \pmod{\ell}$. Now let $m|\ell$, $m' = \ell/m$. After [AL] the m -th Atkin-Lehner involution $\Gamma_0(\ell)W_m$, a coset of $\Gamma_0(\ell)$ in its normalizer in $SL_2(\mathbf{R})$, is given by any matrix of the form

$$W_m = S \begin{pmatrix} \sqrt{m} & 0 \\ 0 & 1/\sqrt{m} \end{pmatrix}, \quad S = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(m'), \quad d \equiv 0 \pmod{m}.$$

We specifically choose $a = 1$, $c = m'$, solve $tm + t'm' = 1$, and put $b = -t'$, $d = tm$. In any case,

$$W_m^2 \equiv 1, \quad W_m W_{m'} \equiv W_{\ell} \pmod{\Gamma_0(\ell)}.$$

Now these involutions are connected, via Gaussian sums, to those in Section 1 by the following important relation which seems to have been proved first by Kitaoka ([Ki] where, however, the constant factor is not worked out; cf. also [BS], p. 77, for the statement of a generalization).

ATKIN-LEHNER IDENTITY:

$$(2.3) \quad \Theta_{\Lambda}|_k W_m = \left(\frac{\det \Lambda_m}{\det \Lambda} \right)^{\frac{1}{4}} i^{-k} g_{m'}(\Lambda_{m'}) \Theta_{\Lambda_m}.$$

How this identity will be used here is readily explained. Suppose that Λ (and then also Λ_m) has determinant ℓ^k . We will be interested in the following gradually stronger conditions

- (1) $\Theta_{\Lambda} = \Theta_{\Lambda_m}$ for all $m|\ell$
- (2) Λ is strongly modular.

So condition (1) says that Θ_{Λ} is an eigenform of all the Atkin-Lehner involutions; Section 3 deals with such modular forms.

Proof of (2.3). Recall that here S is chosen such that $a = 1$, $c = m'$ and $m|d$. Write $\bar{v} \in D$ in the form $\bar{v} = \bar{w} + \bar{y}$ where $\bar{w} \in D(m)$, $\bar{y} \in D(m')$. Then $\phi_S(0, \bar{v}) = \phi_S(0, \bar{y})$ by (2.1), and (2.2) gives

$$\begin{aligned} (\det \Lambda)^{\frac{1}{2}} (im')^k \Theta_{\Lambda}|_k S &= \sum_{\bar{v} \in D} \phi_S(0, \bar{v}) \Theta_{\bar{v}} \\ &= \sum_{\bar{y} \in D(m')} \phi_S(0, \bar{y}) \sum_{\bar{w} \in D(m)} \Theta_{\bar{w} + \bar{y}}. \end{aligned}$$

We first must know that $\phi_S(0, \bar{y}) = 0$ for $\bar{y} \neq 0$. To see this, write $x \in \Lambda$ in the form $x = x_1 + m'x'$ with x_1 from some system of representatives of $\Lambda/\sqrt{m'}\Lambda_{m'}$ and $\sqrt{m'}x' \in \Lambda_{m'}$. Then

$$\phi_S(0, y) e(-dy \cdot y/2m') = \sum_{x_1} e((x_1 \cdot x_1 + 2x_1 \cdot y)/2m') \sum_{\bar{x}' \in D(m')} e(x' \cdot y).$$

But the sum over $D(m')$ vanishes for $\bar{y} \neq 0$ (since $D(m')$ is regular with respect to the discriminant form). It remains to determine $\phi_S(0, 0)$. Put $E = \Lambda_{m'}^*/\Lambda_{m'}$. Since $(1/\sqrt{m'})\Lambda/\Lambda_{m'}$ is $E(m')$, the last formula for $y = 0$ gives

$$\begin{aligned} \phi_S(0, 0) &= \#D(m') \sum_{\bar{v} \in E(m')} e\left(\frac{1}{2}v \cdot v\right) \\ &= \#D(m') (\#E(m'))^{\frac{1}{2}} g_{m'}(\Lambda_{m'}) \end{aligned}$$

where

$$\begin{aligned} \#D(m') (\#E(m'))^{\frac{1}{2}} &= (\Lambda_{m'} : \sqrt{m'}\Lambda) (\sqrt{m'}\Lambda : m'\Lambda_{m'})^{\frac{1}{2}} \\ &= (\#D(m'))^{\frac{1}{2}} (m')^k \\ &= (\#D(m))^{-\frac{1}{2}} (\det \Lambda)^{\frac{1}{2}} (m')^k \\ &= m^{-\frac{k}{2}} \left(\frac{\det \Lambda_m}{\det \Lambda}\right)^{\frac{1}{4}} (\det \Lambda)^{\frac{1}{2}} (m')^k \quad \square \end{aligned}$$

Note that, by Propositions 1 and 2, (2.3) in the case $\det \Lambda = \ell^k$ becomes

$$(2.4) \quad \Theta_\Lambda|_k W_m = g_m(\Lambda_{m'})^{-1} \Theta_{\Lambda_m}.$$

3. SOME USE OF MODULAR FORMS

Let $W(\ell)$ be the elementary abelian 2-group formed by the Atkin-Lehner involutions $w_m = \Gamma_0(\ell)W_m$ for all $m \parallel \ell$. Let k be even, and let $\mathcal{S}_k(\ell)$ denote the space of cusp forms of weight k on $\Gamma_0(\ell)$. Then $W(\ell)$ acts on this space. For a character χ of $W(\ell)$ we let $\mathcal{S}_k(\ell)_\chi$ denote the subspace on which $W(\ell)$ acts by χ . If Λ and M are lattices of dimension $2k$ and level ℓ belonging to the same genus, then $f = \Theta_\Lambda - \Theta_M$ is known to be in $\mathcal{S}_k(\ell)$, and when both lattices are strongly modular identity (2.4) implies that f is in $\mathcal{S}_k(\ell)_\chi$ for the character $\chi(w_m) = g_m(\Lambda)$. So we are interested in such spaces now.

Fortunately, the dimension of $\mathcal{S}_k(\ell)_\chi$ is known; I am indebted to N.-P. Skoruppa for pointing out the reference. Let $s(\ell)$ denote the number of prime