

Objektyp: **ReferenceList**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

a normal subgroup of H then $D(\gamma)$ is in this subgroup (note that there are finitely many such normal subgroups).

Since $D(\gamma)$ has infinite order (if γ is non-trivial), $\langle D(\gamma) \rangle$ has positive dimension so that it contains a non-trivial one-parameter group. Hence every non trivial semi-simple element in $D(\Gamma)$ yields a one-parameter group contained in H_0 . We now show that these one-parameter subgroups generate the connected component of the identity in H . Observe the following elementary fact: if a family of vectors spans the Lie algebra of a Lie group, then the one-parameter groups generated by these vectors generate the connected component of the identity. Therefore, we consider the linear span \mathfrak{E} in the Lie algebra \mathfrak{H} of H of the Lie algebras of all the subgroups $\langle D(\gamma) \rangle$ for γ semi-simple. It is enough to show that $\mathfrak{E} = \mathfrak{H}$. Note that \mathfrak{E} is certainly non-trivial since semi-simple elements are Zariski dense in H . Note also that \mathfrak{E} is invariant under the adjoint action of $D(\Gamma)$, hence under the adjoint action of H since $D(\Gamma)$ is Zariski dense in H . It follows that \mathfrak{E} coincides with the product of some of the simple factors of \mathfrak{H} . The only possibility is that $\mathfrak{E} = \mathfrak{H}$ since otherwise, all the semi-simple $D(\gamma)$ would have some power contained in the same product of some but not all of the simple factors of H (note that the algebraic Abelian group $\langle D(\gamma) \rangle$ has a finite number of connected components). This implies that all semi-simple elements of $D(\Gamma)$ are contained in some non trivial normal subgroup of H . This is not possible by the following argument. In the algebraic group H , there is a non-empty open Zariski set consisting of semi-simple elements which are not contained in any non-trivial normal subgroup of H . Since $D(\Gamma)$ is Zariski dense in H , it intersects non-trivially this open set.

It follows that H_0 contains the connected component of the identity of H . Therefore H_0 is a semi-simple Lie group of finite index in H . By Kushnirenko's theorem, we can analytically linearize $\phi(H)$ (one also uses Remark 2.1) and in particular $\Phi(\Gamma)$.

Theorem 10.4 is proved.

REFERENCES

- [1] ABRAHAM, R. and J. E. MARSDEN. *Foundations of Mechanics*. 2nd ed., Benjamin/Cummings, 1978.
- [2] D'AMBRA, G. and M. GROMOV. Lectures on transformation groups: geometry and dynamics. *Surveys in Differential Geometry, No. 1* (1991), 19–111. Supplement to the *Journal of Differential Geometry*.
- [3] BAKER, A. *Transcendental numbers*. Cambridge University Press, 1975.

- [4] BING, R. H. A homeomorphism between the 3-sphere and the sum of two solid horned spheres. *Ann. of Math.* 56 (1952), 354–362.
- [5] — Inequivalent families of periodic homeomorphisms of E^3 . *Ann. of Math.* 80 (1964), 78–93.
- [6] BREDON, G. E. *Topology and Geometry*. Springer-Verlag, 1993.
- [7] BRJUNO, A. D. Analytical form of differential equations, I and II. *Trans. Moscow Math. Soc.* 25 (1971), 131–288, and 26 (1972), 199–239.
- [8] DYNKIN, E. B. Maximal subgroups of semi-simple Lie algebras. *Amer. Math. Soc. Transl. (Series 2)* 6 (1957), 111–244.
- [9] EILENBERG, S. Sur les transformations périodiques de la surface de sphère. *Fund. Math.* 22 (1934), 28–41.
- [10] FLATO, M., G. PINCZON and J. SIMON. Non linear representations of Lie groups. *Ann. Sci. École Norm. Sup.* 10 (1977), 405–418.
- [11] GUILLEMIN, V. W. and S. STERNBERG. Remarks on a paper of Hermann. *Trans. Amer. Math. Soc.* 130 (1968), 110–116.
- [12] HARTMAN, P. *Ordinary differential equations*. 2nd ed., Birkhäuser, 1982.
- [13] HERMANN, R. The formal linearization of a semisimple Lie algebra of vector fields about a singular point. *Trans. Amer. Math. Soc.* 130 (1968), 105–109.
- [14] HILTON, P. J. and U. STAMMBACH. *A Course in Homological Algebra*. Springer-Verlag, 1971.
- [15] HSIANG, W.-C. A note on free differentiable actions of S^1 and S^3 on homotopy spheres. *Ann. of Math.* 83 (1966), 266–272.
- [16] JACOBSON, N. *Lie Algebras*. Dover, 1979.
- [17] KATOK, A. and B. HASSELBLATT. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, 1995.
- [18] KIRBY, R. C. and L. C. SIEBENMANN. *Foundational essays on topological manifolds, smoothings and triangulations*. Princeton University Press, 1977.
- [19] KOBAYASHI, S. *Transformation Groups in Differential Geometry*. Springer-Verlag, 1972.
- [20] KORAS, M. Linearization of reductive group-actions. *Lecture Notes Math.* 956 (1982), 92–98.
- [21] KRAFT, H., T. PETRIE and G. W. SCHWARZ. *Topological Methods in Algebraic Transformation Groups*. Birkhäuser, 1989.
- [22] KUSHNIRENKO, A. G. Linear-equivalent action of a semisimple Lie group in the neighborhood of a stationary point. *Functional Anal. Appl.* 1 (1967), 89–90.
- [23] LIE, S. Theorie der Transformationsgruppen. *Math. Ann.* 16 (1880), 441–528. See also *Sophus Lie's Transformation group paper*, translated by M. Ackermann, comments by R. Hermann, Math. Sci. Press (1975).
- [24] LIVINGSTON, E. S. and D. L. ELLIOTT. Linearization of families of vector fields. *J. Diff. Equations* 55 (1984), 289–299.
- [25] LUBOTZKY, A. *Discrete Groups, Expanding Graphs and Invariant Measures*. Birkhäuser, 1994.

- [26] LYCHAGIN, V. V. Singularities of solutions, spectral sequences, and normal forms of Lie algebras of vector fields. *Math. USSR Izvestiya* 30 (1988), 549–575.
- [27] MARGULIS, G. A. *Discrete Subgroups of Semisimple Lie Groups*. Springer-Verlag, 1991.
- [28] MARTINET, J. Normalisation des champs de vecteurs holomorphes, séminaire Bourbaki, exposé 564, novembre 1980. *Lecture Notes Math.* 901.
- [29] MONTGOMERY, D. and C. T. YANG. Differentiable actions on homotopy seven spheres. *Trans. Amer. Math. Soc.* 122 (1966), 480–498.
- [30] MONTGOMERY, D. and L. ZIPPIN. *Topological Transformation Groups*. Interscience Publ., 1955.
- [31] NARASIMHAN, R. *Analysis on real and complex manifolds*. Masson & Cie and North-Holland, 1973.
- [32] NEUMAN, M. *Integral Matrices*. Academic Press, 1972.
- [33] ONISHCHIK, A. L. *Lie Groups and Lie Algebras I*. Springer-Verlag, 1993.
- [34] POINCARÉ, H. Sur les propriétés des fonctions définies par les équations aux différences partielles. *Thèse*, Paris, (1879). *Œuvres*, tome I, pp. XLIV à CXXIX, Gauthier-Villars, 1951.
- [35] POUREZZA, I. and J. HUBBARD. The space of closed subgroups of \mathbf{R}^2 . *Topology* 18 (1979), 143–146.
- [36] ROBINSON, D. J. S. *A Course in the Theory of Groups*. Springer-Verlag, 1993.
- [37] SCHNEIDER, C. R. $SL(2, \mathbf{R})$ actions on surfaces. *Amer. J. Math.* 96 (1974), 511–528.
- [38] SCHWARZ, G. W. Exotic algebraic group actions. *C. R. Acad. Sci. Paris, Sér. I*, 309 (1989), 89–94.
- [39] SMALE, S. Generalized Poincaré conjecture in dimensions greater than four. *Ann. of Math.* 74 (1961), 391–406.
- [40] STEINBERG, R. Some consequences of the elementary relations in SL_n , Finite groups—coming of age (Montreal, Que., 1982). *Contemp. Math.* 45. Amer. Math. Soc., Providence, R.I. (1985), 335–350.
- [41] STERNBERG, S. On local C^n contractions of the real line. *Duke Math. J.* 24 (1957), 97–102.
- [42] ——— Local contractions and a theorem of Poincaré. *Amer. J. Math.* 79 (1957), 809–824.
- [43] ——— On the structure of local homeomorphisms of Euclidean n -space II. *Amer. J. Math.* 80 (1958), 623–631.
- [44] ——— The structure of local homeomorphisms III. *Amer. J. Math.* 81 (1959), 578–604.
- [45] THURSTON, W. P. A generalization of the Reeb stability theorem. *Topology* 13 (1974), 347–352.
- [46] UCHIDA, F. Real analytic $SL(n, \mathbf{R})$ actions on spheres. *Tohoku Math. J.* 33 (1981), 145–175.
- [47] ——— Construction of a continuous $SL(3, \mathbf{R})$ action on 4-sphere. *Publ. Res. Inst. Math. Sci.* 21 (1985), 425–431.
- [48] ——— Real analytic actions of complex symplectic groups and other classical Lie groups on spheres. *J. Math. Soc. Japan* 38 (1986), 661–677.

- [49] VAN EST, W. T. On the algebraic cohomology concepts in Lie groups, I and II. *Indagationes Math.* 15 (1955), 225–233 and 286–294.
- [50] ZIMMER, R. J. *Ergodic Theory and Semisimple Groups*. Birkhäuser, 1984.

(Reçu le 19 décembre 1996; version révisée reçue le 10 mars 1997)

Grant Cairns

School of Mathematics
La Trobe University
Melbourne 3083
Australia
E-mail: G.Cairns@latrobe.edu.au

Étienne Ghys

École Normale Supérieure de Lyon
UMPA, UMR 128 CNRS
69364 Lyon Cedex 07
France
E-mail: ghys@umpa.ens-lyon.fr

Vide-leer-empty