Zeitschrift: L'Enseignement Mathématique
Band: 44 (1998)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HOMFLY POLYNOMIAL VIA AN INVARIANT OF COLORED PLANE GRAPHS

Bibliographie

Autor: Murakami, Hitoshi / Yamada, Shuji
DOI: https://doi.org/10.5169/seals-63908

8_{17}	$q^{n+2} - 2q^{n+1} + 3q^n - 2q^{n-1} + q^{n-2} - q^3 + 2q^2 - 4q + 5 - 4q^{-1} + 2q^{-2} - q^{-3} + q^{-n+2} - 2q^{-n+1} + 3q^{-n} - 2q^{-n-1} + q^{-n-2}$
8_{18}	$q^{n+2} - 3q^{n+1} + 3q^n - 3q^{n-1} + q^{n-2} - q^3 + 3q^2 - 4q + 7 - 4q^{-1} + 3q^{-2} - q^{-3} + q^{-n+2} - 3q^{-n+1} + 3q^{-n} - 3q^{-n-1} + q^{-n-2}$
8_{19}	$q^{-3n+3} + q^{-3n+1} + q^{-3n} + q^{-3n-1} + q^{-3n-3} - q^{-4n+2} - q^{-4n+1} - q^{-4n} - q^{-4n-1} - q^{-4n-2} + q^{-5n}$
8_{20}	$-q^{2n+1} - q^{2n-1} + q^{n+2} + 2q^n + q^{n-2} - q + 1 - q^{-1}$
8_{21}	$q^{3n+1} - q^{3n} + q^{3n-1} - q^{2n+2} + q^{2n+1} - 3q^{2n} + q^{2n-1} - q^{2n-2} + 2q^{n+1} - q^n + 2q^{n-1}$

REFERENCES

[1] FREYD, P., D. YETTER, J. HOSTE, W. B. R. LICKORISH, K. MILLETT and A. OCNEANU. A new polynomial invariant of knots and links. *Bull. Amer. Math. Soc. (N.S.) 12* (1985), 239–246.

[2] JIMBO, M. Quantum R matrix for the generalized Toda system. *Comm. Math. Phys. 102* (1986), 537–547.

[3] JONES, V. F. R. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.) 12* (1985), 103–111.

[4] —— On knot invariants related to some statistical mechanical models. *Pacific J. Math. 137* (1989), 311–336.

[5] KAUFFMAN, L. H. State models and the Jones polynomial. *Topology 26* (1987), 395–407.

[6] KIRBY, R. and P. MELVIN. The 3-manifold invariants of Witten and Reshetikhin–Turaev for $\mathfrak{sl}(2, \mathbf{C})$. *Invent. Math. 105* (1991), 473–545.

[7] KIRILLOV, A. N. and N. YU. RESHETIKHIN. Representations of the algebra $U_q(\mathfrak{sl}(2))$, q-orthogonal polynomials and invariants of links. *Infinite Dimensional Lie Algebras and Groups*. V.G. Kac, ed., Advanced Series in Mathematical Physics, vol. 7, World Scientifics, Singapore, 1989.

[8] KUPERBERG, G. The quantum G_2 link invariant. *Internat. J. Math. 5* (1994), 61–85.

[9] LICKORISH, W. B. R. The skein method for three-manifold invariants. *J. Knot Theory Ramifications 2* (1993), 171–194.

[10] —— *An Introduction to Knot Theory*. Graduate Texts in Mathematics, vol. 175, Springer-Verlag, 1997.

[11] MORTON, H. R. Invariants of links and 3-manifolds from skein theory and from quantum groups. *Topics in Knot Theory*. M.E. Bozhüyük, ed., Kluwer, 1993.

[12] OHTSUKI, T. and S. YAMADA. Quantum $SU(3)$ invariants of 3-manifolds via linear skein theory. *J. Knot Theory Ramifications 6* (1997), 373–404.

[13] PRZYTYCKI, J. H. and P. TRACZYK. Invariant of links of Conway type. *Kobe J. Math. 4* (1988), 115–139.

[14] RESHETIKHIN, N. YU. and V. G. TURAEV. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys. 127* (1990), 1–26.

[15] ROLFSEN. D. *Knots and Links*. Mathematics Lecture Series, vol. 7, Publish or Perish, Inc., Houston, Texas, 1990. Second printing, with corrections.

[16] TURAEV, V. G. The Yang-Baxter equation and invariants of links. *Invent. Math. 92* (1988), 527–553.

[17] YOKOTA, Y. Skeins and quantum $SU(N)$ invariants of 3-manifolds. *Math. Ann. 307* (1997), 109–138.

(Reçu le 10 mars 1998)

Hitoshi Murakami

Department of Mathematics
School of Science and Engineering
Waseda University
Ohkubo, Shinjuku-ku
Tokyo 169
Japan
email: hitoshi@uguisu.co.jp

Tomotada Ohtsuki

Department of Mathematical and Computing Sciences
Tokyo Institute of Technology
Oh-Okayama, Meguro-ku
Tokyo 152
Japan
email: tomotada@is.titech.ac.jp

Shuji Yamada

Department of Computer Science
Kyoto Sangyo University
Kamigamo Motoyama, Kita-ku
Kyoto 603
Japan
email: yamada@cc.kyoto-su.ac.jp