

5. The recurrence function

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **44 (1998)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

is the Sturmian sequence obtained as the coding of the orbit of $-\rho/\alpha$ under the rotation by angle $1/\alpha$, with respect to the partition

$$\{]0, 1 - 1/\alpha],]1 - 1/\alpha, 1]\} .$$

Indeed, if $n = \lfloor \alpha m + \rho \rfloor$, then $\lceil 1/\alpha(n+1) - \rho/\alpha \rceil = m+1 = 1 + \lceil n/\alpha - \rho/\alpha \rceil$, and if $\lfloor \alpha m + \rho \rfloor < n < \lfloor \alpha(m+1) + \rho \rfloor$, then $\lceil 1/\alpha(n+1) - \rho/\alpha \rceil = \lceil n/\alpha - \rho/\alpha \rceil$.

5. THE RECURRENCE FUNCTION

Let us deduce now from the three distance and three gap theorems a simple proof of the following result originally due to Morse and Hedlund concerning the recurrence function of a Sturmian sequence (see [40]).

Recall that a sequence u is called *minimal* or *uniformly recurrent* if every factor of u appears infinitely often and with bounded gaps or, equivalently, if for any integer n , there exists an integer m such that every factor of u of length m contains every factor of u of length n . Note that it is equivalent (see [30]) to the *minimality* of the dynamical system $(\overline{\mathcal{O}(u)}, T)$, i.e., the orbit of every element of $\overline{\mathcal{O}(u)}$ is dense, or equivalently every sequence in the orbit closure of u has the same set of factors as u .

The recurrence function φ of a minimal sequence u is defined by

$$\varphi(n) = \min \{m \in \mathbf{N} \text{ such that } \forall B \in L_m, \forall A \in L_n, A \text{ is a factor of } B\} ,$$

where L_n denotes the set of factors of u of length n , i.e., $\varphi(n)$ is the size of the smallest window that contains all factors of length n whatever its position in the sequence.

THEOREM 11. *Let u be a Sturmian sequence with angle α . Let $(q_k)_{k \in \mathbf{N}}$ denote the sequence of denominators of the convergents of the continued fraction expansion of α . The recurrence function φ of this sequence satisfies for any non zero integer n :*

$$\varphi(n) = n - 1 + q_k + q_{k-1}, \text{ where } q_{k-1} \leq n < q_k .$$

Proof of Theorem 11. Let $u \in \{0, 1\}^{\mathbf{N}}$ be a Sturmian sequence. There exist a real number x and an irrational number α in $]0, 1[$ such that $u_n = 0 \Leftrightarrow \{x + n\alpha\} \in I_0$, with $I_0 = [0, \alpha[$ or $I_0 =]0, \alpha]$ (see Section 2.1). Let $I_1 = [\alpha, 1[$ (respectively, $] \alpha, 1]$) if $I_0 = [0, \alpha[$ (respectively, $I_0 =]0, \alpha]$). Let us denote by R the rotation of the circle by angle α . Assume we are given

a positive integer n . We have seen in Section 2.1 that the word $w_1w_2\cdots w_n$ defined on $\{0, 1\}$ appears in u if and only if

$$I(w_1, \dots, w_n) = \bigcap_{j=0}^{n-1} R^{-j}(I_{w_{j+1}}) \neq \emptyset.$$

We deduce from this that every Sturmian sequence of angle α has the same factors as u and thus belongs to the orbit closure of u . Conversely, each sequence of the orbit closure of u is a Sturmian sequence of angle α . Hence the closed orbit of any Sturmian sequence is equal to the set of all Sturmian sequences of the same angle. This implies the minimality of any Sturmian sequence and that Sturmian sequences of the same angle have the same recurrence function; hence we can suppose here that $x = 0$.

Theorem 11 can easily be deduced from the following two lemmata. We omit the proof of Lemma 5, which is straightforward.

LEMMA 4. *Let δ_n be the smallest length of the nonempty intervals $I(w_1, \dots, w_n)$, where w_1, \dots, w_n belong to $\{0, 1\}$. Let l_n be the greatest gap between the successive integers k such that $\{k\alpha\} \in [0, \delta_n[$. We have*

$$\varphi(n) = n - 1 + l_n.$$

LEMMA 5. *Let $(q_k)_{k \in \mathbb{N}}$ denote the sequence of denominators of the convergents of the continued fraction expansion of α . Let k be an integer such that $q_{k-1} \leq n < q_k$. Then we have*

$$\delta_n = \eta_{k-1} \quad \text{and} \quad l_n = q_k + q_{k-1}.$$

Proof of Lemma 4. A set of points is said to visit an interval if one of these points belongs to this interval. By definition of l_n , every set of l_n consecutive points of the sequence $(\{k\alpha\})_{k \in \mathbb{N}}$ visits every interval of length δ_n (see above Remarks). Therefore they visit every nonempty interval of the form $I(w_1, \dots, w_n)$, by definition of δ_n . Let B be a factor of u of length $n - 1 + l_n$; there exists an integer K such that B corresponds to the $n - 1 + l_n$ consecutive points $\{K\alpha\}, \dots, \{(K + n - 1 + l_n - 1)\alpha\}$. The set of the l_n first points of this sequence of points visits every interval of the form $I(w_1, \dots, w_n)$, thus B contains every factor of u of length n . This implies that $\varphi(n) \leq n - 1 + l_n$.

By definition of l_n and by density of $(\{k\alpha\})_{k \in \mathbb{N}}$, there exists a sequence of $l_n - 1$ points of the sequence $(\{k\alpha\})_{k \in \mathbb{N}}$ which do not visit an interval of

the form $I(w_1, \dots, w_n)$ of length δ_n ; therefore, there exists a factor of u of length $l_n - 2 + n$ which does not contain the factor $w_1 \cdots w_n$. This shows that $\varphi(n) \geq n - 1 + l_n$. The lemma is thus proved.

REMARK. Note that in the case of the Fibonacci sequence ($\alpha = \frac{\sqrt{5}-1}{2}$), the recurrence function satisfies, for $F_{k-1} < n \leq F_k$,

$$\varphi(n) = n - 1 + F_{k+1},$$

where $(F_n)_{n \in \mathbb{N}}$ denotes the Fibonacci sequence $F_{n+1} = F_n + F_{n-1}$, with $F_0 = 1$ and $F_1 = 2$.

This result is extended in [13] to the fixed point of the substitution σ introduced by Rauzy which generalizes the Fibonacci substitution and is defined by $\sigma(0) = 01$, $\sigma(1) = 02$, $\sigma(2) = 0$.

THEOREM 12. Let T_n denote the so-called Tribonacci sequence defined as follows: $T_{k+3} = T_{k+2} + T_{k+1} + T_k$, with $T_0 = 0$, $T_1 = 0$, $T_2 = 1$. The recurrence function φ of the fixed point beginning with 0 of the Rauzy substitution satisfies for any positive integer n :

$$\varphi(n) = n - 1 + T_{k+6}, \quad \text{where} \quad \sum_0^{k+1} T_i < n \leq \sum_0^{k+2} T_i.$$

6. HIGHER-DIMENSIONAL GENERALIZATIONS

6.1 TWO-DIMENSIONAL GENERALIZATIONS AND BEATTY SEQUENCES

Let us consider now some two-dimensional versions of the three distance and three gap theorems. Such generalizations were introduced by Fraenkel and Holzman in [26] in order to give an upper bound for the number of gaps in the intersection of two Beatty sequences. They first reduce this problem to a two-dimensional version of the three distance theorem, conjectured by Simpson and Holzman and proved by Geelen and Simpson (see [29]). Then they deduce from this theorem a bound for the number of gaps in the intersection of two Beatty sequences, when at least one of the moduli is rational.

Let us first give the two-dimensional version of the three gap theorem introduced by Fraenkel and Holzman. We will use the same notation as in [26]: for any pair of real numbers (x, y) , $\{(x, y)\}$ means the equivalence class of $(x, y) \bmod \mathbf{Z}^2$, i.e., $\{(x, y)\}$ belongs to the torus \mathbf{T}^2 .