

4.1 Free groups

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

4. NORMS OF RANDOM WALK OPERATORS

Now we will show how Theorem 3 can be used in the problem of computing the norm of the random walk operator P on some groups. Our strategy is as follows: we want to find a positive eigenfunction for the operator P which satisfies the generalized Følner condition. By Theorem 3 such an eigenfunction always exists and the eigenvalue corresponding to this eigenfunction is equal to the norm of the operator P . Theorem 3 is a particular case of Theorem 2 which can also be helpful in computing the norms of more general operators as shown in Section 4.3.

4.1 FREE GROUPS

First of all, as a simple illustration of this method, we will compute the norm of the simple random walk operator on free groups, which was first done by Kesten (see [9]) using a different method.

THEOREM 6 (Kesten). *Let Γ be the free group generated by the standard symmetric set of generators S . The norm of the simple random walk operator P associated to (Γ, S) is equal to*

$$\|P\| = \frac{2\sqrt{\#S - 1}}{\#S}.$$

Proof. The Cayley graph of (Γ, S) is a homogeneous tree T_k of degree $k = \#S$. We draw the tree T_k with level lines as in Figure 3 (level lines are marked by dotted lines). Let us choose arbitrarily a line as the line of level 0. We construct a function on vertices of this tree which depends only on the level of the vertex. For a vertex $v \in T_k$ we denote by $|v|$ its level. We define $f: T_k \rightarrow \mathbf{R}_+$ as follows

$$f(v) = \left(\frac{1}{\sqrt{k-1}} \right)^{|v|}.$$

One has

$$Pf = \frac{2\sqrt{k-1}}{k}f.$$

Let A_n be the set of vertices in T_k consisting of a chosen vertex e from the level 0 and the vertices lying below e up to the level n (in Figure 3 the vertices of A_2 are marked with circles). Then

$$\sum_{v \in A_n} f^2(v) = n + 1,$$

$$\sum_{v \in \partial A_n} f^2(v) = 2.$$

This shows that $\{A_n\}_{n=1}^{\infty}$ is a generalized Følner sequence and by Theorem 3

$$\|P\| = \frac{2\sqrt{k-1}}{k}. \quad \square$$

4.1.1 REMARKS ON GENERALIZED GROWTH

Let Γ be a group generated by a finite, symmetric set S . For $id \neq \gamma \in \Gamma$ we define its length $|\gamma|$ as the minimal number of generators from S needed to represent γ , i.e.

$$|\gamma| = \min\{n; \gamma = s_{i_1} \dots s_{i_n}, s_{i_j} \in S\},$$

and we declare $|id| = 0$.

The growth function (see [10], [18]) of the pair (Γ, S) associates to each integer $n \geq 0$ the number $\beta(\Gamma, S)(n)$ of elements $\gamma \in \Gamma$ such that $|\gamma| \leq n$, i.e.

$$\beta(\Gamma, S)(n) = \#\{\gamma \in \Gamma; |\gamma| \leq n\}.$$

One is often interested only in the type of the growth function. For instance, we say that the group Γ is of polynomial growth if there exist constants c and D such that

$$c^{-1}n^D \leq \beta(\Gamma, S)(n) \leq cn^D.$$

The exponent D does not depend on the set of generators S . If the growth function is bounded by a polynomial, it is known (see [6]) that Γ is of polynomial growth and D is an integer. For a group of polynomial growth with the exponent D , it is known (see [19]) that there exists a constant c such that

$$(5) \quad c^{-1}n^{-\frac{D}{2}} \leq P^{2n}(id, id) \leq cn^{-\frac{D}{2}},$$

where $P^{2n}(id, id)$ is the probability of the return to the identity element of the simple random walk after $2n$ steps.

It seems natural to define a generalized growth function, using an eigenfunction of P . Let f be a positive eigenfunction of P corresponding to the eigenvalue equal to the norm of P , i.e.

$$Pf = \|P\|f.$$

The generalized growth function $\beta(\Gamma, S, f)$ associates to each positive integer n the number

$$\beta(\Gamma, S, f)(n) = \sum_{\gamma \in \Gamma, |\gamma| \leq n} f^2(\gamma),$$

i.e., each element in the ball of radius n is counted with weight f^2 .

Let us compute the generalized growth function in a particular case. Let P be the simple random walk operator on the free group with the standard set of generators of cardinality k as in Section 4.1. Let g be the unique radial eigenfunction of P corresponding to the eigenvalue $\|P\|$ and such that $g(id) = 1$. Explicitly we have:

$$g(\gamma) = \left(\frac{k-2}{k} |\gamma| + 1 \right) \left(\frac{1}{\sqrt{k-1}} \right)^{|\gamma|}.$$

Then we have

$$\sum_{\gamma \in \Gamma, |\gamma| \leq n} g^2(\gamma) = n^3 \left(\frac{k^2 - 4k + 4}{3k^2 - 3k} \right) + n^2 \left(\frac{3k^2 - 8k + 4}{2k^2 - 2k} \right) + n \left(\frac{7k^2 - 16k + 4}{6k^2 - 6k} \right) + 1.$$

This shows that the generalized growth is like n^3 . In particular the sequence of balls is a generalized Følner sequence.

By analogy to (5) we conjecture that the fact that the generalized growth function for the free groups is like n^3 explains that for the free groups one has (see [16]):

$$c^{-1} \lambda^{2n} n^{-\frac{3}{2}} \leq P^{2n}(id, id) \leq c \lambda^{2n} n^{-\frac{3}{2}},$$

where c is a constant and λ is the norm of P .

4.2 FREE PRODUCTS OF FINITE GROUPS

Random walks on free products of finite groups were already considered in [1], [3], [17] and [21].

Let us consider the group $\mathbf{Z}_m \star \mathbf{Z}_n$ with the following generating set:

- if $m \neq 2$ we take $\{\pm 1\}$ as generators of $\mathbf{Z}_m = \{0, 1, \dots, m - 1\}$;
- we take $\{1\}$ as a generator of $\mathbf{Z}_2 = \{0, 1\}$.