

Introduction

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

A FREE GROUP ACTING ON \mathbf{Z}^2 WITHOUT FIXED POINTS

by SATÔ Kenzi

ABSTRACT. The group of all orientation-preserving affine transformations of the plane has a non-abelian free subgroup which stabilizes \mathbf{Z}^2 and which acts on \mathbf{Z}^2 without non-trivial fixed points.

INTRODUCTION

Let G be a group acting on a non-empty set X . The following two conditions are known to be equivalent (see [D], and Theorems 4.5 and 4.8 in [W]):

- (a) *there exists a non-abelian free subgroup of G whose action on X is locally commutative;*
- (b) *there exists a G -paradoxical decomposition of X using 4 pieces, namely a partition of X in parts P_0, P_1, P_2, P_3 and elements $\alpha_0, \alpha_1, \alpha_2, \alpha_3$ in G such that*

$$X = P_0 \sqcup P_1 \sqcup P_2 \sqcup P_3 = \alpha_0(P_0) \sqcup \alpha_1(P_1) = \alpha_2(P_2) \sqcup \alpha_3(P_3).$$

Moreover, in the situation of (b), it can be shown that the subgroup of G generated by $\alpha_0^{-1}\alpha_1$ and $\alpha_2^{-1}\alpha_3$ is free of rank 2. (The symbol \sqcup denotes disjoint union. Recall that an action of a group H on X is *locally commutative* if the stabilizer $\{h \in H \mid h(x) = x\}$ is commutative for all $x \in X$, i.e. if two elements of H which have a common fixed point commute; trivial examples of locally commutative actions are actions *without non-trivial fixed points*, for which $\{h \in H \mid h(x) = x\}$ is reduced to $\{1\}$ for all $x \in X$.)

For example, the group $SO_3(\mathbf{R})$ of rotations of the unit sphere \mathbf{S}^2 has such a free subgroup: this was discovered by F. Hausdorff (see, e.g., [Š], or Theorem 2.1 in [W]). It implies the following result, for which we refer to [BT] and Theorem 3.11 in [W]; we denote by $SG_3(\mathbf{R})$ the group of all orientation-preserving isometries of \mathbf{R}^3 .

THE BANACH-TARSKI PARADOX. *Any two bounded subsets U and V of the 3-dimensional Euclidean space \mathbf{R}^3 with non-empty interiors are $SG_3(\mathbf{R})$ -equidecomposable. In other words, one can partition U into a finite number of pieces and reconstruct V from the same number of respectively $SG_3(\mathbf{R})$ -congruent pieces.*

The Banach-Tarski paradox holds similarly for higher dimensional Euclidean spaces, but not for \mathbf{R} and \mathbf{R}^2 ; the reason is that neither $SG_1(\mathbf{R})$ nor $SG_2(\mathbf{R})$, which are soluble groups, contain free subgroups of rank 2. (There are other known examples of free groups acting without non-trivial fixed points on familiar spaces. See e.g., [B], [DS], and [S2]. The proof of the Banach-Tarski paradox requires the axiom of choice, because the proof of the equivalence of conditions (a) and (b) requires it. But similar paradoxes hold for rational spheres of the form $(\sqrt{q}\mathbf{S}^2) \cap \mathbf{Q}^3$, as can be shown *without* the axiom of choice from the countability of rational spheres. See [S1], and [S3].) In dimension 2, von Neumann has exhibited a Banach-Tarski paradox with respect to the group $SA_2(\mathbf{R})$ of affine transformations of \mathbf{R}^2 that preserve area and orientation ([V], and Theorem 7.3 of [W]). The following problem was raised in [MW]; see also the discussion which follows Proposition 7.1 in [W].

PROBLEM ([MW], [W]). *Does $SA_2(\mathbf{R})$ contain a free subgroup of rank 2 whose action on \mathbf{R}^2 is locally commutative?*

Indeed, these authors asked more specifically if the group generated by

$$\alpha: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

and

$$\beta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

satisfies the requirements of the problem. We observe here that the answer is “no”, because both $\alpha^{-2}\beta^2$ and $\alpha^{-1}\beta^{-1}\alpha\beta$ fix the origin.

Though we cannot solve the above problem, the purpose of this note is to show that, if one replaces \mathbf{R}^2 by \mathbf{Z}^2 , the new problem has a positive solution. In fact, we will prove the following result, which shows somewhat more, namely that the action on \mathbf{Z}^2 may be an action without non-trivial fixed points, rather than only locally commutative. We denote by $\text{SA}_2(\mathbf{Z})$ the group of all transformations $\vec{x} \mapsto A\vec{x} + \vec{a}$ of \mathbf{Z}^2 , with $A \in \text{SL}_2(\mathbf{Z})$ and $\vec{a} \in \mathbf{Z}^2$.

THEOREM. *The group $\text{SA}_2(\mathbf{Z})$ has a free subgroup F_2 of rank 2 which acts on \mathbf{Z}^2 without non-trivial fixed points, namely the subgroup generated by*

$$\zeta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

and

$$\eta: \begin{pmatrix} x \\ y \end{pmatrix} \mapsto \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 3 \\ 2 \end{pmatrix}.$$

The theorem implies the existence of a partition of \mathbf{Z}^2 into three pieces P , Q and R such that the six pieces P , Q , R , $P \sqcup Q$, $Q \sqcup R$, $R \sqcup P$ are pairwise F_2 -congruent, without the axiom of choice ([S0], and Corollary 4.12 in [W]).

As observed in the discussion which follows Proposition 7.1 in [W], it is known that the above theorem does not carry over to \mathbf{R}^2 ; more precisely, it is known that a subgroup of $\text{SA}_2(\mathbf{R})$ which acts on \mathbf{R}^2 without non-trivial fixed points is soluble, and consequently does not contain non-commutative free subgroups.

PROOF OF THE MAIN RESULT

Recall that a matrix in $\text{SL}_2(\mathbf{Z})$ is *hyperbolic* if the absolute value of its trace is strictly larger than 2, or equivalently if it has an eigenvalue of absolute value strictly larger than 1.

LEMMA 0. *The subgroup of $\text{SL}_2(\mathbf{Z})$ generated by*

$$\begin{pmatrix} 7 & 3 \\ 9 & 4 \end{pmatrix} \text{ and } \begin{pmatrix} 94 & 39 \\ 147 & 61 \end{pmatrix}$$

is free of rank 2 and all its elements distinct from the identity are hyperbolic.