

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 46 (2000)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THREE REMARKS ON GEODESIC DYNAMICS AND FUNDAMENTAL GROUP
Autor: Gromov, Mikhaïl
Anhang: Why the Appendices were not written : author's apologies to the readers
DOI: <https://doi.org/10.5169/seals-64809>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 18.04.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

- [Ma] MANNING, A. Topological entropy and the first homology group. *Springer Lecture Notes in Math.* 468 (1974), 185–191.
- [Nie] NIELSEN, J. Über topologische Abbildungen geschlossener Flächen. *Abh. Math. Sem. Univ. Hamburg* 3 (1924), 246–260.
- [Pu] PUGH, C. On the entropy conjecture. *Springer Lecture Notes in Math.* 468 (1974), 257–262.
- [Sh] SHUB, M. Endomorphisms of compact manifolds. *Amer. J. Math.* 91 (1969), 185–199.
- [Sh-S] SHUB, M. and D. SULLIVAN. A remark on the Lefschetz fixed point formula. *Topology* 13 (1974), 189–191.

WHY THE APPENDICES WERE NOT WRITTEN :

AUTHOR'S APOLOGIES TO THE READERS

APPENDIX 2. The stable homeomorphism suggests a geometric link between the homotopy and topological invariance of Pontryagin classes, at least for manifolds with negative curvature but I did not manage to forge this to my satisfaction till 1996 (see [Gro₂]); also see [Fa-Jo] for a deeper analysis.

APPENDIX 3. One can define a notion of hyperbolicity for an automorphism α of an arbitrary finitely generated group Γ , such that (Γ, α) functorially defines a Bowen-Franks hyperbolic system (see [Gro₁]). Unfortunately, this class of (Γ, α) is rather limited, e.g. is not closed under free products and does not include hyperbolic automorphisms of surface groups. I still do not know what the right setting is.

APPENDIX 4. An obvious example of semi-hyperbolicity is provided by non-strictly expanding endomorphisms, where the geometric picture is rather clear. However, I still do not see a functorial description, in the spirit of the symbolic dynamics, of more general semi-hyperbolic systems, not even for the geodesic (or Weil chamber) flows on locally symmetric spaces (compare [B-G-S] and [Br-Ha]).

APPENDIX 5. The section on entropy was inspired by Manning's paper [Ma₁], but I was unaware of the prior paper by Dinaburg (see [Din]) that essentially contained the entropy estimate for geodesic flows (also discussed in [Ma₂]). On the other hand, estimating the entropy of an endomorphism (or an automorphism) f in terms of $f_*: \pi_1 \rightarrow \pi_1$ appears now much less clear than it seemed to me back in 1976. It is not hard to bound the entropy from below

via the “asymptotic stretch” of $f_*: \pi_1 \rightarrow \pi_1$ with respect to the word metric in π_1 (see [Bow]). But this is not sharp even for linear automorphisms of tori T^n , where the entropy is expressed by the “ k -dimensional stretch” on H_1 for some $k \leq n$ that equals the spectral radius of f_* on H_k . Such k -stretch can be defined, in general, in terms of $f_*: \pi_1(S) \rightarrow \pi_1(S)$ and the classifying map $S \rightarrow K(\pi_1, 1)$ (refining the spectral radius of f_* on H^k coming from $K(\pi_1, 1)$), but my obvious “proof” of the lower bound on the entropy by this k -stretch missed a hidden trap. This was also overlooked in [Ma₃] (for $f_*: H_1 \rightarrow H_1$, where a proper identification of the “ k -stretch” with the spectral radius needs extra work), as was pointed out to me much later by David Fried. (The difficulty already appears for closed subsets S in the torus T^n invariant under linear automorphisms f of T^n , where one wishes to estimate the entropy of $f|_S$ in terms of f_* acting on the *spectral* cohomology of S coming from T^n . On the other hand, the case of $T^n \rightarrow T^n$ is settled in [Mi-Pr].)

APPENDIX 6. Probably, the recent progress in Nielsen theory allows a description of the cases, where $\text{card}(\text{Fix } f)$ is well controlled from below by some twisted Lefschetz number (see [Fel]).

APPENDIX 7. Nothing interesting to say.

APPENDIX 8. Minima of geometric functionals related to the logical complexity have been studied in depth by A. Nabutovski (see [Na] and references therein). Yet I do not feel ready yet to write this Appendix. For example, I do not see what is the actual influence of a suitable (?) complexity measure of $\pi_1(V)$ on the Plateau problem in V .

BIBLIOGRAPHY

- [B-G-S] BALLMANN, W., M. GROMOV and V. SCHROEDER. *Manifolds of Non-Positive Curvature*. Birkhäuser, Boston, 1985.
- [Bow] BOWEN, R. Entropy and the fundamental group. In: *The Structure of Attractors in Dynamical Systems*. (Proc. Conf., North Dakota State Univ., Fargo, N.D. (1977)), 21–29. Lecture Notes in Math., 668, Springer, Berlin, 1978.
- [Br-Ha] BRIDSON, M. R. and A. HAEFLIGER. *Metric Spaces of Non-Positive Curvature*. Grundlehren der mathematischen Wissenschaften, Vol. 319. Springer, Berlin, 1999.
- [Din] DINABURG, E. I. A connection between various entropy characterizations of dynamical systems (Russian). *Izv. Akad. Nauk SSSR Ser. Mat.* 35 (1971), 324–366.