

6.6 Examples

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.07.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

6.6 EXAMPLES

Keeping the notations of the previous section, we shall illustrate Theorem 17.

a. TOTALLY GEODESIC TRANSFORM. As in Section 4.1 **a**, let $X = G/K$ be a Riemannian symmetric space of the noncompact type and $y_o = \text{Exp } \mathfrak{s}$ the origin in the dual space $Y = G/H$. By (3) we have $\mathfrak{k} + \mathfrak{h} = \mathfrak{k} \oplus \mathfrak{s}$, therefore Theorem 17 (i) applies with $\mathfrak{t} = \mathfrak{s}^\perp$, the orthogonal of \mathfrak{s} in \mathfrak{p} .

b. HOROCYCLE TRANSFORM. Again $X = G/K$ is a Riemannian symmetric space of the noncompact type (see Notations, **d**), but the dual space is now the space of horocycles $Y = G/MN$. We recall Harish-Chandra's isomorphism of algebras ([9], p. 306)

$$\Gamma : \mathbf{D}(X) \longrightarrow \mathbf{D}(A)^W,$$

where $\mathbf{D}(A)^W$ is the subalgebra of W -invariant differential operators in $\mathbf{D}(A)$. The definition of Γ will be recalled during the next proof.

PROPOSITION 18. *Given $v \in C^\infty(Y)$, the function of $x = gK$ and $a \in A$ given by*

$$w(x, a) = a^\rho R_a^* v(x) = a^\rho \int_K v(gkaN) dk$$

is a solution of the system of multitemporal wave equations

$$P_{(x)} w(x, a) = \Gamma(P)_{(a)} w(x, a), \quad P \in \mathbf{D}(X), x \in X, a \in A.$$

Proof. Theorem 17 (ii) applies here with $T = A$, the abelian subgroup from the Iwasawa decomposition $G = KAN$; indeed $\mathfrak{k} + \mathfrak{h} = \mathfrak{k} + \mathfrak{m} + \mathfrak{n} = \mathfrak{k} \oplus \mathfrak{n}$, and $\mathfrak{g} = (\mathfrak{k} \oplus \mathfrak{n}) \oplus \mathfrak{a}$, $[\mathfrak{a}, \mathfrak{h}] \subset [\mathfrak{a}, \mathfrak{m}] + [\mathfrak{a}, \mathfrak{n}] \subset \mathfrak{n} \subset \mathfrak{h}$. By (31) we thus have

$$(32) \quad P_{(x)} R_a^* v(x) = D'_{(a)} R_a^* v(x),$$

where $D \in \mathbf{D}(G)^K$ is related to P by (28) and $D' \in \mathbf{D}(A)$ was characterized by

$$(33) \quad D - D' \in \mathfrak{k}\mathbf{D}(G) + \mathbf{D}(G)\mathfrak{n}.$$

To compare D' and $\Gamma(P)$ we recall that $\Gamma(P) = a^{-\rho} D_a \circ a^\rho$, where $D_a \in \mathbf{D}(A)$ is characterized by

$$(34) \quad D - D_a \in \mathfrak{n}\mathbf{D}(G) + \mathbf{D}(G)\mathfrak{k}.$$

Moreover $(Df)(a) = D_a(f(a))$ for $a \in A$, if $f \in C^\infty(G)$ is such that $f(ngk) = f(g)$ for any $g \in G, k \in K, n \in N$ ([9], p. 302 sq.).

Taking $u \in \mathcal{D}(G)$ we have, by a classical integral formula,

$$(35) \quad \int_G Df(g) \cdot u(g) dg = \int_{N \times A \times K} Df(a) \cdot u(nak) a^{-2\rho} dn da dk \\ = \int_{N \times A \times K} D_a f(a) \cdot u(nak) a^{-2\rho} dn da dk.$$

On the other hand, this integral can be written with the transpose operator tD as

$$\int_G Df(g) \cdot u(g) dg = \int_G f(g) {}^tDu(g) dg \\ = \int_A f(a) a^{-2\rho} da \int_{N \times K} ({}^tDu)(nak) dn dk.$$

But ${}^tD \in \mathbf{D}(G)^K$ therefore, for any $g \in G$,

$$\int_{N \times K} ({}^tDu)(ngk) dn dk = ({}^tD)_{(g)} \left(\int_{N \times K} u(ngk) dn dk \right).$$

The latter integral, as a function of g , is left N -invariant and right K -invariant so that

$$\int_{N \times K} ({}^tDu)(nak) dn dk = ({}^tD)_a \left(\int_{N \times K} u(nak) dn dk \right).$$

Since $({}^tD)_a = {}^t(D')$ obviously by (33) and (34), we obtain

$$\int_G Df(g) \cdot u(g) dg = \int_A D'(f(a)a^{-2\rho}) da \int_{N \times K} u(nak) dn dk \\ = \int_{N \times A \times K} (a^{2\rho} D' \circ a^{-2\rho}) f(a) \cdot u(nak) a^{-2\rho} dn da dk,$$

for any $f \in C^\infty(A)$ and any $u \in \mathcal{D}(G)$. Comparing with (35) it follows that

$$D_a = a^{2\rho} D' \circ a^{-2\rho}, \quad D' = a^{-\rho} \Gamma(P) \circ a^\rho,$$

whence the result by (32). \square

A slightly different proof can be obtained by decomposing the wave $a^\rho R_a^* v(gK)$ into *elementary horocycle waves* as follows. For $g \in G$ we denote by $A(g) \in A$ the A -component of g in the Iwasawa decompositions $G = NAK = ANK$ (we recall that A normalizes N), and by $K(g) \in K$ its K -component in the decompositions $G = KAN = KNA$.

PROPOSITION 19. (i) Given $f \in C^\infty(A)$ and $k \in K$, the function

$$w(gK, a) = a^{-\rho} f(A(k^{-1}g)a)$$

is a solution of the system of multitemporal wave equations

$$P_{(x)} w(x, a) = \Gamma(P)_{(a)} w(x, a), \quad P \in \mathbf{D}(X), \quad x \in X, \quad a \in A.$$

(ii) Given $v \in C^\infty(Y)$, the function of $x = gK$ and $a \in A$ given by

$$a^\rho R_a^* v(gK) = \int_K a^\rho v(gkaN) dk$$

is a solution of the same equations.

REMARKS. Part (i) is Proposition 8.5 in [12], p.118. Note that, k being fixed, the “wave surfaces” $A(k^{-1}g) = \text{constant}$ are parallel horocycles with the same normal $kM \in K/M$ (cf. [11], p.81). Indeed the equality $A(k^{-1}g) = a_o \in A$ is equivalent to $k^{-1}g \in a_oNK$, i.e. $g \cdot x_o \in ka_o \cdot y_o$.

If λ is a linear form on \mathfrak{a} and $f(a) = a^{i\lambda + \rho}$, the result (i) implies that $A(k^{-1}g)^{i\lambda + \rho}$ is, as a function of gK , an eigenfunction of all invariant operators $P \in \mathbf{D}(X)$; this is a fundamental result for harmonic analysis on X .

Part (ii) provides a simpler proof and a generalization of Proposition 8.6 in [12], p.118, where v was the Radon transform Ru of some $u \in \mathcal{D}(X)$. We refer to [12] or [13] for a detailed study of those multitemporal wave equations.

Proof of Proposition 19. (i) Both sides of the wave equation are invariant under the action of K on X ; we can therefore assume $k = e$. Now $w(gK, a) = a^{-\rho} f(A(g)a)$ is left N -invariant and right K -invariant as a function of g , and it will suffice to prove the result for $g = a \in A$.

By the decomposition (34) of D we have, for any $b \in A$,

$$D_{(g)} (f(A(g)b))|_{g=a} = (D_a)_{(a)} (f(ab)) = a^\rho \Gamma(P)_{(a)} (a^{-\rho} f(ab)) .$$

But $\Gamma(P)$ is an invariant differential operator on A , isomorphic to the additive group of a vector space, and we obtain

$$\begin{aligned} D_{(g)} (b^{-\rho} f(A(g)b))|_{g=a} &= a^\rho \Gamma(P)_{(a)} ((ab)^{-\rho} f(ab)) \\ &= a^\rho \Gamma(P)_{(b)} ((ab)^{-\rho} f(ab)) \\ &= \Gamma(P)_{(b)} (b^{-\rho} f(ab)) = \Gamma(P)_{(b)} (b^{-\rho} f(A(g)b))|_{g=a} . \end{aligned}$$

Thus (i) is proved for $g = a$.

(ii) Let $g \in G$, $k \in K$ and $k' = K(gk)$. Then $gk = k'a'n'$ with $a' \in A$ and $n' \in N$. It follows that $k'^{-1}g = a'n'k^{-1}$, therefore $a' = A(k'^{-1}g)$ and

$$gkaN = k'A(k'^{-1}g)aN.$$

For fixed g the map $k \mapsto K(gk) = k'$ is a diffeomorphism of K onto itself and, by the integral formula ([9], p. 197)

$$\int_K F(k') dk = \int_K A(k'^{-1}g)^{2\rho} F(k') dk',$$

we have

$$\begin{aligned} a^\rho R_a^* v(gK) &= a^\rho \int_K v(gkaN) dk \\ &= a^\rho \int_K v(k'A(k'^{-1}g)aN) dk \\ &= a^{-\rho} \int_K (A(k'^{-1}g)a)^{2\rho} v(k'A(k'^{-1}g)aN) dk'. \end{aligned}$$

By (i) applied to the functions $f(a) = a^{2\rho}v(k'aN)$, $k' \in K$, this is a solution of the wave equations. \square

COROLLARY 20 (Helgason). *If \mathfrak{g} has only one conjugacy class of Cartan subalgebras, there exists a differential operator $P \in \mathbf{D}(X)$ such that the horocycle Radon transform of $X = G/K$ is inverted by*

$$u(x) = PR^*Ru(x)$$

for $u \in \mathcal{D}(X)$, $x \in X$.

We prove it here by means of shifted transforms and wave equations; see [11], p. 116 for Helgason's original proof.

Proof. The assumption on \mathfrak{g} implies that, in the notation of (15), $C \cdot |c(\lambda)|^{-2}$ is a W -invariant polynomial on \mathfrak{a}^* . Let $P \in \mathbf{D}(X)$ be the corresponding operator under the isomorphism $\Gamma: \mathbf{D}(X) \rightarrow \mathbf{D}(A)^W$, so that $\Gamma(P)(i\lambda) = C \cdot |c(\lambda)|^{-2}$. By Theorem 13 and Proposition 19 (ii) (with $v = Ru$)

we have

$$\begin{aligned} u(x) &= \langle T_{(a)}, a^\rho R_a^* Ru(x) \rangle = \Gamma(D)_{(a)} (a^\rho R_a^* Ru(x)) \Big|_{a=e} \\ &= P_{(x)} (a^\rho R_a^* Ru(x)) \Big|_{a=e} = P_{(x)} R^* Ru(x). \quad \square \end{aligned}$$

REFERENCES

- [1] BERENSTEIN, C. and E. C. TARABUSI. Inversion formulas for the k -dimensional Radon transform in real hyperbolic spaces. *Duke Math. J.* 62 (1991), 613–631.
- [2] — An inversion formula for the horocyclic Radon transform on the real hyperbolic space. *Lectures in Appl. Math.* 30 (1994), 1–6.
- [3] ERDELYI, A., W. MAGNUS, F. OBERHETTINGER and F. TRICOMI. *Higher Transcendental Functions. Vol. I.* McGraw-Hill, 1953.
- [4] GEL'FAND, I., M. GRAEV and Z. SHAPIRO. Differential forms and integral geometry. *Funct. Anal. Appl.* 3 (1969), 101–114.
- [5] GRINBERG, E. Spherical harmonics and integral geometry on projective spaces. *Trans. Amer. Math. Soc.* 279 (1983), 187–203.
- [6] GUILLEMIN, V. and S. STERNBERG. *Geometric Asymptotics.* Math. Surveys and Monographs no. 14. Amer. Math. Soc., 1990.
- [7] HELGASON, S. The Radon transform on Euclidean spaces, compact two-point homogeneous spaces and Grassmann manifolds. *Acta Math.* 113 (1965), 153–180.
- [8] — *Differential Geometry, Lie Groups and Symmetric Spaces.* Academic Press, 1978.
- [9] — *Groups and Geometric Analysis.* Academic Press, 1984.
- [10] — The totally-geodesic Radon transform on constant curvature spaces. In: *Integral Geometry and Tomography. Contemp. Math.* 113 (1990), 141–149.
- [11] — *Geometric Analysis on Symmetric Spaces.* Math. Surveys and Monographs no. 39. Amer. Math. Soc., 1994.
- [12] — Radon transforms and wave equations. *Lecture Notes in Math.* 1684 (1998), 99–121.
- [13] — Integral geometry and multitemporal wave equations. *Comm. Pure Appl. Math.* 51 (1998), 1035–1071.
- [14] — *The Radon Transform.* 2nd edition, Birkhäuser, 1999.
- [15] KOBAYASHI, S. and K. NOMIZU. *Foundations of Differential Geometry. Vol. II.* Wiley, 1969.
- [16] RADON, J. Über die Bestimmung von Funktionen durch ihre Integralwerte längs gewisser Mannigfaltigkeiten. *Ber. Verh. Sächs. Akad. Wiss. Leipzig, Math. Nat. Kl.* 69 (1917), 262–277.