

ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT AND AUTOMORPHISMS OF $\sharp_{i=1}^b (S^2 \times S^5)$

Autor(en): **Schmitt, Alexander**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **48 (2002)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.07.2024**

Persistenter Link: <https://doi.org/10.5169/seals-66077>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

ON THE CLASSIFICATION OF CERTAIN PIECEWISE LINEAR
AND DIFFERENTIABLE MANIFOLDS IN DIMENSION EIGHT
AND AUTOMORPHISMS OF $\#_{i=1}^b(S^2 \times S^5)$

by Alexander SCHMITT

ABSTRACT. In this paper, we will be concerned with the explicit classification of closed, oriented, simply-connected spin manifolds in dimension eight with vanishing cohomology in the odd dimensions. The study of such manifolds was begun by Stefan Müller. In order to understand the structure of these manifolds, we will analyze their minimal handle presentations and describe explicitly to what extent these handle presentations are determined by the cohomology ring and the characteristic classes. It turns out that the cohomology ring and the characteristic classes do not suffice to reconstruct a manifold of the above type completely. In fact, the group $\text{Aut}_0(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0(\#_{i=1}^b(S^2 \times D^6))$ of automorphisms of $\#_{i=1}^b(S^2 \times S^5)$ which induce the identity on cohomology modulo those which extend to $\#_{i=1}^b(S^2 \times D^6)$ acts on the set of oriented homeomorphism classes of manifolds with fixed cohomology ring and characteristic classes, and we will be also concerned with describing this group and some facts about the above action.

1. INTRODUCTION

The classification of topological manifolds up to homeomorphism is an extremely interesting and important problem. Let us restrict our attention to the case of closed (i.e., compact without boundary) and oriented simply connected manifolds. As a general classification scheme, surgery theory [1] solves this problem for manifolds within a given homotopy type, e.g., that of a sphere. Another approach to the classification “up to finite indeterminacy”, using rational homotopy theory, is due to Sullivan [34]. Nevertheless, there are only a few explicit results which characterize the oriented homeomorphism type of a manifold in terms of easily computable invariants. They usually require many simplifying assumptions, such as high connectedness [36]. In this paper, we will

consider *even cohomology manifolds* (or *E-manifolds*, for short) in dimension eight, by which we mean closed, oriented, simply connected, piecewise linear or smooth manifolds all of whose odd-dimensional homology groups with integer coefficients vanish. The universal coefficient theorem implies that all homology groups of an E-manifold are without torsion. Moreover, since $H^3(X, \mathbf{Z}_2) = 0$ for an E-manifold, two E-manifolds of dimension at least 6 are homeomorphic (as topological manifolds) if and only if they are piecewise linearly homeomorphic [16].

Though the class of E-manifolds is fairly restricted, it still contains many interesting examples from various areas of mathematics, such as the piecewise linear manifolds underlying the toric manifolds from Algebraic Geometry [4] or the polygon spaces [12], to mention a few. So far, E-manifolds have been classified up to dimension 6. Of course, in dimension 2 there is only S^2 , in dimension 4, there is the famous classification result of Freedman various interesting aspects of which are discussed in [15], and finally in dimension 6, the classification was achieved by Wall [37] and Jupp [14]. Various applications of the latter result to Algebraic Geometry are surveyed in [26]. Finally, we refer to [2], [3], and [29] for the determination of projective algebraic structures on certain 6- and 8-dimensional E-manifolds.

ACKNOWLEDGEMENTS. My thanks go to J.-C. Hausmann for his interest in this work and for pointing out to me that an E-manifold of dimension eight is actually not determined by its classical invariants.

2. STATEMENT OF THE RESULT

We now discuss the main result of this note, namely the classification of E-manifolds of dimension 8 with vanishing second Stiefel-Whitney class in the form of an exact sequence of pointed sets. This result was motivated by the work [24]. In order to state it in a more elegant form, we will work with based manifolds. By a *based piecewise linear (smooth) E-manifold*, we mean a triple $(X, \underline{x}, \underline{y})$, consisting of a piecewise linear (smooth) E-manifold X and bases $\underline{x} = (x_1, \dots, x_{b_2(M)})$ for $H^2(X, \mathbf{Z})$ and $\underline{y} = (y_1, \dots, y_{b_4(M)})$ for $H^4(X, \mathbf{Z})$. Recall that by definition E-manifolds are oriented, so that the above data specify a basis for $H^*(X, \mathbf{Z})$, such that the bases for $H^i(X, \mathbf{Z})$ and $H^{8-i}(X, \mathbf{Z})$ are dual to each other with respect to the cup product. An *isomorphism between piecewise linear (smooth) based E-manifolds* $(X, \underline{x}, \underline{y})$ and $(X', \underline{x}', \underline{y}')$ is an orientation preserving piecewise linear (smooth) isomorphism $f: X \rightarrow X'$ with $f^*(\underline{x}') = \underline{x}$

and $f^*(\underline{y}') = \underline{y}$. We denote by $\mathfrak{J}^{\text{PL}(C^\infty)}(b, b')$ the set of isomorphy classes of piecewise linear (smooth) based E-manifolds $(X, \underline{x}, \underline{y})$ of dimension eight with vanishing second Stiefel-Whitney class, $b_2(X) = b$, and $b_4(X) = b'$.

2.1 THE CLASSICAL INVARIANTS

In the terminology of [24], the classical invariants of an E-manifold consist of its cohomology ring, the Stiefel-Whitney classes, the Wu classes, the Pontrjagin classes, the Euler class, the Steenrod squares, the reduced Steenrod powers, and the Pontrjagin powers. For an eight-dimensional E-manifold X with vanishing second Stiefel-Whitney class, the main result of [24] states that the classical invariants are fully determined by the following invariants:

1. The cup product map

$$\begin{aligned} \delta_X: S^2H^2(X, \mathbf{Z}) &\longrightarrow H^4(X, \mathbf{Z}) \\ x \otimes x' &\longmapsto x \cup x'. \end{aligned}$$

2. The intersection form

$$\begin{aligned} \gamma_X: S^2H^4(X, \mathbf{Z}) &\longrightarrow \mathbf{Z} \\ y \otimes y' &\longmapsto (y \cup y')[X]. \end{aligned}$$

Here, $[X] \in H_8(X, \mathbf{Z})$ is the fundamental class determined by the orientation.

3. The first Pontrjagin class $p_1(X) \in H^4(X, \mathbf{Q})$.

REMARK 2.1. The above invariants are not independent. By associativity of the cohomology ring, the following relation holds

$$(1) \quad \delta_X^*(\gamma_X) \in S^4H^2(X, \mathbf{Z})^\vee,$$

i.e.,

$$\gamma_X(\delta_X(x_1 \otimes x_2) \otimes \delta_X(x_3 \otimes x_4)) = \gamma_X(\delta_X(x_1 \otimes x_3) \otimes \delta_X(x_2 \otimes x_4)),$$

for all $x_1, x_2, x_3, x_4 \in H^2(X, \mathbf{Z})$. Furthermore,

PROPOSITION ([24], Prop. A.7 or Cor. 3.14 below). *For every element $y \in H^4(X, \mathbf{Z})$ we have*

$$(2) \quad p_1(X)y \equiv 2y^2 \pmod{4}.$$

Note that this implies $p_1(X) \in H^4(X, \mathbf{Z})$. If, in addition, X is differentiable then for every integral lift $W \in H^2(X, \mathbf{Z})$ of $w_2(X)$ one has

$$(3) \quad 3p_1(X)^2 - 14p_1(X)W^2 + 7W^4 \equiv 12 \operatorname{Sign}(\gamma_X) \pmod{2688}.$$

Müller has also shown [24] that these relations imply all the other relations among the classical invariants of X . Conversely, a piecewise linear manifold X whose invariants obey relation (3) admits a differentiable structure [18], [24].

We are led to the following algebraic concept: A *system of invariants of type (b, b')* is a triple (δ, γ, p) , consisting of

- a homomorphism $\delta: S^2\mathbf{Z}^{\oplus b} \rightarrow \mathbf{Z}^{\oplus b'}$,
- a unimodular symmetric bilinear form $\gamma: S^2\mathbf{Z}^{\oplus b'} \rightarrow \mathbf{Z}$, and
- an element $p \in \mathbf{Z}^{\oplus b'}$.

We denote by $Z(b, b')$ the set of systems of invariants of type (b, b') .

Now, let $(X, \underline{x}, \underline{y})$ be a based eight-dimensional E-manifold. This defines a set of invariants $Z_{(X, \underline{x}, \underline{y})} := (\delta_X, \gamma_X, p_1(X))$ of type $(b_2(X), b_4(X))$. Thus, we have natural maps

$$\begin{aligned} Z^{\text{PL}(C^\infty)}(b, b') &: \mathfrak{J}^{\text{PL}(C^\infty)}(b, b') \longrightarrow Z(b, b') \\ [X, \underline{x}, \underline{y}] &\longmapsto Z_{(X, \underline{x}, \underline{y})}. \end{aligned}$$

It will be the concern of our paper to understand the maps $Z^{\text{PL}(C^\infty)}$ as well as possible. The first result can be easily derived from Wall's work [36] and deals with the case $b = 0$. It will be proved in detail in Section 4.1.

THEOREM 2.2. i) *The map $Z^{\text{PL}}(0, b')$ is injective. Its image consists precisely of those elements which satisfy the relations (1) and (2).*

ii) *Given two smooth based E-manifolds (X, \underline{y}) and (X', \underline{y}') with $b_2(X) = 0 = b_2(X')$ and $Z_{(X, \underline{y})} = Z_{(X', \underline{y}')}$, there exists an exotic 8-sphere Σ such that $(X \# \Sigma, \underline{y})$ and (X', \underline{y}') are smoothly isomorphic. In particular, the fibres of Z^{C^∞} have cardinality at most two. The image of Z^{C^∞} consists exactly of those elements which satisfy the relations (1), (2), and (3).*

2.2 MANIFOLDS WITH TRIVIAL CUP FORM δ_X

In addition to describing the explicit geometric meaning of the system of invariants of an E-manifold X with $w_2(X) = 0$, we will describe those manifolds X for which the cup form δ_X vanishes.

For any $b > 0$, let $\mathcal{FC}_b^{\text{PL}(C^\infty)}$ be the group of isotopy classes of piecewise linear (smooth) embeddings of b disjoint copies of $S^5 \times D^3$ into S^8 . The following result will be established in Section 3.7.

PROPOSITION 2.3. $\text{FL}_b := \mathcal{FC}_b^{C^\infty} \cong \mathcal{FC}_b^{\text{PL}}$.

Given an element $[l] \in \text{FL}_b$, we can perform surgery along the link l and get a smooth based E-manifold $(X(l), \underline{x}(l))$ with $w_2(X) = 0$ and $b_4(X) = 0$, which is well defined up to smooth isomorphism of based manifolds.

We will also use the following notation: Fix a pair $(\gamma, p) \in Z(0, b')$ which satisfies relation (2) (and (3)) and denote by $\mathcal{J}^{\text{PL}(C^\infty)}(b, \gamma, p)$ the set of isomorphism classes of based piecewise linear (smooth) E-manifolds $(X, \underline{x}, \underline{y})$ with $w_2(X) = 0$, $b_2(X) = b$, $\gamma_X = \gamma$, and $p_1(X) = p$. Pick a three-connected piecewise linear (smooth) based E-manifold (X_0, \underline{y}_0) with $\gamma_{X_0} = \gamma$ and $p_1(X_0) = p$. In the smooth case, let $\vartheta^8 \cong \mathbf{Z}_2$ [17] be the group of exotic 8-spheres, and set $\vartheta(X_0) := \vartheta^8$, if X_0 is not diffeomorphic to $X_0 \# \Sigma$, Σ a generator for ϑ^8 , and $\vartheta(X_0) := \{[S^8]\} \subset \vartheta^8$ otherwise. Now, we define maps

$$K^{\text{PL}}(b, \gamma, p): \text{FL}_b \longrightarrow \mathcal{J}^{\text{PL}}(b, \gamma, p)$$

$$[l] \longmapsto [X(l) \# X_0, \underline{x}(l), \underline{y}_0]$$

and

$$K_{X_0}^{C^\infty}(b, \gamma, p): \text{FL}_b \oplus \vartheta(X_0) \longrightarrow \mathcal{J}^{C^\infty}(b, \gamma, p)$$

$$([l], [\Sigma]) \longmapsto [X(l) \# X_0 \# \Sigma, \underline{x}(l), \underline{y}_0].$$

In $\mathcal{J}^{\text{PL}(C^\infty)}(b, \gamma, p)$, we mark the class $[(\#_{i=1}^b (S^2 \times S^6)) \# X_0, \underline{x}, \underline{y}_0]$, where \underline{x} comes from the natural basis of $H^2(\#_{i=1}^b (S^2 \times S^6), \mathbf{Z})$. Then our main result is the following

THEOREM 2.4. i) For every $b > 0$ and every pair (γ, p) which satisfies the relation (2),

$$\begin{array}{ccccc} \{1\} & \longrightarrow & \text{FL}_b & \xrightarrow{K^{\text{PL}}(b, \gamma, p)} & \mathfrak{J}^{\text{PL}}(b, \gamma, \varphi) & \longrightarrow & \text{Hom}(S^2 \mathbf{Z}^b, \mathbf{Z}^{b'}) \\ 1 & \longmapsto & [\text{trivial link}] & & [X, \underline{x}, \underline{y}] & \longmapsto & \delta_X \end{array}$$

is an exact sequence of pointed sets.

ii) For every $b > 0$ and every pair (γ, p) which satisfies the relations (2) and (3),

$$\begin{array}{ccccc} \{1\} & \longrightarrow & \text{FL}_b \oplus \mathcal{V}(X_0) & \xrightarrow{K_{x_0}^{\text{C}\infty}(b, \gamma, p)} & \mathfrak{J}^{\text{C}\infty}(b, \gamma, \varphi) & \longrightarrow & \text{Hom}(S^2 \mathbf{Z}^b, \mathbf{Z}^{b'}) \\ 1 & \longmapsto & [\text{trivial link}] & & [X, \underline{x}, \underline{y}] & \longmapsto & \delta_X \end{array}$$

is an exact sequence of pointed sets.

The proof will be given in Sections 4.2 and 4.3.

REMARK 2.5. i) In the PL setting, we will show that the inclusion of FL_b into $\mathfrak{J}^{\text{PL}}(b, \gamma, \varphi)$ extends to an action of FL_b on $\mathfrak{J}^{\text{PL}}(b, \gamma, \varphi)$, such that the orbits are the fibres of the map $[X, \underline{x}, \underline{y}] \longmapsto \delta_X$.

ii) On all the sets occurring in Theorem 2.4 there are natural $(\text{GL}_b(\mathbf{Z}) \times \text{GL}_{b'}(\mathbf{Z}))$ -actions, and the maps are equivariant for these actions. Therefore, by forming the equivalence classes with respect to these actions, we get the classification of E-manifolds with vanishing second Stiefel-Whitney class up to orientation preserving piecewise linear (smooth) isomorphy.

iii) We will discuss in Section 3.7 the structure of the group FL_b . It turns out that it is finite if and only if $b = 1$. It follows easily that the set of $\text{GL}_b(\mathbf{Z})$ -equivalence classes in FL_b is infinite for $b \geq 2$. Thus, the cohomology ring and the characteristic classes classify E-manifolds of dimension eight up to finite indeterminacy only if the second Betti number is at most one.

The starting point of our proof of the above results will be the theory of minimal handle decompositions of Smale which states that X can be obtained from D^8 by first attaching $b_2(X)$ 2-handles, then $b_4(X)$ 4-handles, then $b_6(X)$ 6-handles and finally one 8-handle. At each step, the attachment will be determined by the isotopy class of a certain framed link in a 7-manifold, and we will first explain how to read off the isotopy class of the attaching links for the 2- and 4-handles from the invariants.

3. PRELIMINARIES

We collect in this paragraph the background material and some preliminary results which we will use in our proof. Most of the results are by now standard results from various parts of algebraic, differential, and piecewise linear topology.

3.1 THE STRUCTURE OF MANIFOLDS: HANDLE ATTACHMENT AND SURGERY

Let M be an m -dimensional manifold with boundary. Suppose we are given an embedding $f: S^{\lambda-1} \times D^{m-\lambda} \rightarrow \partial M$. We then define

$$M' := M \cup_f (D^\lambda \times D^{m-\lambda})$$

and say that M' is obtained from M by the *attachment of a λ -handle along f* . Moreover, $f(S^{\lambda-1} \times \{0\})$ is called the *attaching sphere*, $D^\lambda \times \{0\}$ the *core disc*, and $\{0\} \times S^{m-\lambda-1}$ the *belt sphere*. We will often simply write $M' = M \cup H^\lambda$.

REMARK 3.1. i) If M is assumed to be differentiable and f to be a differentiable embedding, handle attachment can be described in such a way that the resulting manifold is again differentiable (see [17], VI, §§6 and 8), i.e., no “smoothing of the corners” is required.

ii) If M is oriented, then M' will inherit an orientation which is compatible with the given orientation of M and the natural orientation of $D^\lambda \times D^{m-\lambda}$, if and only if f *reverses* the orientations.

The next operation we consider was introduced by Milnor [21] and Wallace [38] and goes back to Thom. For this, let N be a manifold of dimension $m-1$ and $f: S^{\lambda-1} \times D^{m-\lambda} \rightarrow N$ an embedding. Denote by \bar{f} the restriction of f to $S^{\lambda-1} \times S^{m-\lambda-1}$, and set

$$\chi(N, f) := (N \setminus \text{int } f(S^{\lambda-1} \times D^{m-\lambda})) \cup_{\bar{f}} (D^\lambda \times S^{m-\lambda-1}).$$

We say that $\chi(N, f)$ is constructed from N by *surgery along f* . Informally speaking, we remove from N a $(\lambda-1)$ -sphere with trivial normal bundle and replace it with an $(m-\lambda-1)$ -sphere, again with trivial normal bundle.

REMARK 3.2. i) If N is oriented, then f has to be orientation preserving for $\chi(N, f)$ to inherit a natural orientation from those of N and $D^\lambda \times S^{m-\lambda-1}$. This is because $S^{\lambda-1} \times S^{m-\lambda-1}$ inherits the reversed orientation as boundary of $N \setminus \text{int } f(S^{\lambda-1} \times D^{m-\lambda})$.

ii) The operations of handle attachment and surgery are closely related: Let M be an m -dimensional manifold with boundary $N := \partial M$ and $f: S^{\lambda-1} \times D^{m-\lambda} \rightarrow N$ an embedding. Now, attach a λ -handle along f in order to obtain M' . Then $\partial M' = \chi(N, f)$.

We will also perform a “surgery in pairs”. For this, N is assumed to be an $(m-1)$ -dimensional manifold, and K a submanifold of dimension $k-1$. Assume that, for some $\lambda \leq k$, we are given an embedding $f: S^{\lambda-1} \times D^{m-\lambda} \rightarrow N$ which induces an embedding $f^* := f|_{S^{\lambda-1} \times D^{k-\lambda}}: S^{\lambda-1} \times D^{k-\lambda} \rightarrow K$. Then $\chi(K, f^*)$ is naturally contained as a submanifold in $\chi(N, f)$.

The next result is a special case of the “minimal presentation theorem” of Smale [31] and is crucial for the explicit analysis of the structure of a manifold.

THEOREM 3.3. *Let X be a closed simply connected manifold of dimension $m \geq 6$ with torsion free homology. Then there exists a sequence of submanifolds*

$$D^m \cong W_0 \subset W_1 \subset W_2 \subset \cdots \subset W_m = X,$$

such that W_i is obtained from W_{i-1} by attaching $b_i(X)$ i -handles, $i = 1, \dots, m$.

Moreover, for any such sequence, there exists a dual sequence

$$\bar{W}_0 \subset \bar{W}_1 \subset \cdots \subset \bar{W}_m = X,$$

such that the attaching $(i-1)$ -spheres in \bar{W}_{i-1} coincide with the belt spheres in W_{m-i} , $i = 1, \dots, m$.

Proof. For differentiable manifolds, an attractive presentation of the relevant material is contained in Chapters VII and VIII of [17]. In the piecewise linear category, handle decompositions are discussed in [27] (cf. also [13]). However, the statement concerning the number of handles is not explicitly given there. Nevertheless, one verifies that the necessary tools (such as Whitney lemma and handle sliding) are also proved in [27]. \square

REMARK 3.4. i) Retracting all λ -handles to their core discs, starting with $\lambda = 0$, yields a CW-complex which is homotopy equivalent to X (cf. [27], p. 83).

ii) Observe that, by i), a handle decomposition as in Theorem 3.3 yields a *preferred basis* for $H_*(X, \mathbf{Z})$. By renumbering, orientation reversal in the attaching spheres, and handle sliding, one can obtain any basis of $H_*(X, \mathbf{Z})$ as the preferred basis of a handle decomposition ([17], (1.7), p. 148).

iii) If X comes with an orientation, we may assume that D^m is orientation preservingly embedded and that all attaching maps are orientation reversing.

3.2 CONSEQUENCES FOR E-MANIFOLDS OF DIMENSION EIGHT WITH $w_2 = 0$

Let X be a piecewise linear (smooth) E-manifold of dimension eight with $w_2(X) = 0$. The first observation concerns the structure of W_2 .

LEMMA 3.5. *One has $W_2 \cong \#_{i=1}^b(S^2 \times D^6)$.*

Proof. The manifold W_2 is an $(8, 1)$ -handle body and as such homeomorphic to the boundary connected sum of b D^6 -bundles over S^2 ([17], §11, p.115). As $\pi_1(\text{SO}(4)) \cong \mathbf{Z}_2$ and we have requested $w_2(X) = 0$, the claim follows. \square

The next consequence is

The manifold W_4 is determined by a framed link of $b_4(X)$ three-dimensional spheres in $\#_{i=1}^b(S^2 \times S^5)$.

We shall look into the classification of such links below. The third consequence is

LEMMA 3.6. i) $\partial W_4 \cong \#_{i=1}^b(S^2 \times S^5)$.

ii) *The manifold X is of the form $W_4 \cup_f \#_{i=1}^b(S^2 \times D^6)$ where*

$$f: \#_{i=1}^b(S^2 \times S^5) \longrightarrow \partial W_4$$

is a piecewise linear (smooth) isomorphism, such that f_ maps the canonical basis of $H_2(\#_{i=1}^b(S^2 \times S^5), \mathbf{Z})$ to the preferred basis of $H_2(\partial W_4, \mathbf{Z})$.*

Proof. i) This follows because $\partial W_4 \cong \partial \bar{W}_2$. ii) follows because $X = W_4 \cup \bar{W}_2$, and $\bar{W}_2 \cong \#_{i=1}^b(S^2 \times D^6)$, by Lemma 3.5. \square

3.3 HOMOTOPY VS. ISOTOPY

By Theorem 3.3, the manifold is determined by the ambient isotopy classes of the attaching maps. However, the topological invariants of the manifold give us at best their homotopy classes. It is, therefore, important to have theorems granting that this is enough. In the setting of differentiable manifolds, we have

THEOREM 3.7 (Haefliger [6], [7]). *Let S be a closed differentiable manifold of dimension n and M an m -dimensional differentiable manifold without boundary. Let $f: S \rightarrow M$ be a continuous map and $k \geq 0$, such that*

$$\pi_i(f): \pi_i(S) \rightarrow \pi_i(M)$$

is an isomorphism for $0 \leq i \leq k$ and surjective for $i = k + 1$. Then the following is satisfied:

1. *If $m \geq 2n - k$ and $n > 2k + 2$, then f is homotopic to a differentiable embedding.*
2. *If $m > 2n - k$ and $n \geq 2k + 2$, then two differentiable embeddings of S into M which are homotopic are also ambient isotopic.*

In the setting of piecewise linear manifolds, similar results hold true. We refer to Haefliger's survey article [9]. For our purposes, the result stated below will be sufficient.

THEOREM 3.8. *Suppose S is a closed n -dimensional manifold, M an m -dimensional manifold without boundary, and $f: S \rightarrow M$ a continuous map. Assume one has*

- $m - n \geq 3$;
- S is $(2n - m + 1)$ -connected;
- M is $(2n - m + 2)$ -connected.

Then:

- (1) *f is homotopic to an embedding;*
- (2) *two embeddings which are homotopic to f are ambient isotopic.*

Proof. The theorem of Irwin ([27], Thm. 7.12 and Ex. 7.14, [13], Thm. 8.1) yields (1) and that f_1 and f_2 as in (2) are concordant. But, since $m - n \geq 3$, concordance implies ambient isotopy ([13], Chap. IX). \square

COROLLARY 3.9. *Let $S := S^3$ and M a simply connected differentiable or piecewise linear manifold of dimension 7 without boundary. Then $\pi_3(M)$ classifies differentiable and piecewise linear embeddings, respectively, of S^3 into M up to ambient isotopy.*

COROLLARY 3.10 (Zeeman's unknotting theorem [39]). *For every m, n with $m - n \geq 3$, any piecewise linear embedding of S^n into S^m is isotopic to the standard embedding.*

3.4 SOME 4-DIMENSIONAL CW-COMPLEXES

By Remark 3.4, a handle decomposition of X gives us a CW-complex which is homotopy equivalent to X . The following discussion will enable us to understand the 4-skeleton of that complex.

Let $W := S^2 \vee \dots \vee S^2$ be the b -fold wedge product of 2-spheres. Suppose X is the CW-complex obtained by attaching a 4-cell to W via the map $g \in \pi_3(W)$. The Hilton-Milnor theorem ([30], Thm. 7.9.4) asserts that

$$\pi_3(W) = \bigoplus_{i=1}^b \pi_3(S^2) \oplus \bigoplus_{1 \leq i < j \leq b} \pi_3(S^3).$$

Choosing the standard generators for $\pi_3(S^2)$ and $\pi_3(S^3)$, we can describe g by a tuple $(l_i, i = 1, \dots, b; l_{ij}, 1 \leq i < j \leq b)$ of integers. These integers determine the cohomology ring of $X = W \cup_g D^4$ as follows:

PROPOSITION 3.11. *Let $y \in H^4(X, \mathbf{Z})$ be the generator of $H^4(X, \mathbf{Z})$ given by the attached 4-cell and x_1, \dots, x_b the canonical basis of $H^2(X, \mathbf{Z}) = H^2(W, \mathbf{Z})$. Then*

$$\begin{aligned} x_i \cup x_j &= l_{ij} \cdot y, & 1 \leq i < j \leq b, \\ x_i \cup x_i &= l_i \cdot y, & i = 1, \dots, b. \end{aligned}$$

This is proved like [22], (1.5), p. 103. We recall the proof in the following example.

EXAMPLE 3.12. We treat the case $b = 2$. Consider the embedding

$$\iota: S^2 \vee S^2 \hookrightarrow S^2 \times S^2 \hookrightarrow \mathbf{CP}^\infty \times \mathbf{CP}^\infty.$$

The standard basis for $H^4(\mathbf{CP}^\infty \times \mathbf{CP}^\infty, \mathbf{Z}) \cong \mathbf{Z}^{\oplus 3}$ is given by the elements y_1, y_2, y_3 obtained from attaching D^4 via $(1, 0; 0)$, $(0, 0; 1)$, and $(0, 1; 0)$, respectively. Let $h: D^4 \rightarrow D^4 \vee D^4 \vee D^4$ be the canonical map followed by

$$(\vartheta \cdot x \mapsto \vartheta \cdot m_{l_1}(x)) \vee (\vartheta \cdot x \mapsto \vartheta \cdot m_{l_{12}}(x)) \vee (\vartheta \cdot x \mapsto \vartheta \cdot m_{l_2}(x)).$$

Here, m_k stands for a representative of $[k \cdot \text{id}_{S^3}] \in \pi_3(S^3)$ and $D^4 = \{ \vartheta \cdot x \mid x \in S^3, \vartheta \in [0, 1] \}$. Now, h and ι glue to a map $f: X \rightarrow \mathbf{CP}^\infty \times \mathbf{CP}^\infty$, and

$$\begin{aligned} f^*: H^4(\mathbf{CP}^\infty \times \mathbf{CP}^\infty, \mathbf{Z}) &\rightarrow H^4(X, \mathbf{Z}) \\ a_1 y_1 + a_2 y_2 + a_3 y_3 &\mapsto (a_1 l_1 + a_2 l_{12} + a_3 l_2) y, \end{aligned}$$

so that the assertion follows from the naturality of the cup-product.

3.5 PONTRJAGIN CLASSES AND $\pi_3(\text{SO}(4))$

Vector bundles of rank 4 over S^4 are classified by elements in $\pi_3(\text{SO}(4))$. In our setting, such vector bundles will appear as normal bundles. We recall, therefore, the description of that group and relate it to Pontrjagin classes and self intersection numbers.

First, look at the natural map $\pi_3(\text{SO}(4)) \longrightarrow \pi_3(\text{SO}(4)/\text{SO}(3)) = \pi_3(S^3)$. This map has a splitting ([32], §22.6) which induces an isomorphism

$$\pi_3(\text{SO}(4)) = \pi_3(\text{SO}(3)) \oplus \pi_3(S^3).$$

Let α_3 be the generator for $\pi_3(\text{SO}(3)) \cong \mathbf{Z}$ from [32], §22.3, and $\beta_3 := [\text{id}_{S^3}] \in \pi_3(S^3)$, so that we obtain the isomorphism $\mathbf{Z} \oplus \mathbf{Z} \longrightarrow \pi_3(\text{SO}(4))$, $(k_1, k_2) \longmapsto k_1\alpha_3 + k_2\beta_3$. Finally, the kernel of the map $\pi_3(\text{SO}(4)) \longrightarrow \pi_3(\text{SO})$ to the stable homotopy group is generated by $-\alpha_3 + 2\beta_3$ ([32], §23.6), whence [23], (20.9), implies

PROPOSITION 3.13. *Let E be the vector bundle over S^4 defined by the element $k_1\alpha_3 + k_2\beta_3 \in \pi_3(\text{SO}(4))$. Then*

$$p_1(E) = \pm(2k_1 + 4k_2).$$

COROLLARY 3.14. *Let $f: S^4 \longrightarrow M$ be a differentiable embedding of S^4 into the differentiable 8-manifold M . Let $E := f^*T_M/T_{S^4}$ be the normal bundle. Then the self intersection number s of $f(S^4)$ in M satisfies*

$$2s \equiv p_1(E) \pmod{4}.$$

Proof. If E is given by the element $k_1\alpha_3 + k_2\beta_3 \in \pi_3(\text{SO}(4))$, then $s = k_2$ ([17], (5.4), p.72). Since $p_1(E) = \pm(2k_2 + 4k_1)$, the claim follows. \square

3.6 LINKS OF 3-SPHERES IN $\#_{i=1}^b(S^2 \times S^5)$

If X is a closed E-manifold of dimension 8 with $w_2(X) = 0$, then $W_2 := \#_{i=1}^b(S^2 \times D^6)$, $b = b_2(X)$, by Lemma 3.5. Thus, W_4 is determined by a framed link of 3-spheres in $\partial W_2 = \#_{i=1}^b(S^2 \times S^5)$. Therefore, we will now classify such links.

So, let $W := \#_{i=1}^b(S^2 \times S^5)$ be a b -fold connected sum. We can choose b disjoint 2-spheres S_i^2 , $i = 1, \dots, b$, embedded in W and representing the natural basis of $H_2(W, \mathbf{Z})$. One checks that the homotopy type of W is given up to dimension 4 by the b -fold wedge product $S^2 \vee \dots \vee S^2$. Suppose we are given a link of b' three-dimensional spheres, i.e., we are given b' differentiable embeddings $g_i: S^3 \longrightarrow W$, $i = 1, \dots, b'$, with $g_i(S^3) \cap g_j(S^3) = \emptyset$ for $i \neq j$.

By the transversality theorem ([17], IV.(2.4)), one sees that we may assume $S_i^2 \cap g_j(S^3) = \emptyset$ for all i and j .

By Corollary 3.9, the ambient isotopy class of the embedding g_k is determined by the element $\varphi_k := [g_k] \in \pi_3(W_k)$, $W_k := W \setminus \bigcup_{j \neq k} g_j(S^3)$, $k = 1, \dots, b'$. We clearly have (compare [8])

$$\pi_3(W_k) = \pi_3\left(\underbrace{S^2 \vee \dots \vee S^2}_{b \times} \vee \underbrace{S^3 \vee \dots \vee S^3}_{(b'-1) \times}\right),$$

so that the Hilton-Milnor theorem yields

$$\pi_3(W_k) = \bigoplus_{i=1}^b \pi_3(S^2) \oplus \bigoplus_{1 \leq i < j \leq b} \pi_3(S^3) \oplus \bigoplus_{j \neq k} \pi_3(S^3).$$

Hence, we write φ_k as a tuple of integers:

$$\varphi_k = (l_i^k, i = 1, \dots, b; l_{ij}^k, 1 \leq i < j \leq b; \lambda_{kj}, j \neq k).$$

Observe that, for $j \neq k$, φ_k is mapped under the natural homomorphism

$$\pi_3(W_k) \longrightarrow H_3(W_k, \mathbf{Z}) \longrightarrow H_3(W \setminus g_j(S^3), \mathbf{Z}) (\cong \mathbf{Z})$$

to the image of the fundamental class of S^3 under g_{j*} . Thus, λ_{kj} is just the ‘usual’ linking number of the spheres $g_k(S^3)$ and $g_j(S^3)$ in W (compare [8]).

3.7 LINKS OF 5-SPHERES IN S^8

Let $\mathcal{FC}_b^{\text{PL}(C^\infty)}$ be as before, and let $\mathcal{C}_b^{\text{PL}(C^\infty)}$ be the group of isotopy classes of piecewise linear (smooth) embeddings of b disjoint copies of S^5 into S^8 . For $b = 1$, these groups are studied in [10], [19], and [20]. A brief summary with references of results in the case $b > 1$ is contained in Section 2.6 of [11]. We will review some of this material below.

PROPOSITION 3.15. *We have $\mathcal{FC}_1^{C^\infty} \cong \mathcal{FC}_1^{\text{PL}} \cong \mathbf{Z}_2$.*

Proof. Since $\pi_5(\text{SO}(3)) \cong \mathbf{Z}_2$, the standard embedding of S^5 into S^8 with its two possible framings provides an injection of \mathbf{Z}_2 into $\mathcal{FC}_1^{\text{PL}(C^\infty)}$. By Zeeman’s unknotting theorem 3.10, the map $\mathbf{Z}_2 \longrightarrow \mathcal{FC}_1^{\text{PL}}$ is an isomorphism. As remarked in Section 2.6 of [11], $\mathcal{FC}_1^{\text{PL}}$ is isomorphic to $\mathcal{F}\vartheta$, the group of h-cobordism classes of framed submanifolds of S^8 which are homotopy 5-spheres. Moreover, by [10] and [19], there is an exact sequence

$$\dots \longrightarrow \vartheta^6 \longrightarrow \mathcal{FC}_1^{C^\infty} \longrightarrow \mathcal{F}\vartheta \longrightarrow \vartheta^5 \longrightarrow \dots$$

As the groups ϑ^5 and ϑ^6 of exotic 5- and 6-spheres are trivial [17], our claim is settled. \square

Let $L_b \subset C_b^{C^\infty}$ be the subgroup of those embeddings for which the restriction to each component is isotopic to the standard embedding. As observed in Section 2.6 of [11], Zeeman's unknotting theorem 3.10 implies that $L_b = C_b^{PL}$. The following result settles Proposition 2.3:

$$\text{COROLLARY 3.16. } \mathcal{FC}_b^{C^\infty} \cong \mathcal{FC}_b^{PL} \cong L_b \oplus \bigoplus_{i=1}^b \mathbf{Z}_2.$$

For the group L_b , Theorem 1.3 of [11] provides a fairly explicit description as an extension of abelian groups. For this, consider the b -fold wedge product $\bigvee_{i=1}^b S^2$ of 2-spheres together with its inclusion $i: \bigvee_{i=1}^b S^2 \hookrightarrow \times_{i=1}^b S^2$ into the b -fold product of 2-spheres. Finally, let $p_i: \bigvee_{i=1}^b S^2 \rightarrow S^2$ be the projection onto the i^{th} factor, $i = 1, \dots, b$. Set, for $m = 1, 2, \dots$,

$$\Lambda_{b,j}^m := \text{Ker}(\pi_m(p_j): \pi_m(\bigvee_{i=1}^b S^2) \rightarrow \pi_m(S^2)), \quad j = 1, \dots, b,$$

$$\Lambda_b^m := \bigoplus_{j=1}^b \Lambda_{b,j}^m$$

and

$$\Pi_b^m := \text{Ker}(\pi_m(i): \pi_m(\bigvee_{i=1}^b S^2) \rightarrow \bigoplus_{i=1}^b \pi_m(S^2)),$$

and define

$$w_b^m: \Lambda_b^m \rightarrow \Pi_b^{m+1}$$

on $\Lambda_{b,j}^m$ by $w_b^m(\alpha) := [\alpha, \iota_i]$. Here, $[\cdot, \cdot]$ stands for the Whitehead product inside the homotopy groups of $\bigvee_{i=1}^b S^2$ and $\iota_i: S^2 \hookrightarrow \bigvee_{i=1}^b S^2$ for the inclusion of the i^{th} factor, $i = 1, \dots, b$. Theorem 1.3 of [11] yields in our situation

THEOREM 3.17. *There is an exact sequence of abelian groups*

$$0 \rightarrow \text{Coker}(w_b^6) \rightarrow L_b \rightarrow \text{Ker}(w_b^5) \rightarrow 0.$$

We remark that the formulas of Steer [33] might be used for the explicit computation of Whitehead products and thus for the determination

of $\text{Coker}(w_b^6)$ and $\text{Ker}(w_b^5)$. The free part of L_b , e.g., can be obtained quite easily. We confine ourselves to prove the following important fact.

COROLLARY 3.18. *The group L_b has positive rank for $b \geq 2$.*

Proof. Let $L_b := \bigoplus_{l \geq 1} L_{b,l}$ be the free graded Lie algebra with $L_{b,1} := \bigoplus_{i=1}^b \mathbf{Z} \cdot e_i$. For $l = 2, 3, \dots$, let $e_1^l, \dots, e_{d_l}^l$ be a basis for $L_{b,l}$ consisting of iterated commutators of the e_i . By assigning ι_i to e_i , every iterated commutator $c \in L_{b,l}$ in the e_i defines an element $\alpha(c) \in \pi_{l+1}(\bigvee_{i=1}^b S^2)$.

To settle our claim, it is certainly sufficient to show that $\text{Coker}(w_b^6)$ has positive rank. Now, by the Hilton-Milnor theorem

$$\Pi_b^7 \cong \bigoplus_{l=3}^7 \bigoplus_{k=1}^{d_{l-1}} \pi_7(S^l) \cdot \alpha(e_k^{l-1}).$$

Note that $\pi_7(S^l)$ is finite for $l \notin \{4, 7\}$ (see [32] and [35] for the explicit description of those groups). The Hopf fibration $S^7 \rightarrow S^4$ [32], on the other hand, yields a decomposition $\pi_7(S^4) \cong \pi_6(S^3) \oplus \pi_7(S^7) \cong \mathbf{Z}_{12} \oplus \mathbf{Z}$. Therefore, it will suffice to show that the free part of Λ_b^6 is mapped to $\bigoplus_{j=1}^{d_6} \pi_7(S^7) \cdot \alpha(e_j^6)$. For $j = 1, \dots, b$, we have

$$\Lambda_{b,j}^6 \cong \bigoplus_{i \neq j} \pi_6(S^2) \cdot \iota_i \oplus \bigoplus_{l=3}^6 \bigoplus_{k=1}^{d_{l-1}} \pi_6(S^l) \cdot \alpha(e_k^{l-1}).$$

The group $\pi_6(S^l)$ is finite for $l < 6$, and we obviously have $[\alpha(e_k^5), \iota_j] = \alpha([e_k^5, e_j])$. If we expand the commutator $[e_k^5, e_j]$ in the basis $e_1^6, \dots, e_{d_6}^6$, we find an expansion for $[\alpha(e_k^6), \iota_j]$ in terms of the $\alpha(e_k^6)$. \square

COROLLARY 3.19. *The set of $\text{GL}_b(\mathbf{Z})$ -equivalence classes of elements in L_b is infinite for $b \geq 2$.*

Proof. We have seen that the $\text{GL}_b(\mathbf{Z})$ -set $L_{b,3}$ is contained in the $\text{GL}_b(\mathbf{Z})$ -set L_b . The $\text{GL}_b(\mathbf{Z})$ -action on $L_{b,3}$ originates from a homomorphism $\text{GL}_b(\mathbf{Z}) \rightarrow \text{GL}(L_{b,3}) := \text{Aut}_{\mathbf{Z}}(L_{b,3})$. In particular, any matrix $g \in \text{GL}_b(\mathbf{Z})$ preserves the absolute value of the determinant of any d_3 elements in $L_{b,3}$. This implies, for instance, that $a \cdot e_1^3$ and $b \cdot e_1^3$ cannot lie in the same $\text{GL}_b(\mathbf{Z})$ -orbit, if $0 \leq a < b$. \square

4. PROOF OF THEOREM 2.2 AND THEOREM 2.4

From now on, X stands for an eight-dimensional E-manifold with $w_2(X) = 0$.

4.1 PROOF OF THEOREM 2.2

The classification result for 3-connected E-manifolds of dimension eight is a special case of a result of Wall's [36] and can be easily obtained with the methods described in [17], VII, §12. Let us recall the details, because we will need them later on.

We fix a basis \underline{b} for $H_4(X, \mathbf{Z})$ and let \underline{y} be the dual basis of $H^4(X, \mathbf{Z})$. Then there is a handle presentation $X = D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4 \cup D^8$ with \underline{b} as the preferred basis. The manifold $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$ is determined by the ambient isotopy class of a framed link of 3-spheres in S^7 , having b' components. Let us first look at such a link, forgetting the framing, i.e., suppose we are given embeddings $g_i: S^3 \rightarrow S^7$ with $S_i \cap S_j = \emptyset$ for $i \neq j$, $S_i := g_i(S^3)$, $i = 1, \dots, b'$. By 3.9, we may assume that the g_i are differentiable. Observe that the normal bundles of the S_i are trivial.

We equip S_i with the orientation induced via g_i by the standard orientation of S^3 and the normal bundle of S_i with the orientation which is determined by requiring that the orientation of S_i followed by that of its normal bundle coincide with the orientation of S^7 . Therefore, a 3-sphere F_i which bounds the fibre of a tubular neighborhood of S_i in S^7 inherits an orientation and thus provides a generator e_i for $H_3(S^7 \setminus S_i, \mathbf{Z}) \cong \mathbf{Z}$, $i = 1, \dots, b'$. For $i \neq j$, the image of the fundamental class $[S_i]$ in $H_3(S^7 \setminus S_j, \mathbf{Z})$ is of the form $\lambda_{ij} \cdot e_j$. The integer λ_{ij} is called *the linking number of S_i and S_j* .

For $i = 1, \dots, b'$, the manifold $S^7 \setminus \bigcup_{j \neq i} S_j$ is up to dimension 5 homotopy equivalent to $\bigvee_{j \neq i} F_j$, and

$$\pi_3(S^7 \setminus \bigcup_{j \neq i} S_j) \cong \pi_3(\bigvee_{j \neq i} F_j) \cong \bigoplus_{j \neq i} H_3(S^7 \setminus S_j, \mathbf{Z}).$$

Under this identification, we have $[g_i] = \sum_{j \neq i} \lambda_{ij} \cdot e_j$. The $[g_i]$ determine the ambient isotopy class of the given link (3.9), and we deduce

PROPOSITION 4.1. *The linking numbers λ_{ij} , $1 \leq i < j \leq b'$, determine the given link up to ambient isotopy.*

The sphere S_i bounds a 4-dimensional disc D_i^- in D^8 , $i = 1, \dots, b'$, which we equip with the induced orientation. We may, furthermore, assume

that the D_i^- intersect transversely in the interior of D^8 . Then the λ_{ij} coincide with the intersection numbers $D_i^- \cdot D_j^-$, $1 \leq i < j \leq b'$. For an intuitive proof (in dimension 4), see [28], p.67. Now, every disc D_i^- is completed by the core disc D_i^+ of the i^{th} 4-handle to an embedded 4-sphere Σ_i in T , $i = 1, \dots, b'$, and, since all the core discs are pairwise disjoint, the λ_{ij} coincide with the intersection numbers $\Sigma_i \cdot \Sigma_j$, $1 \leq i < j \leq b'$. Finally, X is obtained by gluing an 8-disc to T along ∂T , and the spheres Σ_i represent the elements of the chosen basis \underline{b} of $H_4(X, \mathbf{Z})$. Identifying the intersection ring with the cohomology ring of X via Poincaré-duality, we see

COROLLARY 4.2. *The linking numbers λ_{ij} coincide with the cup products $(y_i \cup y_j)[X]$, $1 \leq i < j \leq b'$, i.e., the link of the attaching spheres is determined up to ambient isotopy by the basis \underline{b} and the cup products.*

As we have remarked before, the normal bundles of the S_i are trivial, whence there exist embeddings $f_i^0: S^3 \times D^4 \rightarrow S^7$ with $f_i^0|_{S^3 \times \{0\}} = g_i$, $i = 1, \dots, b'$. From the uniqueness of tubular (in differential topology) or regular (in piecewise linear topology) neighbourhoods, every other embedding $f_i: S^3 \times D^4 \rightarrow S^7$ with $f_i|_{S^3 \times \{0\}} = g_i$ is ambient isotopic to one of the form $f_i^{[h_i]} := ((x, y) \mapsto (x, h_i \cdot y))$, $[h_i] \in \pi_3(\text{SO}(4))$, $i = 1, \dots, b'$. Corollary 3.14 implies that we can choose the f_i^0 , $i = 1, \dots, b'$, in such a way that the following holds:

LEMMA 4.3. *Suppose T is obtained by attaching 4-handles along $f_i^{[h_i]}$ with $[h_i] = k_1^i \alpha_3 + k_2^i \beta_3$, $i = 1, \dots, b'$, then*

$$\Sigma_i \cdot \Sigma_i = k_2^i \quad \text{and} \quad p_1(T_{T|\Sigma_i}) = \pm(2k_2^i + 4k_1^i).$$

This shows that also the framed link used for constructing T and X is determined by the system of invariants associated to (X, \underline{y}) , proving the injectivity in Part i) of the theorem. Moreover, the assertion about the fibres in Part ii) is clear.

Conversely, given a system Z of invariants in $Z(0, b')$, satisfying relation (2), there exists a based 3-connected manifold (X, \underline{y}) realizing Z . Indeed, by the above identification of the invariants, Z determines a framed link in S^7 and thus the manifold $T := D^8 \cup H_1^4 \cup \dots \cup H_{b'}^4$. The boundary of T is a 7-dimensional homotopy sphere ([17], (12.2), p.119) and, therefore, piecewise linearly homeomorphic to S^7 . Hence, $X = T \cup_{S^7} D^8$ is a piecewise linear manifold with the desired system of invariants, settling Part i). If, in

addition, relation (3) holds, then [18] ensures that X will carry a smooth structure (compare Theorem A.4 of [24]), finishing the proof of Part ii). \square

4.2 THE DETERMINATION OF W_4 IN THE GENERAL CASE

We have a handle decomposition $W_0 \subset W_2 \subset W_4 \subset W_6 \subset X$ of X providing preferred bases \underline{b} of $H_2(X, \mathbf{Z})$ and \underline{c} of $H_4(X, \mathbf{Z})$, respectively. Let \underline{x} and \underline{y} be the dual bases of $H^2(X, \mathbf{Z})$ and $H^4(X, \mathbf{Z})$, respectively. Finally, let \underline{y}^* be the basis of $H^4(X, \mathbf{Z})$ which is dual to \underline{y} via γ_X .

We find $\partial W_2 \cong \#_{i=1}^b (S^2 \times S^5)$, and W_4 is determined by the ambient isotopy class of a framed link of 3-spheres in ∂W_2 with b' components. Let $f_k: S^3 \times D^4 \rightarrow \partial W_2$ be the k^{th} component of that link and $g_k := f_k|_{S^3 \times \{0\}}$, $k = 1, \dots, b'$. In the notation of Section 3.6, we write $[g_k] \in \pi_3(\partial W_2 \setminus \bigcup_{k \neq j} S_j)$ in the form $(l_i^k, i = 1, \dots, b, l_{ij}^k, 1 \leq i < j \leq b; \lambda_{kj}, j \neq k)$, $k = 1, \dots, b'$. To see the significance of the l_i^k and l_{ij}^k , note that, by Remark 3.4, $W_2 \cup H_k^4 \subset X$ is homotopy equivalent to $(\bigvee_{i=1}^b S^2) \cup_{g_k} D^4$. The cohomology ring of that complex has been computed in Proposition 3.11, so that the naturality of the cup product implies the following formulae for the cup products in X :

$$x_i \cup x_j = \sum_{k=1}^{b'} l_{ij}^k \cdot y_k^*, \quad i \neq j,$$

$$x_i \cup x_i = \sum_{k=1}^{b'} l_i^k \cdot y_k^*, \quad i = 1, \dots, b.$$

Therefore, the l_i^k and l_{ij}^k are determined by δ_X and γ_X (used to compute \underline{y}^*), in fact $l_i^k = \gamma_X(\delta(x_i \otimes x_i) \otimes y_k)$ and $l_{ij}^k = \gamma_X(\delta(x_i \otimes x_j) \otimes y_k)$.

To determine the λ_{ij} and the framings, we proceed as follows: Look at the embedding $\#_{i=1}^b (S^2 \times S^5) \hookrightarrow X$. There exist b embedded 2-spheres S_1^2, \dots, S_b^2 which represent the basis \underline{b} and which do not meet the given link. Finally, $\#_{i=1}^b (S^2 \times S^5)$ obviously possesses a regular neighborhood in X which is homeomorphic to $\#_{i=1}^b (S^2 \times S^5) \times D^1$. Thus, we can perform "surgery in pairs" as described in Section 3.1. The result is a 3-connected manifold X^* containing S^7 . It is by construction the manifold obtained from the framed link in S^7 derived from the given one in $\#_{i=1}^b (S^2 \times S^5)$ (cf. Section 4.1). We will be finished, once we are able to compare the invariants of X to those of X^* . To do so, we look at the *trace of the surgery*, i.e., at $Y = (X \times I) \cup H_1^5 \cup \dots \cup H_{b'}^5$, the 5-handles being attached along tubular neighborhoods of the $S_i \times \{1\}$ in $X \times \{1\}$. Then $\partial Y = X \sqcup \bar{X}^*$.

The Mayer-Vietoris sequence provides the isomorphisms

$$H_4(X, \mathbf{Z}) \cong H_4\left(X \setminus \bigsqcup_{i=1}^{b'} (S_i \times D^6), \mathbf{Z}\right) \cong H_4(X^*, \mathbf{Z}).$$

Set $H := H_4\left(X \setminus \bigsqcup_{i=1}^{b'} (S_i \times D^6), \mathbf{Z}\right)$. By Lefschetz duality ([5], (28.18)), there is for each $q \in \mathbf{N}$ a diagram (omitting \mathbf{Z} -coefficients)

$$(4) \quad \begin{array}{ccccccc} H^{q-1}(Y) & \longrightarrow & H^{q-1}(\partial Y) & \longrightarrow & H^q(Y, \partial Y) & \longrightarrow & H^q(Y) \\ \downarrow \cong & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ H_{10-q}(Y, \partial Y) & \longrightarrow & H_{9-q}(\partial Y) & \longrightarrow & H_{9-q}(Y) & \longrightarrow & H_{9-q}(Y, \partial Y) \end{array}$$

where the left square commutes up to the sign $(-1)^{q-1}$ and the other two commute. We first use it in the case $q = 5$. Look at the commutative diagram

$$\begin{array}{ccc} H & \xrightarrow{\cong} & H_4(X^*, \mathbf{Z}) \\ \downarrow \cong & & \downarrow \\ H_4(X, \mathbf{Z}) & \longrightarrow & H_4(Y, \mathbf{Z}), \end{array}$$

in which all arrows are injective, because $H_5(Y, X; \mathbf{Z}) = 0 = H_5(Y, X^*; \mathbf{Z})$ (cf. [17], p.198). Using the identification $H_4(\partial Y, \mathbf{Z}) = H \oplus H$, we find

$$(5) \quad \text{Im}(H_5(Y, \partial Y; \mathbf{Z})) = \{ (y, -y) \in H \oplus H \}.$$

Similar considerations apply to the case $q = 9$. Taking into account that X^* sits in Y with the reversed orientation, (4) shows that the forms γ_X and γ_{X^*} , both defined with respect to the preferred bases, coincide. In the same manner, the pullbacks of $p_1(Y)$ to $H^4(X, \mathbf{Z})$ and $H^4(X^*, \mathbf{Z})$, respectively, agree. Since X and X^* are the boundary components of Y , these pullbacks are $p_1(X)$ and $p_1(X^*)$, respectively, and we are done. \square

4.3 MANIFOLDS WITH GIVEN INVARIANTS

One might speculate, especially in view of the classification of E-manifolds in dimension 4 and 6, that the invariants δ_X , γ_X , and $p_1(X)$ might suffice to classify E-manifolds with $w_2(X) = 0$ in dimension 8. However,

Lemma 3.6 shows that these invariants determine only W_4 and we still have the choice of an isomorphism in gluing $\#_{i=1}^b(S^2 \times S^5)$ to W_4 , and different gluings may lead to different results. The following example, which was communicated to me by J.-C. Hausmann, illustrates this phenomenon.

EXAMPLE 4.4. One has $\pi_5(\mathrm{SO}(3)) \cong \mathbf{Z}_2$ [32]. Therefore, there are two different S^2 -bundles over S^6 , call them $X := S^6 \times S^2$ and $X' := S^6 \tilde{\times} S^2$. Obviously, X and X' are spin-manifolds with trivial invariants, but one computes $\pi_5(X) \cong \mathbf{Z}_2$ and $\pi_5(X') = \{0\}$.

Fix b, b' , and a system Z of invariants in the image of the map $Z^{\mathrm{PL}(\mathcal{C}^\infty)}(b, b')$. As we have seen, Z determines a certain manifold W_4 whose boundary is diffeomorphic to $\#_{i=1}^b(S^2 \times S^5)$ together with a basis \underline{b} for $H_2(\partial W_4, \mathbf{Z})$. Let \underline{b}_0 be the natural basis for $H_2(\#_{i=1}^b(S^2 \times S^5), \mathbf{Z})$, and denote by $\mathrm{Iso}_0^{\mathrm{PL}(\mathcal{C}^\infty)}$ the set of piecewise linear (smooth) isomorphisms $f: \#_{i=1}^b(S^2 \times S^5) \rightarrow \partial W_4$ with $f_*(\underline{b}_0) = \underline{b}$. Our results show that every based piecewise linear (smooth) manifold $(X, \underline{x}, \underline{y})$ with system of invariants Z is piecewise linearly (smoothly) isomorphic to a manifold of the form

$$X(f) := \partial W_4 \cup_f \#_{i=1}^b(S^2 \times S^5) \quad \text{for some } f \in \mathrm{Iso}_0^{\mathrm{PL}(\mathcal{C}^\infty)}$$

with its given bases for $H^2(X(f), \mathbf{Z})$ and $H^4(X(f), \mathbf{Z})$. Conversely, every manifold of the form $X(f)$ is a piecewise linear (smooth) based E-manifold with invariants Z .

Now, suppose we are given $f, f' \in \mathrm{Iso}_0^{\mathrm{PL}(\mathcal{C}^\infty)}$, such that $X(f)$ and $X(f')$ are isomorphic as piecewise linear (smooth) based manifolds. We claim that we can find an isomorphism $\varphi: X(f) \rightarrow X(f')$ with $\varphi(W_4) = W_4$. For this, look at the handle decomposition $W_0 \subset W_2 \subset W_4$. Since W_0 is just an embedded 8-disc in $X(f)$ and $X(f')$, respectively, we can choose φ with $\varphi(W_0) = W_0$. Let $l \subset \partial W_0$ be the framed link for attaching the 2-handles. Then $\varphi(l)$ and l are isotopic. Therefore, we can find a level preserving diffeomorphism $\tilde{\psi}: [-1, 1] \times \partial W_0 \rightarrow [-1, 1] \times \partial W_0$ with $\tilde{\psi}|_{\{\pm 1\} \times \partial W_0} = \mathrm{id}_{\partial W_0}$ and $\tilde{\psi}|_{\{0\} \times \partial W_0}(\varphi(l)) = l$. If we choose a tubular neighborhood ($\cong [-1, 1] \times \partial W_0$) of ∂W_0 in $X(f')$, we can use $\tilde{\psi}$ to define an automorphism $\psi: X(f') \rightarrow X(f')$ with $\psi(\varphi(l)) = l$. Thus, $\psi \circ \varphi$ maps W_2 onto W_2 . A similar argument shows that we can achieve $\varphi(W_4) = W_4$.

Let $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6))$ be the group of piecewise linear (smooth) automorphisms g of $\#_{i=1}^b(S^2 \times D^6)$ with $H^2(g, \mathbf{Z}) = \text{id}$ and similarly define $\text{Aut}_0^{\text{PL}(C^\infty)}(W_4)$. Then we have just established

PROPOSITION 4.5. *The set of isomorphism classes of based piecewise linear (smooth) E-manifolds with invariants Z is in bijection to the set of equivalence classes in $\text{Iso}_0^{\text{PL}(C^\infty)}$ with respect to the equivalence relation coming from the group action*

$$\text{Aut}_0^{\text{PL}(C^\infty)}(W_4) \times \text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6)) \times \text{Iso}_0^{\text{PL}(C^\infty)} \longrightarrow \text{Iso}_0^{\text{PL}(C^\infty)}$$

$$(h, g, f) \longmapsto h|_{\partial W_4} \circ f \circ g|_{\#_{i=1}^b(S^2 \times S^5)}^{-1}.$$

We shall see in Lemma 5.1 that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ contains the commutator subgroup of $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$.

COROLLARY 4.6. *The set of isomorphism classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the abelian group*

$$\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6)).$$

I have been informed by experts that the structure of the groups $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times S^5))$ and $\text{Aut}_0^{\text{PL}(C^\infty)}(\#_{i=1}^b(S^2 \times D^6))$ has not yet been determined and that this would be a rather difficult task. Therefore, we choose the viewpoint of framed links in order to finish our considerations. In Theorem 5.2, we will then use this viewpoint to compute the group $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$.

As above, let $(X, \underline{x}, \underline{y})$ be a based piecewise linear (smooth) E-manifold with zero second Stiefel-Whitney class and system of invariants $Z_{(X, \underline{x}, \underline{y})} = (\delta, \gamma, p)$. We have seen that we can find a framed link l_X of 2-spheres in X which represents the basis \underline{x} and perform surgery along this link in order to get a 3-connected piecewise linear (smooth) based manifold (X^*, \underline{y}) together with a framed link l_{X^*} of 5-spheres in it. If $(X', \underline{x}', \underline{y}', l_{X'})$ is another such object where $(X', \underline{x}', \underline{y}')$ is isomorphic to $(X, \underline{x}, \underline{y})$, then clearly we can find an isomorphism $\varphi: (X, \underline{x}, \underline{y}) \longrightarrow (X', \underline{x}', \underline{y}')$ with $\varphi(l_X) = l_{X'}$. Such an isomorphism φ yields, after surgery, an isomorphism $\varphi^*: (X^*, \underline{y}) \longrightarrow (X'^*, \underline{y}')$ with $\varphi^*(l_{X^*}) = l_{X'^*}$. In particular, the manifold (X^*, \underline{y}) is determined up to piecewise linear (smooth) isomorphy. We call it the *type of $(X, \underline{x}, \underline{y})$* . Note that this notion matters only in the smooth case, by Theorem 2.2.

To summarize, we have

PROPOSITION 4.7. *The set of isomorphism classes of based piecewise linear (smooth) E-manifolds of type (X^*, \underline{y}) is in bijection to the set of equivalence classes of framed links of 5-spheres in X^* where two such links l and l' are considered equivalent, if there is a piecewise linear (smooth) automorphism $\varphi^* : (X^*, \underline{y}) \rightarrow (X^*, \underline{y})$ with $\varphi^*(l) = l'$.*

EXAMPLE 4.8. The group $\mathbf{Z}_2^{\oplus b}$ acts freely on the set of isotopy classes of framed links of b spheres of dimension 5 in X^* by altering the framings of the components. Note that the two possible framings of the trivial bundle on a 5-sphere are distinguished by the fact that one extends over D^6 and the other does not. This property is preserved under piecewise linear homeomorphisms, so that we conclude that $\mathbf{Z}_2^{\oplus b}$ acts also freely on the set of equivalence classes of framed links of b spheres of dimension 5 in X^* .

Note that this completes the classification of Spin-E-manifolds of dimension eight with second Betti number one.

Let us look at manifolds of type S^8 . We claim that two framed links l and l' of 5-spheres are equivalent in the above sense, if and only if they are isotopic. Clearly, after replacing l and l' by isotopic links, we may assume that both of them are contained in the Southern hemisphere and that φ^* is the identity on the Northern hemisphere. Now, choose a representative φ^\dagger for the isotopy class of φ^{*-1} which is the identity on the Southern hemisphere. Then $\varphi^\dagger \circ \varphi^*$ is isotopic to the identity and carries l into l' .

For differentiable manifolds, the operation $X \mapsto X\#\Sigma$, Σ an exotic 8-sphere, establishes a bijection between the set of isomorphism classes of based smooth E-manifolds of type S^8 and the set of isomorphism classes of based smooth E-manifolds of type Σ . We conclude

COROLLARY 4.9. i) *The set of isomorphism classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the group $\text{FL}_b = \text{L}_b \oplus \bigoplus_{i=1}^b \mathbf{Z}_2$.*

ii) *The set of isomorphism classes of based smooth E-manifolds with $b_2 = b$ and $b_4 = 0$ is in bijection to the group $\mathcal{V}^8 \oplus \text{FL}_b$.*

Finally, we have to deal with those manifolds for which the cup form δ is trivial. Our investigations in Sections 3.6 and 4.2 show that the framed link of 3-spheres in ∂W_2 can be chosen to be contained in a small disc.

In other words, a manifold X with $\delta_X \equiv 0$ is piecewise linear (smoothly) isomorphic $X^\dagger \# X^*$ where X^* is the type of X and $b_4(X^\dagger) = 0$. As our surgery arguments above reveal, an isomorphism between $X^\dagger \# X^*$ and $X'^\dagger \# X'^*$ can be chosen of the form $\varphi^\dagger \# \varphi^*$ where $\varphi^\dagger: X^\dagger \rightarrow X'^\dagger$ and $\varphi^*: X^* \rightarrow X'^*$ are isomorphisms. Therefore, the set of isomorphy classes of based piecewise linear E-manifolds of type X^* with $b_2 = b$ is in bijection to the set of isomorphy classes of based piecewise linear E-manifolds with $b_2 = b$ and $b_4 = 0$. The same goes for differentiable manifolds of type X^* , if X^* is not diffeomorphic to $X^* \# \Sigma$, Σ an exotic 8-sphere. Otherwise, we have to divide by the action of ϑ^8 . This observation together with Corollary 4.9 settles Theorem 2.4. \square

5. STRUCTURE OF THE GROUP $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$

In this section we prove that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is an abelian group which is, moreover, isomorphic to the group FL_b defined before. This result should be of some independent interest, especially because the group FL_b is quite well understood by Haefliger's work. For $b = 1$, we refer to [20] for more specific information.

We begin with the elementary

LEMMA 5.1. *Let $k \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ be a commutator. Then k extends to an automorphism of $\#_{i=1}^b(S^2 \times D^6)$.*

Proof. For the proof, we depict $\#_{i=1}^b(S^2 \times S^5)$ as follows: Let V_i , $i = 1, \dots, b$, be b copies of $S^2 \times D^6$, and we join V_i and V_{i+1} by a tube $T_i \cong [-1, 1] \times D^7$, $i = 1, \dots, b-1$. The result is a manifold W whose boundary is isomorphic to $\#_{i=1}^b(S^2 \times S^5)$. We make the following normalizations: Write ∂V_i as $(S^2 \times D_+^i) \cup (S^2 \times D_-^i)$, let n_i and s_i be the centers of D_+^i and D_-^i , respectively, and set $S_+^i := S^2 \times n_i$ and $S_-^i := S^2 \times s_i$, $i = 1, \dots, b$. Choose furthermore points $e_i \neq w_i$ in $(S^2 \times D_+^i) \cap (S^2 \times D_-^i)$, $i = 1, \dots, b$, and suppose that $\{-1\} \times D^7 \subset T_i$ is attached to a disc around w_i in ∂V_i and $\{1\} \times D^7 \subset T_i$ to a disc around e_{i+1} in ∂V_{i+1} , $i = 1, \dots, b-1$. Set $T := \bigsqcup_{i=1}^{b-1} T_i$.

Now, let $k = f \circ g \circ f^{-1} \circ g^{-1}$ with $f, g \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$. As $H_2(h, \mathbf{Z})$ is the identity for every element $h \in \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$ and S_\pm^i , $i = 1, \dots, b$, both represent the same basis for $H_2(\partial W, \mathbf{Z})$, h is isotopic to a map h' which satisfies either assumption (A) or (B) below.

- (A) : h' is trivial on a tubular neighborhood of S^i_+ which contains $(S^2 \times D^i_+) \setminus \text{Int}(T)$, $i = 1, \dots, b$.
- (B) : h' is trivial on a tubular neighborhood of S^i_- which contains $(S^2 \times D^i_-) \setminus \text{Int}(T)$, $i = 1, \dots, b$.

Next, replace f by an isotopic map f' satisfying (A), and g by an isotopic map g' satisfying (B). Then k' is isotopic to $f' \circ g' \circ f'^{-1} \circ g'^{-1}$. The map k' is the identity outside $\text{Int}(\partial T)$. It is, furthermore, the identity on a collar of $(\{-1\} \sqcup \{1\}) \times S^6$ in $R_i := [-1, 1] \times S^6 \subset \partial T_i$, $i = 1, \dots, b - 1$. Let k'_i be the restriction of k' to R_i , $i = 1, \dots, b$. We know that each k'_i is the identity on a collar of $\{-1, 1\} \times S^6$ in R_i . Thus, we extend every k'_i to a PL automorphism \tilde{k}_i of $D^7 \times \{-1\} \cup R_i \cup D^7 \times \{1\} \cong S^7$ through $\text{id}_{D^7 \times \{-1\} \cup D^7 \times \{1\}}$. Now, by [27], Lemma 1.10, p. 8, \tilde{k}_i extends to an automorphism κ_i of $D^8 \cong D^7 \times [-1, 1]$, $i = 1, \dots, b$. Thus, the maps id_{V_i} and α_i , $i = 1, \dots, b$, glue to an automorphism of $\#_{i=1}^b(S^2 \times D^6)$ whose restriction to the boundary is just k' . \square

This lemma shows that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is a normal subgroup of $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5))$, and that $\text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$ is abelian. Moreover, in Section 4.3, we have already defined a set theoretic bijection

$$\beta: \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6)) \longrightarrow \text{FL}_b .$$

THEOREM 5.2. *The map β is a group isomorphism.*

Proof. Since β is bijective, we have to verify that β is a homomorphism. In order to do so, we will construct a group \mathbf{G} together with surjective homomorphisms

$$\chi_1: \mathbf{G} \longrightarrow \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6))$$

and

$$\chi_2: \mathbf{G} \longrightarrow \text{FL}_b ,$$

such that $\chi_2 = \beta \circ \chi_1$. This will clearly settle the claim.

Before we define \mathbf{G} , we recall some constructions and conventions from [11]. Let $S^8 = \{(x_0, \dots, x_9) \in \mathbf{R}^9 \mid x_0^2 + \dots + x_9^2 = 1\}$ be the unit sphere, write $S^8 = D^8_+ \cup D^8_-$, and let $\sigma: S^8 \longrightarrow S^8$ be the reflection at $S^7 = D^8_+ \cap D^8_-$, interchanging the Northern and the Southern hemispheres. First, let $S_b := (S^5_1, \dots, S^5_b)$ be a ‘standard link’ in S^8 defined as follows: Fix real numbers $-1/2 < a_1 < \dots < a_b < 1/2$, and set

$$S_i^5 := \{ (x_0, \dots, x_9) \in S^8 \mid x_6 = x_7 = x_8 = 0, x_9 = a_i \}.$$

We choose, furthermore, framings $\tau_i: S_i^5 \times D^3 \rightarrow S^8$ which extend over D^6 , such that $\tau_i(D_{i,\pm}^5 \times D^3) \subset D_{\pm}^8$ and $\sigma \circ \tau_i = \tau_i \circ (\sigma|_{S_i^5} \times \text{id}_{D^3})$, $i = 1, \dots, b$. Let l_b^0 be the resulting framed link in S^8 with $l_{b,\pm}^0 := l_b^0 \cap D_{\pm}^8$. Recall from Section 1 of [11] that

1. Every framed link l of b five-dimensional spheres in S^8 is isotopic to a link l' , such that either (A) $l' \cap D_+^8 = l_{b,+}^0$ or (B) $l' \cap D_-^8 = l_{b,-}^0$.
2. If l_1 satisfies (A) and l_2 satisfies (B), then $l_1 + l_2$ is represented by the link l with $l \cap D_+^8 = l_1 \cap D_+^8$ and $l \cap D_-^8 = l_2 \cap D_-^8$.

Note that, if we perform surgery along l_b^0 , we get a manifold $W = W_+ \cup W_-$ which is isomorphic to $\#_{i=1}^b(S^2 \times S^6)$, and

$$W_{\pm} := (D_{\pm}^8 \setminus \text{Int}(l_b^0)) \cup \left(\bigsqcup_{i=1}^b (S_i^2 \times D_{\pm}^6) \right)$$

is canonically isomorphic to $\#_{i=1}^b(S^2 \times D^6)$. For the rest of the proof, we will use the description of $\#_{i=1}^b(S^2 \times S^5)$ as $\partial W_+ = \partial W_-$. Set

$$\mathbf{G} := \{ \text{PL-maps } f: S^7 \setminus \text{Int}(l_b^0) \rightarrow S^7 \setminus \text{Int}(l_b^0): f|_{\text{boundary}} = \text{id} \}.$$

For every $f \in \mathbf{G}$, we define $\varphi(f): \#_{i=1}^b(S^2 \times S^5) \rightarrow \#_{i=1}^b(S^2 \times S^5)$, by extending f through the identity on $\bigsqcup_{i=1}^b(S_i^2 \times D^5)$. Similarly, define $\psi(f): S^7 \rightarrow S^7$. Obviously,

$$\begin{aligned} \chi_1: \mathbf{G} &\longrightarrow \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times S^5)) / \text{Aut}_0^{\text{PL}}(\#_{i=1}^b(S^2 \times D^6)) \\ f &\longmapsto [\varphi(f)] \end{aligned}$$

is a surjective homomorphism.

Next, we associate to $f \in \mathbf{G}$ an element $\chi_2(f) \in \text{FL}_b$ as follows: First, we define $\Sigma(f) := D_+^8 \cup_{\psi(f)} D_-^8$ and the link $l'(f) := l_{b,+}^0 \cup_{\psi(f)} l_{b,-}^0$. Then we choose a piecewise linear homeomorphism $F: \Sigma(f) \rightarrow S^8$ and set $l_F(f) := F(l'(f))$. We have checked before that the isotopy class of $l_F(f)$ does not depend on the chosen homeomorphism, so that $\chi_2(f) := [l_F(f)] \in \text{FL}_b$ is well defined. To see that $\chi_2: \mathbf{G} \rightarrow \text{FL}_b$ is a homomorphism, let f, f' be in \mathbf{G} . Choose extensions $\bar{\psi}: D_+^8 \rightarrow D_+^8$ and $\bar{\psi}': D_-^8 \rightarrow D_-^8$ of $\psi(f)$ and $\psi(f')$, respectively. We then define $F: \Sigma(f) \rightarrow S^8$ as $\bar{\psi}$ on D_+^8 and as the identity on D_-^8 , $F': \Sigma(f) \rightarrow S^8$ as the identity on D_+^8 and $(\bar{\psi}')^{-1}$ on D_-^8 , and $F'': \Sigma(f' \circ f) \rightarrow S^8$ as $\bar{\psi}$ on D_+^8 and $(\bar{\psi}')^{-1}$ on D_-^8 . Then the link $l_F(f)$ satisfies (B), the link $l_{F'}(f')$ satisfies (A), and (2) above shows that $[l_{F''}(f' \circ f)] = [l_{F'}(f')] + [l_F(f)]$.

Finally, for given $f \in \mathbf{G}$, we can perform surgery on $\Sigma(f)$ along $l'(f)$. The result is $W_+ \cup_{\varphi(f)} W_-$. Reading this backwards means nothing else but $\beta(\chi_1(f)) = \chi_2(f)$ and we are done. \square

REFERENCES

- [1] BROWDER, W. *Surgery on Simply-Connected Manifolds*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer, New York–Heidelberg, 1972.
- [2] CAMPANA, F. and TH. PETERNELL. Rigidity theorems for primitive Fano 3-folds. *Comm. Anal. Geom.* 2 (1994), 173–201.
- [3] FREITAG, P. Rigidity properties of imprimitive Fano 3-folds. *Math. Ann.* 309 (1997), 179–198.
- [4] FULTON, W. *Introduction to Toric Varieties*. Annals of Mathematics Studies 131. The William H. Roever Lectures in Geometry, Princeton University Press, Princeton (NJ), 1993.
- [5] GREENBERG, M. *Lectures on Algebraic Topology*. W. A. Benjamin, New York – Amsterdam, 1967.
- [6] HAEFLIGER, A. Differentiable imbeddings. *Bull. Amer. Math. Soc.* 67 (1961), 109–112.
- [7] — Plongements différentiables de variétés dans variétés. *Comment. Math. Helv.* 36 (1961), 47–82.
- [8] — Differentiable links. *Topology* 1 (1962), 241–244.
- [9] — Knotted spheres and related geometric problems. *Proc. Internat. Congr. Math. (Moscow, 1966)*, 437–445.
- [10] — Differential embeddings of S^n in S^{n+q} for $q > 2$. *Ann. of Math. (2)* 83 (1966), 402–436.
- [11] — Enlacements de sphères en codimension supérieure à 2. *Comment. Math. Helv.* 41 (1966/1967), 51–72.
- [12] HAUSMANN, J.-C. and A. KNUTSON. Polygon spaces and Grassmannians. *L'Enseignement Math.* 43 (1997), 173–198.
- [13] HUDSON, J. F. P. *Piecewise Linear Topology*. University of Chicago Lecture Notes, prepared with the assistance of J. L. Shaneson and J. Lees. W. A. Benjamin, Inc., New York–Amsterdam, 1969.
- [14] JUPP, P. Classification of certain 6-manifolds. *Proc. Cambridge Philos. Soc.* 73 (1973), 293–300.
- [15] KIRBY, R. *The Topology of 4-Manifolds*. Lecture Notes in Mathematics, 1374. Springer-Verlag, Berlin, 1989.
- [16] KIRBY, R. and L. SIEBENMANN. *Foundational Essays on Topological Manifolds, Smoothings, and Triangulations*, with notes by John Milnor and Michael Atiyah. Annals of Mathematics Studies, Princeton University Press, Princeton – University of Tokyo Press, Tokyo, 1977.
- [17] KOSINSKI, A. *Differential Manifolds*. Academic Press, Boston, 1993.
- [18] KUIPER, N. C. Algebraic equations for nonsmoothable 8-manifolds. *Inst. Hautes Études Sci. Publ. Math.* 33 (1967), 139–155.
- [19] LEVINE, J. A classification of differentiable knots. *Ann. of Math. (2)* 82 (1965), 15–50.
- [20] — Self-equivalences of $S^n \times S^k$. *Trans. Amer. Math. Soc.* 143 (1969), 523–543.
- [21] MILNOR, J. A procedure for killing homotopy groups of differentiable manifolds. *Proc. Sympos. Pure Math.* 3 (1963), 39–55.

- [22] MILNOR, J. and D. HUSEMOLLER. *Symmetric Bilinear Forms*. Springer, New York – Heidelberg, 1973.
- [23] MILNOR, J. and J.D. STASHEFF. *Characteristic Classes*. Princeton University Press, Princeton (N.J.) – University of Tokyo Press, Tokyo, 1974.
- [24] MÜLLER, S. *Zur Topologie einfacher 8-Mannigfaltigkeiten*. Zürich, 1998.
- [25] NOVIKOV, S. Topological invariance of rational Pontrjagin classes. *Soviet Math. Doklady* 6 (1965), 921–923.
- [26] OKONEK, CH. and A. VAN DE VEN. Cubic forms and complex 3-folds. *L'Enseignement Math.* (2) 41 (1995), 297–333.
- [27] ROURKE, C.P. and B.J. SANDERSON. *Introduction to Piecewise-Linear Topology*. Springer, New York – Heidelberg, 1972.
- [28] SAVELIEV, N. *Lectures on the Topology of 3-Manifolds: An Introduction to the Casson Invariant*. Walter de Gruyter, Berlin, 1999.
- [29] SCHMITT, A. Topological rigidity of holomorphic \mathbf{P}_3 -bundles over curves. *Ann. Math. Pura Appl.* (to appear).
- [30] SELICK, P. *Introduction to Homotopy Theory*. Amer. Math. Soc., Providence (RI), 1997.
- [31] SMALE, S. On the structure of manifolds. *Amer. J. Math.* 84 (1962), 387–399.
- [32] STEENROD, N. *The Topology of Fibre Bundles*. Princeton University Press, Princeton (N.J.), 1951.
- [33] STEER, B. Generalized Whitehead products. *Quart. J. Math. Oxford Ser. (2)* 14 (1963), 29–40.
- [34] SULLIVAN, D. Inside and outside manifolds. In: *Proc. International Congress of Mathematicians (Vancouver, B.C., 1974)*, Vol. 1, pp.201–207. Canad. Math. Congress, Montréal, 1975.
- [35] TODA, H. *Composition Methods in Homotopy Groups of Spheres*. Annals of Mathematics Studies, No. 49. Princeton University Press, Princeton (N.J.), 1962.
- [36] WALL, C. T. C. Classification of $(n-1)$ -connected $2n$ -manifolds. *Ann. of Math.* (2) 75 (1962), 163–189.
- [37] ——— Classification problems in differential topology. V. On certain 6-manifolds. *Invent. Math.* 1 (1966), 355–374.
- [38] WALLACE, A. H. Modifications and cobounding manifolds. *Canad. J. Math.* 12 (1960), 503–528.
- [39] ZEEMAN, E. C. Unknotting combinatorial balls. *Ann. of Math.* (2) 78 (1963), 501–526.

(Reçu le 18 janvier 2002)

Alexander Schmitt

Universität GH Essen

FB6 Mathematik & Informatik

D-45141 Essen

Germany

e-mail: alexander.schmitt@uni-essen.de

Vide-leer-empty